

GANGA GUIDE

MATHEMATICS

10

Authors

P. CHOCKALINGAM, M.Sc., B.Ed.,
Palayamkottai.

D. SAMUEL DEVARAJ,
M.Sc., B.Ed., M.Phil.,
Chennai.

M. ARUL SELVAN, M.Sc., B.Ed., Ph.D.,
Tiruvannamalai.

B. VENKATESH, M.Sc., B.Ed.,
Madurai.



SRI GANGA PUBLICATIONS®

(A UNIT OF SHYAMALA GROUP)

Corporate Office : ↑ Registered Office :

No. 1, Sugar Mill Colony,
Salai Kumaran illam,
Madurai Road,
Tirunelveli - 627001.

Phone : 0462 - 233 8899 / 233 8484
Mobile : 94431 58484 / 95978 39822
Email : suryaguides@yahoo.com

New No. 59, 4th Avenue,
Opp. to Govt. Boys Hr. Sec. School,
Ashok Nagar,
Chennai - 600 083.

Phone : 044 - 2474 4484
Mobile : 94421 58484 / 94425 58484
Email : srigangapublications5@gmail.com

Website : www.suryapublications.in

Price : ₹ 250/-

(ii)

Published By
B. ARUMUGAM
SRI GANGA PUBLICATIONS
(A unit of **Shyamala** Group)

BANK DETAILS

TIRUNELVELI ACCOUNT

Account Name : **Surya Publications**
Account Number: **446971431**
Bank Name : **Indian Bank**
IFSC Code : **IDIB000T034**
Branch Name : **Tirunelveli Junction**

Account Name : **Surya Publications**
Account Number: **510909010051752**
Bank Name : **City Union Bank**
IFSC Code : **CIUB0000230**
Branch Name : **Palayamkottai**

CHENNAI ACCOUNT

Account Name : **Sri Ganga Publication**
Account Number: **928507483**
Bank Name : **Indian Bank**
IFSC Code : **IDIB000A031**
Branch Name : **Ashok Nagar**

Account Name : **Sri Ganga Publication**
Account Number : **512020010022514**
Bank Name : **City Union Bank**
IFSC Code : **CIUB0000230**
Branch Name : **Palayamkottai**

FOREWORD

This **GANGA GUIDE** for **Mathematics** for **Class X** has been prepared written in the hope of sharing the excitement experienced in the study of Mathematics strictly in accordance with the text book published by our Government of TamilNadu in 2019.

We have come forward to facilitate learning maths in a more meaningful and pleasant manner.

The most salient features of this comprehensive guide.

beginning of every topic as “KEY POINTS”.

- ♦ **All the textual example problems are given for all the 8 Units.**
- ♦ The exercise problems have been solved with finest solutions illustrated so charmingly by the author of this Guide.
- ♦ The Guide contains newly-designed problems (15 to 20% will be asked in the Public Exam) in the form of Objective type, Short and Long answer types.

Every effort has been taken to learn mathematics tension-free. The Ganga Guide provides exam-oriented learning inputs that will go a long way in assisting the students to come out successful in the public exam.

With the **Ganga Guide** for **Mathematics X** we wish the students and the teachers a very happy, pleasant, fruitful academic year.

-Publisher

10**MATHEMATICS****CONTENTS**

UNIT	TOPIC	PAGE NO.
1	Relations and Functions	1 - 35
2	Numbers and Sequences	36 - 89
	Algebra	90 - 198
4	Geometry	199 - 249
5	Coordinate Geometry	250 - 293
6	Trigonometry	294 - 335
7	Mensuration	336 - 370
8	Statistics and Probability	371 - 415

CHAPTER 1

RELATIONS AND FUNCTIONS

I. ORDERED PAIR AND CARTESIAN PRODUCT :

Key Points

- ✓ If A and B are two non-empty sets, then the set of all ordered pairs (a, b) such that $a \in A, b \in B$ is called the Cartesian Product of A and B, and is denoted by $A \times B$. Thus, $A \times B = \{(a, b) \mid a \in A, b \in B\}$.
- ✓ $A \times B$ is the set of all possible ordered pairs between the elements of A and B such that the first coordinate is an element of A and the second coordinate is an element of B.
- ✓ $B \times A$ is the set of all possible ordered pairs between the elements of A and B such that the first coordinate is an element of B and the second coordinate is an element of A.
- ✓ If $a = b$ then $(a, b) = (b, a)$.
- ✓ In general $A \times B \neq B \times A$, but $n(A \times B) = n(B \times A)$.
- ✓ $A \times B = \phi$ if and only if $A = \phi$ or $B = \phi$.
- ✓ If $n(A) = p$ and $n(B) = q$ then $n(A \times B) = pq$.
- ✓ For any three sets A, B, C we have
 - (i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$
 - (ii) $A \times (B \cap C) = (A \times B) \cap (A \times C)$.

Example 1.1

If $A = \{1, 3, 5\}$ and $B = \{2, 3\}$ then (i) find $A \times B$ and $B \times A$. (ii) Is $A \times B = B \times A$? If not why? (iii) Show that $n(A \times B) = n(B \times A) = n(A) \times n(B)$.

Solution :

Given that $A = \{1, 3, 5\}$ and $B = \{2, 3\}$

- (i) $A \times B = \{1, 3, 5\} \times \{2, 3\}$
 $= \{(1, 2), (1, 3), (3, 2), (3, 3), (5, 2), (5, 3)\}$
 ... (1)

$$B \times A = \{2, 3\} \times \{1, 3, 5\}$$

$$= \{(2, 1), (2, 3), (2, 5), (3, 1), (3, 3), (3, 5)\}$$

.. (2)

- (ii) From (1) and (2) we conclude that $A \times B \neq B \times A$ as $(1, 2) \neq (2, 1)$ and $(1, 3) \neq (3, 1)$, etc.
- (iii) $n(A) = 3; n(B) = 2$.

From (1) and (2) we observe that,

$$n(A \times B) = n(B \times A) = 6;$$

we see that, $n(A) \times n(B) = 3 \times 2 = 6$ and

$$n(B) \times n(A) = 2 \times 3 = 6$$

Hence, $n(A \times B) = n(B \times A) = n(A) \times n(B) = 6$.

Thus, $n(A \times B) = n(B \times A) = n(A) \times n(B)$.

Example 1.2

If $A \times B = \{(3, 2), (3, 4), (5, 2), (5, 4)\}$ then find A and B .

Solution :

$$A \times B = \{(3, 2), (3, 4), (5, 2), (5, 4)\}$$

We have

$A = \{\text{set of all first coordinates of elements of } A \times B\}$. Therefore $A = \{3, 5\}$

$B = \{\text{set of all second coordinates of elements of } A \times B\}$. Therefore $B = \{2, 4\}$

Example 1.3

Let $A = \{x \in \mathbb{N} \mid 1 < x < 4\}$, $B = \{x \in \mathbb{W} \mid 0 \leq x < 2\}$ and $C = \{x \in \mathbb{N} \mid x < 3\}$.

Then verify that (i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$

(ii) $A \times (B \cap C) = (A \times B) \cap (A \times C)$

Solution :

$$A = \{x \in \mathbb{N} \mid 1 < x < 4\} = \{2, 3\},$$

$$B = \{x \in \mathbb{W} \mid 0 \leq x < 2\} = \{0, 1\}.$$

$$C = \{x \in \mathbb{N} \mid x < 3\} = \{1, 2\}$$

$$(i) \quad A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$B \cup C = \{0, 1\} \cup \{1, 2\} = \{0, 1, 2\}$$

$$A \times (B \cup C) = \{2, 3\} \times \{0, 1, 2\}$$

$$= \{(2, 0), (2, 1), (2, 2), (3, 0), (3, 1), (3, 2)\} \quad \dots (1)$$

$$A \times B = \{2, 3\} \times \{0, 1\}$$

$$= \{(2, 0), (2, 1), (3, 0), (3, 1)\}$$

$$A \times C = \{2, 3\} \times \{1, 2\}$$

$$= \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$(A \times B) \cup (A \times C) = \{(2, 0), (2, 1), (3, 0), (3, 1)\} \cup \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$= \{(2, 0), (2, 1), (2, 2), (3, 0), (3, 1), (3, 2)\} \quad \dots (2)$$

From (1) and (2),

$A \times (B \cup C) = (A \times B) \cup (A \times C)$ is verified.

$$(ii) \quad A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$B \cap C = \{0, 1\} \cap \{1, 2\} = \{1\}$$

$$A \times (B \cap C) = \{2, 3\} \times \{1\}$$

$$= \{(2, 1), (3, 1)\} \quad \dots (3)$$

$$A \times B = \{2, 3\} \times \{0, 1\}$$

$$= \{(2, 0), (2, 1), (3, 0), (3, 1)\}$$

$$A \times C = \{2, 3\} \times \{1, 2\}$$

$$= \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$(A \times B) \cap (A \times C)$$

$$= \{(2, 0), (2, 1), (3, 0), (3, 1)\} \cap \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$= \{(2, 1), (3, 1)\} \quad \dots (4)$$

From (3) and (4),

$A \times (B \cap C) = (A \times B) \cap (A \times C)$ is verified.

EXERCISE 1.1

1. Find $A \times B$, $A \times A$ and $B \times A$

(i) $A = \{2, -2, 3\}$ and $B = \{1, -4\}$

(ii) $A = B = \{p, q\}$ (iii) $A = \{m, n\}$; $B = \phi$

Solution:

(i) Given $A = \{2, -2, 3\}$, $B = \{1, -4\}$.

$$A \times B = \{(2, 1), (2, -4), (-2, 1), (-2, -4), (3, 1), (3, -4)\}$$

$$A \times A = \{(2, 2), (2, -2), (2, 3), (-2, 2), (-2, -2), (-2, 3), (3, 2), (3, -2), (3, 3)\}$$

$$B \times A = \{(1, 2), (1, -2), (1, 3), (-4, 2), (-4, -2), (-4, 3)\}$$

(ii) Given $A = B = \{p, q\}$

$$A \times B = \{(p, p), (p, q), (q, p), (q, q)\}$$

$$A \times A = \{(p, p), (p, q), (q, p), (q, q)\}$$

$$B \times A = \{(p, p), (p, q), (q, p), (q, q)\}$$

(iii) $A = \{m, n\}, B = \phi$

If $A = \phi$ (or) $B = \phi$, then $A \times B = \phi$.

and $B \times A = \phi$

$A \times B = \phi$ and $B \times A = \phi$

$$A \times A = \{(m, m), (m, n), (n, m), (n, n)\}$$

2. Let $A = \{1, 2, 3\}$ and $B = \{x \mid x \text{ is a prime number less than } 10\}$. Find $A \times B$ and

Solution :

Given $A = \{1, 2, 3\}, B = \{x \mid x \text{ is a prime number less than } 10\}$.

$$B = \{2, 3, 5, 7\}$$

$$A \times B = \{(1, 2), (1, 3), (1, 5), (1, 7), (2, 2), (2, 3), (2, 5), (2, 7), (3, 2), (3, 3), (3, 5), (3, 7)\}$$

$$B \times A = \{(2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3), (5, 1), (5, 2), (5, 3), (7, 1), (7, 2), (7, 3)\}$$

3. If $B \times A = \{(-2, 3), (-2, 4), (0, 3), (0, 4), (3, 3), (3, 4)\}$ find A and B .

Solution :

Given $B \times A = \{(-2, 3), (-2, 4), (0, 3), (0, 4), (3, 3), (3, 4)\}$

$$\therefore B = \{-2, 0, 3\}, A = \{3, 4\}$$

4. If $A = \{5, 6\}, B = \{4, 5, 6\}, C = \{5, 6, 7\}$. Show that $A \times A = (B \times B) \cap (C \times C)$.

Solution :

Given $A = \{5, 6\}, B = \{4, 5, 6\}, C = \{5, 6, 7\}$

$$\text{LHS : } A \times A = \{5, 6\} \times \{5, 6\}$$

$$= \{(5, 5), (5, 6), (6, 5), (6, 6)\} \dots (1)$$

$$\text{RHS : } B \times B = \{4, 5, 6\} \times \{4, 5, 6\}$$

$$= \{(4, 4), (4, 5), (4, 6), (5, 4), (5, 5), (5, 6), (6, 4), (6, 5), (6, 6)\}$$

$$C \times C = \{5, 6, 7\} \times \{5, 6, 7\}$$

$$= \{(5, 5), (5, 6), (5, 7), (6, 5), (6, 6), (6, 7), (7, 5), (7, 6), (7, 7)\}$$

$$\therefore (B \times B) \cap (C \times C) = \{(5, 5), (5, 6), (6, 5), (6, 6)\} \dots (2)$$

\therefore From (1) and (2).

$$\text{LHS} = \text{RHS}$$

5. Given $A = \{1, 2, 3\}, B = \{2, 3, 5\}, C = \{3, 4\}$ and $D = \{1, 3, 5\}$, check if $(A \cap C) \times (B \cap D) = (A \times B) \cap (C \times D)$ is true ?

Solution :

Given $A = \{1, 2, 3\}, B = \{2, 3, 5\},$

$$C = \{3, 4\}, D = \{1, 3, 5\}$$

$$A \cap C = \{3\}, B \cap D = \{3, 5\}$$

$$\therefore (A \cap C) \times (B \cap D) = \{(3, 3), (3, 5)\} \dots (1)$$

$$A \times B = \{(1, 2), (1, 3), (1, 5), (2, 2), (2, 3), (2, 5), (3, 2), (3, 3), (3, 5)\}$$

$$C \times D = \{(3, 1), (3, 3), (3, 5), (4, 1), (4, 3), (4, 5)\}$$

$$\therefore (A \times B) \cap (C \times D) = \{(3, 3), (3, 5)\} \dots (2)$$

\therefore From (1) and (2)

$$\text{LHS} = \text{RHS}.$$

6. Let $A = \{x \in W \mid x < 2\}$, $B = \{x \in N \mid 1 < x \leq 4\}$ and $C = \{3, 5\}$. Verify that

(i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$

(ii) $A \times (B \cap C) = (A \times B) \cap (A \times C)$

(iii) $(A \cup B) \times C = (A \times C) \cup (B \times C)$

Solution :

Given $A = \{x \in W \mid x < 2\} \Rightarrow A = \{0, 1\}$

$B = \{x \in N \mid 1 < x \leq 4\}$

$\Rightarrow B = \{2, 3, 4\}$

$C = \{3, 5\}$

(i) To verify :

$A \times (B \cup C) = (A \times B) \cup (A \times C)$

$B \cup C = \{2, 3, 4, 5\}$

$\therefore A \times (B \cup C) = \{(0, 2), (0, 3), (0, 4), (0, 5),$

$A \times B = \{(0, 2), (0, 3), (0, 4), (1, 2),$
 $(1, 3), (1, 4)\}$

$A \times C = \{(0, 3), (0, 5), (1, 3), (1, 5)\}$

$\therefore (A \times B) \cup (A \times C) = \{(0, 2), (0, 3), (0, 4),$
 $(0, 5), (1, 2), (1, 3), (1, 4), (1, 5)\} \dots(2)$

\therefore From (1) and (2) LHS = RHS.

(ii) To verify : $A \times (B \cap C) = (A \times B) \cap (A \times C)$

$B \cap C = \{3\}$

$\therefore A \times (B \cap C) = \{(0, 3), (1, 3)\} \dots(1)$

$A \times B = \{(0, 2), (0, 3), (0, 4), (1, 2),$
 $(1, 3), (1, 4)\}$

$A \times C = \{(0, 3), (0, 5), (1, 3), (1, 5)\}$

$\therefore (A \times B) \cap (A \times C) = \{(0, 3), (1, 3)\} \dots(2)$

\therefore From (1) and (2), LHS = RHS.

(iii) $(A \cup B) \times C = (A \times C) \cup (B \times C)$

$A \cup B = \{0, 1, 2, 3, 4\}$

$\therefore (A \cup B) \times C = \{(0, 3), (0, 5), (1, 3), (1, 5),$
 $(2, 3), (2, 5), (3, 3), (3, 5),$
 $(4, 3), (4, 5)\} \dots(1)$

$A \times C = \{(0, 3), (0, 5), (1, 3), (1, 5)\}$

$B \times C = \{(2, 3), (2, 5), (3, 3), (3, 5),$
 $(4, 3), (4, 5)\}$

$\therefore (A \times C) \cup (B \times C) = \{(0, 3), (0, 5), (1, 3),$
 $(1, 5), (2, 3), (2, 5)$
 $(3, 3), (3, 5), (4, 3),$
 $(4, 5)\} \dots(2)$

\therefore From (1) and (2) LHS = RHS.

7. Let $A =$ The set of all natural numbers less than 8, $B =$ The set of all prime num-

number. Verify that

(i) $(A \cap B) \times C = (A \times C) \cap (B \times C)$

(ii) $A \times (B - C) = (A \times B) - (A \times C)$

Solution :

Given $A = \{1, 2, 3, 4, 5, 6, 7\}$

$B = \{1, 3, 5, 7\}$

$C = \{2\}$

(i) To verify : $(A \cap B) \times C = (A \times C) \cap (B \times C)$

$A \cap B = \{1, 3, 5, 7\}$

$\therefore (A \cap B) \times C = \{(1, 2), (3, 2), (5, 2), (7, 2)\}$
 $\dots(1)$

$A \times C = \{(1, 2), (2, 2), (3, 2), (4, 2), (5, 2),$
 $(6, 2), (7, 2)\}$

$B \times C = \{(1, 2), (3, 2), (5, 2), (7, 2)\} \dots(2)$

\therefore From (1) and (2), LHS = RHS.

(ii) To verify : $A \times (B - C) = (A \times B) - (A \times C)$

$$B - C = \{1, 3, 5, 7\}$$

$$\therefore A \times (B - C) = \{(1, 1), (1, 3), (1, 5), (1, 7),$$

$$(2, 1), (2, 3), (2, 5), (2, 7),$$

$$(3, 1), (3, 3), (3, 5), (3, 7),$$

$$(4, 1), (4, 3), (4, 5), (4, 7),$$

$$(5, 1), (5, 3), (5, 5), (5, 7),$$

$$(6, 1), (6, 3), (6, 5), (6, 7),$$

$$(7, 1), (7, 3), (7, 5), (7, 7),\}$$

$$A \times B = \{(1, 1), (1, 3), (1, 5), (1, 7),$$

$$(2, 1), (2, 3), (2, 5), (2, 7),$$

$$(3, 1), (3, 3), (3, 5), (3, 7),$$

$$(4, 1), (4, 3), (4, 5), (4, 7),$$

$$(5, 1), (5, 3), (5, 5), (5, 7),$$

$$(6, 1), (6, 3), (6, 5), (6, 7),$$

$$(7, 1), (7, 3), (7, 5), (7, 7),\}$$

$$A \times C = \{(1, 2), (2, 2), (3, 2), (4, 2), (5, 2), (6, 2), (7, 2)\}$$

$$\therefore (A \times B) - (A \times C)$$

$$= \{(1, 1), (1, 3), (1, 5), (1, 7),$$

$$(2, 1), (2, 3), (2, 5), (2, 7),$$

$$(3, 1), (3, 3), (3, 5), (3, 7),$$

$$(4, 1), (4, 3), (4, 5), (4, 7),$$

$$(5, 1), (5, 3), (5, 5), (5, 7),$$

$$(6, 1), (6, 3), (6, 5), (6, 7),$$

$$(7, 1), (7, 3), (7, 5), (7, 7),\}$$

\therefore From (1) and (2), LHS = RHS.

II. RELATIONS :

Key Points

- ✓ Let A and B be any two non-empty sets. A 'relation' R from A to B is a subset of $A \times B$ satisfying some specified conditions. If $x \in A$ is related to $y \in B$ through R, then we write it as xRy . xRy if and only if $(x, y) \in R$.
- ✓ A relation may be represented algebraically either by the roster method or by the set builder method.
- ✓ An arrow diagram is a visual representation of a relation.
- ✓ A relation which contains no element is called a "Null relation".
- ✓ If $n(A) = p$, $n(B) = q$, then the total number of relations that exist between A and B is 2^{pq} .

Example 1.4

Let $A = \{3, 4, 7, 8\}$ and $B = \{1, 7, 10\}$. Which of the following sets are relations from A to B ?

(i) $R_1 = \{(3, 7), (4, 7), (7, 10), (8, 1)\}$

(ii) $R_2 = \{(3, 1), (4, 12)\}$

(iii) $R_3 = \{(3, 7), (4, 10), (7, 7), (7, 8), (8, 11), (8, 7), (8, 10)\}$

Solution :

$$A \times B = \{(3, 1), (3, 7), (3, 10), (4, 1), (4, 7), (4, 10), (7, 1), (7, 7), (7, 10), (8, 1), (8, 7), (8, 10)\}$$

(i) We note that $R_1 \subseteq A \times B$. Thus, R_1 is a relation from A to B.


(ii) Here, $(4, 12) \in R_2$, but $(4, 12) \notin A \times B$. So, R_2 is not a relation from A to B.

(iii) Here, $(7, 8) \in R_3$, but $(7, 8) \notin A \times B$. So R_3 is not a relation from A to B.

Example 1.5

The arrow diagram shows a relationship between the sets P and Q. Write the relation in (i) Set builder form (ii) Roster form (iii) What is the domain and range of R.

Solution :



(i) Set builder form of

$$R = \{(x, y) \mid y = x - 2, x \in P, y \in Q\}$$

(ii) Roster form $R = \{(5, 3), (6, 4), (7, 5)\}$

(iii) Domain of $R = \{5, 6, 7\}$ and range of $R = \{3, 4, 5\}$

EXERCISE 1.2

1. Let $A = \{1, 2, 3, 7\}$ and $B = \{3, 0, -1, 7\}$, which of the following are relation from A to B ?

(i) $R_1 = \{(2, 1), (7, 1)\}$

(ii) $R_2 = \{(-1, 1)\}$

(iii) $R_3 = \{(2, -1), (7, 7), (1, 3)\}$

(iv) $R_4 = \{(7, -1), (0, 3), (3, 3), (0, 7)\}$

Solution :

Given $A = \{1, 2, 3, 7\}$, $B = \{3, 0, -1, 7\}$

$$\begin{aligned} \therefore A \times B = & \{(1, 3), (1, 0), (1, -1), (1, 7), \\ & (2, 3), (2, 0), (2, -1), (2, 7), \\ & (3, 3), (3, 0), (3, -1), (3, 7), \\ & (7, 3), (7, 0), (7, -1), (7, 7)\} \end{aligned}$$

i) $R_1 = \{(2, 1), (7, 1)\}$

$(2, 1) \in R_1$ but $(2, 1) \notin A \times B$

$\therefore R_1$ is not a relation from A to B.

ii) $R_2 = \{(-1, 1)\}$

$(-1, 1) \in R_2$ but $(-1, 1) \notin A \times B$

$\therefore R_2$ is not a relation from A to B.

iii) $R_3 = \{(2, -1), (7, 7), (1, 3)\}$

We note that $R_3 \subseteq A \times B$

$\therefore R_3$ is a relation.

iv) $R_4 = \{(7, -1), (0, 3), (3, 3), (0, 7)\}$

$(0, 3), (0, 7) \in R_4$ but not in $A \times B$.

$\therefore R_4$ is not a relation.

2. Let $A = \{1, 2, 3, 4, \dots, 45\}$ and R be the relation defined as “is square of” on A. Write R as a subset of $A \times A$. Also, find the domain and range of R.

Solution :

Given $A = \{1, 2, 3, 4, \dots, 45\}$

R : “is square of”

$R = \{1, 4, 9, 16, 25, 36\}$

Clearly R is a subset of A.

$\therefore \text{Domain} = \{1, 2, 3, 4, 5, 6\}$

$\therefore \text{Range} = \{1, 4, 9, 16, 25, 36\}$

3. A Relation R is given by the set $\{(x, y) / y = x + 3, x \in \{0, 1, 2, 3, 4, 5\}\}$. Determine its domain and range.

Solution :

$$\text{Given } R = \{(x, y) / y = x + 3,$$

$$x \in \{0, 1, 2, 3, 4, 5\}\}$$

$$x = 0 \Rightarrow y = 3$$

$$x = 1 \Rightarrow y = 4$$

$$x = 2 \Rightarrow y = 5$$

$$x = 3 \Rightarrow y = 6$$

$$x = 4 \Rightarrow y = 7$$

$$x = 5 \Rightarrow y = 8$$

$$\therefore R = \{(0, 3), (1, 4), (2, 5), (3, 6), (4, 7), (5, 8)\}$$

$$\therefore \text{Domain} : \{0, 1, 2, 3, 4, 5\}$$

$$\text{Range} : \{3, 4, 5, 6, 7, 8\}$$

(a) an arrow diagram, (b) a graph and (c) a set in roster form, wherever possible.

(i) $\{(x, y) \mid x = 2y, x \in \{2, 3, 4, 5\},$

$$y \in \{1, 2, 3, 4\}$$

(ii) $\{(x, y) \mid y = x + 3, x, y \text{ are natural numbers} < 10\}$

Solution :

i) $\{(x, y) \mid x = 2y, x \in \{2, 3, 4, 5\}, y \in \{1, 2, 3, 4\}$


a) Arrow diagram :

$$y = 1 \Rightarrow x = 2$$


$$y = 2 \Rightarrow x = 4$$

$$y = 3 \Rightarrow x = 6$$

$$y = 4 \Rightarrow x = 8$$



b) Graph :




c) a set in roster : $z \{(2, 1), (4, 2)\}$


ii) $\{(x, y) \mid y = x + 3,$

$$x, y \text{ are natural numbers} < 10\}$$

a) Arrow Diagram :



b) Graph :



c) a set in roster :

$$= \{(1, 4), (2, 5), (3, 6), (4, 7), (5, 8), (6, 9)\}$$

5. A company has four categories of employees given by Assistants (A), Clerks (C), Managers (M) and an Executive Officer (E). The company provide ₹10,000, ₹25,000, ₹50,000 and ₹1,00,000 as salaries to the people who work in the categories A, C, M and E respectively. If A_1, A_2, A_3, A_4 and A_5 were Assistants ; C_1, C_2, C_3, C_4 were Clerks ; M_1, M_2, M_3 were managers and E_1, E_2 were Executive officers and if the relation R is defined by xRy , where x is the salary given to person y , express the relation R through an ordered pair and an arrow diagram.


Solution :

a) Ordered Pair :

$\{(10000, A_1), (10000, A_2), (10000, A_3),$
 $(10000, A_4), (10000, A_5), (25000, C_1),$

$(25000, C_2), (25000, C_3), (25000, C_4),$
 $(50000, M_1), (50000, M_2), (50000, M_3),$
 $(100000, E_1), (100000, E_2)\}.$

b) Arrow Diagram :



III. FUNCTIONS :

Key Points


- ✓ A relation f between two non-empty sets X and Y is called a function from X to Y if, for each $x \in X$ there exists only one $y \in Y$ such that $(x, y) \in f$. That is, $f = \{(x, y) \mid \text{for all } x \in X, y \in Y\}$.
- ✓ If $f: X \rightarrow Y$ is a function then the set X is called the domain of the function f and the set Y is called its co-domain.
- ✓ If $f(a) = b$, then b is called 'image' of a under f and a is called a 'pre-image' of b .
- ✓ The set of all images of the elements of X under f is called the 'range' of f .
- ✓ $f: X \rightarrow Y$ is a function only if
 - (i) every element in the domain of f has an image.
 - (ii) the image is unique.
- ✓ If A and B are finite sets such that $n(A) = p$, $n(B) = q$ then the total number of functions that exist between A and B is q^p .
- ✓ The range of a function is a subset of its co-domain.

Example 1.6

Let $X = \{1, 2, 3, 4\}$ and $Y = \{2, 4, 6, 8, 10\}$ and $R = \{(1, 2), (2, 4), (3, 6), (4, 8)\}$. Show that R is a function and find its domain, co-domain and range?

Solution :

Pictorial representation of R is given in Figure. From the diagram, we see that for each $x \in X$, there exists only one $y \in Y$. Thus all elements in X have only one image in Y . Therefore R is a function. Domain $X = \{1, 2, 3, 4\}$; Co-domain $Y = \{2, 4, 6, 8, 10\}$; Range of $f = \{2, 4, 6, 8\}$.

**Example 1.7**

A relation ' f ' is defined by $f(x) = x^2 - 2$ where, $x \in \{-2, -1, 0, 3\}$

- (i) List the elements of f (ii) If f is a function?

Solution :

$$f(x) = x^2 - 2 \text{ where } x \in \{-2, -1, 0, 3\}$$

(i) $f(-2) = (-2)^2 - 2 = 2$;

$$f(-1) = (-1)^2 - 2 = -1$$

$$f(0) = (0)^2 - 2 = -2 ; f(3) = (3)^2 - 2 = 7$$

Therefore, $f = \{(-2, 2), (-1, -1), (0, -2), (3, 7)\}$

- (ii) We note that each element in the domain of f has a unique image. Therefore f is a function.

Example 1.8


If $X = \{-5, 1, 3, 4\}$ and $Y = \{a, b, c\}$, then which of the following relations are functions from X to Y ?

- (i) $R_1 = \{(-5, a), (1, a), (3, b)\}$
 (ii) $R_2 = \{(-5, b), (1, b), (3, a), (4, c)\}$
 (iii) $R_3 = \{(-5, a), (1, a), (3, b), (4, c), (1, b)\}$

Solution :

- (i) $R_1 = \{(-5, a), (1, a), (3, b)\}$


We may represent the relation R_1 in an arrow diagram.



- (ii) $R_2 = \{(-5, b), (1, b), (3, a), (4, c)\}$


Arrow diagram of R_2 is shown in Figure.

R_2 is a function as each element of X has a unique image in Y .



- (iii) $R_3 = \{(-5, a), (1, a), (3, b), (4, c), (1, b)\}$

Representing R_3 in an arrow diagram.



R_3 is not a function as $1 \in X$ has two images $a \in Y$ and $b \in Y$.

Note that the image of an element should always be unique.

Example 1.9

Given $f(x) = 2x - x^2$, find (i) $f(1)$ (ii) $f(x+1)$ (iii) $f(x) + f(1)$

Solution :

(i) Replacing x with 1, we get

$$f(1) = 2(1) - (1)^2 = 2 - 1 = 1$$

(ii) Replacing x with $x + 1$, we get

$$\begin{aligned} f(x+1) &= 2(x+1) - (x+1)^2 \\ &= 2x + 2 - (x^2 + 2x + 1) = -x^2 + 1 \end{aligned}$$

(iii) $f(x) + f(1) = (2x - x^2) + 1 = -x^2 + 2x + 1$

$+ b$ is not equal to $f(a) + f(b)$.

EXERCISE 1.3

1. Let $f = \{(x, y) \mid x, y \in \mathbb{N} \text{ and } y = 2x\}$ be a relation on \mathbb{N} . Find the domain, co-domain and range. Is this relation a function ?


Solution :

Given $f = \{(x, y) \mid x, y \in \mathbb{N} \text{ and } y = 2x\}$

Domain = $\{1, 2, 3, 4, \dots\}$

Co-domain = $\{1, 2, 3, 4, \dots\}$

Range = $\{2, 4, 6, 8, \dots\}$



Yes, f is a function.

Since all the elements of domain will be mapped into a unique element in co-domain.

2. Let $X = \{3, 4, 6, 8\}$. Determine whether the relation $R = \{(x, f(x)) \mid x \in X, f(x) = x^2 + 1\}$ is a function from X to \mathbb{N} ?

Solution :

Given $X = \{3, 4, 6, 8\}$

$$R = \{(x, f(x)) \mid x \in X, f(x) = x^2 + 1\}$$

$$x = 3 \Rightarrow f(x) = f(3) = 9 + 1 = 10$$

$$x = 4 \Rightarrow f(x) = f(4) = 16 + 1 = 17$$

$$x = 6 \Rightarrow f(x) = f(6) = 36 + 1 = 37$$

$$x = 8 \Rightarrow f(x) = f(8) = 64 + 1 = 65$$

$$R = \{(3, 10), (4, 17), (6, 37), (8, 65)\}$$

\therefore The relation $R : X \rightarrow \mathbb{N}$ is a function.

2

evaluate

(i) $f(-1)$ (ii) $f(2a)$ (iii) $f(2)$ (iv) $f(x-1)$

Solution :

$$\text{Given } f: x \rightarrow x^2 - 5x + 6$$

$$\Rightarrow f(x) = x^2 - 5x + 6$$

$$\begin{aligned} \text{(i) } f(-1) &= (-1)^2 - 5(-1) + 6 \\ &= 1 + 5 + 6 \\ &= 12 \end{aligned}$$

$$\begin{aligned} \text{(ii) } f(2a) &= (2a)^2 - 5(2a) + 6 \\ &= 4a^2 - 10a + 6 \end{aligned}$$

$$\begin{aligned} \text{(iii) } f(2) &= 2^2 - 5(2) + 6 \\ &= 4 - 10 + 6 \\ &= 0 \end{aligned}$$

$$\begin{aligned} \text{(iv) } f(x-1) &= (x-1)^2 - 5(x-1) + 6 \\ &= x^2 - 2x + 1 - 5x + 5 + 6 \\ &= x^2 - 7x + 12 \end{aligned}$$

4. A graph representing the function $f(x)$ is given in figure it is clear that $f(9) = 2$.


(i) Find the following values of the function

(a) $f(0)$ (b) $f(7)$ (c) $f(2)$ (d) $f(10)$

(ii) For what value of x is $f(x) = 1$?

(iii) Describe the following (i) Domain (ii) Range.

(iv) What is the image of 6 under f ?



Solution :

(i) a) $f(0) = 9$ b) $f(7) = 6$
c) $f(2) = 6$ d) $f(10) = 0$

(ii) When $x = 9.5$, $f(x) = 1$.

(iii) a) Domain : $\{x / 0 \leq x \leq 10, x \in \mathbb{R}\}$

b) Range : $\{x / 0 \leq x \leq 9, x \in \mathbb{R}\}$

(iv) Image of 6 = $f(6) = 5$.

5. Let $f(x) = 2x + 5$. If $x \neq 0$ then find $\frac{f(x+2) - f(2)}{x}$.

Solution :

$$\begin{aligned} \text{Given } f(x) &= 2x + 5 \\ f(x+2) &= 2(x+2) + 5 \\ &= 2x + 9 \\ f(2) &= 2(2) + 5 \\ &= 9 \end{aligned}$$

$$\therefore \frac{f(x+2) - f(x)}{x} = \frac{2x + 9 - 9}{x} = 2$$

6. A function f is defined by $f(x) = 2x - 3$

(i) find $\frac{f(0) + f(1)}{2}$

(ii) find x such that $f(x) = 0$

(iii) find x such that $f(x) = x$

(iv) find x such that $f(x) = f(1 - x)$.

Solution :

$$\text{Given } f(x) = 2x - 3$$


$$\begin{aligned} \text{i) } \frac{f(0) + f(1)}{2} &= \frac{(-3) + (-1)}{2} \\ &= \frac{-4}{2} \\ &= -2 \end{aligned}$$


$$\begin{aligned} \text{(ii) } f(x) = 0 &\Rightarrow 2x - 3 = 0 \\ &\Rightarrow 2x = 3 \\ &\Rightarrow x = \frac{3}{2} \end{aligned}$$

$$\begin{aligned} \text{(iii) } f(x) = x &\Rightarrow 2x - 3 = x \\ &\Rightarrow 2x - x = 3 \\ &\Rightarrow x = 3 \end{aligned}$$

$$\begin{aligned} \text{(iv) } f(x) = 1 - x &\Rightarrow 2x - 3 = 1 - x \\ &\Rightarrow 2x + x = 1 + 3 \\ &\Rightarrow 3x = 4 \\ &\Rightarrow x = \frac{4}{3} \end{aligned}$$

7. An open box is to be made from a square piece of material, 24 cm on a side, by cutting equal squares from the corners and turning up the sides as shown in the figure. Express the volume V of the box as a function of x .





Solution :

From the given data, it is clear that length l = breadth, $b = 24 - 2x$ cm, height = x cm.

$$\begin{aligned}\therefore \text{Volume of the box, } V &= lbh \\ &= (24 - 2x)^2 x \\ &= (576 + 4x^2 - 96x) x \\ &= 4x^3 - 96x^2 + 576x\end{aligned}$$

\therefore Volume is expressed as a function of x .

8. A function f is defined by $f(x) = 3 - 2x$. Find x such that $f(x^2) = (f(x))^2$.

Solution :

$$\begin{aligned}\text{Given } f(x) &= 3 - 2x \text{ and} \\ f(x^2) &= (f(x))^2 \\ \Rightarrow 3 - 2x^2 &= (3 - 2x)^2 \\ \Rightarrow 3 - 2x^2 &= 9 + 4x^2 - 12x \\ \Rightarrow 6x^2 - 12x + 6 &= 0 \\ \Rightarrow x^2 - 2x + 1 &= 0 \\ \Rightarrow (x - 1)^2 &= 0 \\ \Rightarrow x &= 1 \text{ (twice)}\end{aligned}$$

9. A plane is flying at a speed of 500 km per hour. Express the distance d travelled by the plane as function of time t in hours.

Solution :

$$\begin{aligned}\text{Given, Speed of the plane} &= 500 \text{ km / h} \\ \text{time} &= t \text{ hrs} \\ \text{distance} &= d \text{ km} \\ \therefore \text{Distance} &= \text{Time} \times \text{Speed} \\ \therefore d &= 500t\end{aligned}$$

10. The data in the adjacent table depicts the length of a woman's forearm and her corresponding height. Based on this data, a student finds a relationship between the height (y) and the forearm length (x) as $y = ax + b$, where a, b are constants.

Length 'x' of forearm (in cm)	Height 'y' (in inches)
45.5	65.5
35	56
45	65
50	69.5
55	74

- (i) Check if this relation is a function.
 (ii) Find a and b .
 (iii) Find the height of a woman whose
 (iv) Find the length of forearm of a woman if her height is 53.3 inches.

Solution :

- i) From the table, it is clear that each element of domain is having a unique image in co-domain and hence it is a function.

ii) $y = ax + b$

when $x = 55, y = 75 \Rightarrow 75 = 55a + b$

when $x = 42, y = 62 \Rightarrow 62 = 42a + b$

Subtracting, $13 = 13a$

$a = 1$

$\therefore 75 = 55(1) + b$

$\therefore b = 75 - 55 = 20$

$\therefore a = 1, b = 20$

iii) When $x = 48$; $y = ?$

$$\therefore y = ax + b$$

$$\Rightarrow y = x + 20$$

$$\text{when } x = 48, y = 48 + 20$$

$$\therefore y = 68$$

\therefore Height of a woman = 68 inches.

iv) When $y = 60.54$ inches, $x = ?$

$$\therefore y = x + 20$$

$$\Rightarrow 60.54 = x + 20$$

$$\therefore x = 40.54 \text{ cm}$$

IV. REPRESENTATION OF FUNCTIONS AND TYPE OF FUNCTIONS :

Key Points

- ✓ A function may be represented by
 - (a) a set of ordered pairs (b) a table form (c) an arrow diagram (d) a graphical form
- ✓ Every function can be represented by a curve in a graph. But not every curve drawn in a graph will represent a function.
- ✓ A curve drawn in a graph represents a function, if every vertical line intersects the curve in at most one point.
- ✓ A function $f: A \rightarrow B$ is called one - one function if distinct elements of A have distinct images in B.
- ✓ If for all $a_1, a_2 \in A, f(a_1) = f(a_2)$ implies $a_1 = a_2$, then f is called **one-one function**.
- ✓ A function $f: A \rightarrow B$ is called **many-one function** if two or more elements of A have same image in B.
- ✓ A function $f: A \rightarrow B$ is called many-one if f is not one-one.
- ✓ A function $f: A \rightarrow B$ is said to be **onto function** if the range of f is equal to the co-domain of f .
- ✓ That is a every element in the co-domain B has a pre-image in the domain A.
- ✓ An onto function is also called a **surjection**.
- ✓ If $f: A \rightarrow B$ is an onto function then, the range of $f = B$. That is $f(A) = B$.
- ✓ A function $f: A \rightarrow B$ is called an **into function** if there exists atleast one element in B which is not the image of any element of A.
- ✓ The range of f is a proper subset of the co-domain of f .
- ✓ A function $f: A \rightarrow B$ is called 'into' if it is not 'onto'.
- ✓ If a function $f: A \rightarrow B$ is both **one-one and onto**, then f is called a **bijection** from A to B.

- ✓ A function represented in a graph is one-one, if every horizontal line intersects the curve in at most one point.
- ✓ A function $f: A \rightarrow B$ is called a **constant function** if the range of f contains only one element. That is $f(x) = c$, for all $x \in A$ and for some fixed $c \in B$.
- ✓ Let A be a non-empty set. Then the function $f: A \rightarrow A$ defined by $f(x) = x$ for all $x \in A$ is called an **identity function** on A and is denoted by I_A .
- ✓ A function $f: A \rightarrow B$ is called a **real valued function** if the range of f is a subset of the set of all real numbers R . That is, $f(A) \subseteq R$.

Example 1.10

Using vertical line test, determine which of the following curves (Fig. 1.18(a), 1.18(b), 1.18(c), 1.18(d)) represent a function ?




Fig. (1.18(a))




Fig. 1.18(b)





Fig. 1.18(c)

Solution :

The curves Fig. 1.18(a) and Fig. 1.18(c) do not represent a function as the vertical lines meet the curves in two points P and Q.



The curves in Fig. 1.18(b) and Fig. 1.18(d) represent a function as the vertical lines meet the curve in at most one point.

Example 1.11

Let $A = \{1, 2, 3, 4\}$ and $B = \{2, 5, 8, 11, 14\}$ be two sets. Let $f: A \rightarrow B$ be a function given by $f(x) = 3x - 1$. Represent this function

- (i) by arrow diagram
- (ii) in a table form
- (iii) as a set of ordered pairs
- (iv) in a graphical form

Solution :


$A = \{1, 2, 3, 4\}$; $B = \{2, 5, 8, 11, 14\}$; $f(x) = 3x - 1$

$f(1) = 3(1) - 1 = 3 - 1 = 2$; $f(2) = 3(2) - 1 = 6 - 1 = 5$

$f(3) = 3(3) - 1 = 9 - 1 = 8$; $f(4) = 4(3) - 1 =$

(i) Arrow diagram

Let us represent the function $f: A \rightarrow B$ by an arrow diagram

**(ii) Table form**

The given function f can be represented in a tabular form as given below

x	1	2	3	4
$f(x)$	2	5	8	11

(iii) Set of ordered pairs

The function f can be represented as a set of


ordered pair as

$$f = \{(1, 2), (2, 5), (3, 8), (4, 11)\}$$

(iv) Graphical form

In the adjacent xy -plane the points

$(1, 2), (2, 5), (3, 8), (4, 11)$ are plotted

**Example 1.12**

Using horizontal line test (Fig. 1.35(a), 1.35(b), 1.35(c)), determine which of the following functions are one-one.




Fig. 1.35 (a)




Fig. 1.35(b)




Fig. 1.35(c)

Solution :

The curves in Fig.1.35(a) and Fig. 1.35(c) represent a one-one function as the horizontal lines meet the curves in only one point P.

The curve Fig.1.35(b) does not represent a one-one function, since, the horizontal line meet the curve in two points P and Q.

Example 1.13


Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and $f = \{(1, 4), (2, 5), (3, 6)\}$

that f is one-one but not onto function.

Solution:

$A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$; $f = \{(1, 4), (2, 5), (3, 6)\}$

Then f is a function from A to B and for different elements in A , there are different images in B . Hence f is one-one function. Note that the element 7 in the co-domain does not have any pre-image in the domain. Hence f is not onto.

**Example 1.14**

If $A = \{-2, -1, 0, 1, 2\}$ and $f: A \rightarrow B$ is an onto function defined by $f(x) = x^2 + x + 1$ then find B .

Solution :

Given $A = \{-2, -1, 0, 1, 2\}$ and $f(x) = x^2 + x + 1$

$$f(-2) = (-2)^2 + (-2) + 1 = 3 ;$$

$$f(-1) = (-1)^2 + (-1) + 1 = 1$$

$$f(0) = 0^2 + 0 + 1 = 1 ;$$

$$f(1) = 1^2 + 1 + 1 = 3$$

$$f(2) = 2^2 + 2 + 1 = 7$$

Since, f is an onto function, range of $f = B =$ co-domain of f .

Therefore, $B = \{1, 3, 7\}$.

Example 1.15

Let f be a function $f: N \rightarrow N$ be defined by $f(x) = 3x + 2, x \in N$

(i) Find the images of 1, 2, 3 (i i)

Find the pre-images of 29, 53

(iii) Identify the type of function

Solution : $f(x) = 3x + 2, x \in N$

(i) If $x = 1$, $f(1) = 3(1) + 2 = 5$

If $x = 2$, $f(2) = 3(2) + 2 = 8$

If $x = 3$, $f(3) = 3(3) + 2 = 11$

The images of 1, 2, 3 are 5, 8, 11 respectively.

(ii) If x is the pre-image of 29, then $f(x) = 29$.

$$\text{Hence } 3x + 2 = 29$$

$$3x = 27 \Rightarrow x = 9.$$

Similarly, if x is the pre-image of 53, then $f(x) = 53$. Hence $3x + 2 = 53$

$$3x = 51 \Rightarrow x = 17.$$

Thus the pre-images of 29 and 53 are 9 and 17 respectively.

(iii) Since different elements of N have different images in the co-domain, the function f is one-one function.

The co-domain of f is N .

But the range of $f = \{5, 8, 11, 14, 17, \dots\}$ is a proper subset of N . Therefore f is not an onto function. That is, f is an into function.

Thus f is one-one and into function.

Example 1.16

Forensic scientists can determine the height (in cms) of a person based on the length of their thigh bone. They usually do so using the function $h(b) = 2.47b + 54.10$ where b is the length of the thigh bone.

(i) Check if the function h is one-one

(ii) Also find the height of a person if the length of his thigh bone is 50 cms.

(iii) Find the length of the thigh bone if the height of a person is 147.96 cms.

Solution :

(i) To check if h is one-one, we assume that $h(b_1) = h(b_2)$.

Then we get, $2.47b_1 + 54.10 = 2.47b_2 + 54.10$

$$2.47b_1 = 2.47b_2 \Rightarrow b_1 = b_2$$

Thus, $h(b_1) = h(b_2) \Rightarrow b_1 = b_2$. So, the function h is one-one.

(ii) If the length of the thigh bone $b = 50$, then the height is

$$h(50) = (2.47 \times 50) + 54.10 = 177.6 \text{ cms.}$$

(iii) If the height of a person is 147.96 cms, then $h(b) = 147.96$ and so the length of the thigh bone is given by $2.47b + 54.10 = 147.96$.

$$b = \frac{93.86}{2.47} = 38$$

Therefore, the length of the thigh bone is 38 cms.

Example 1.17

Let f be a function from R to R defined by $f(x) = 3x - 5$. Find the values of a and b given that $(a, 4)$ and $(1, b)$ belong to f .

Solution :

$f(x) = 3x - 5$ can be written as

$$f = \{(x, 3x - 5) \mid x \in R\}$$

$(a, 4)$ means the image of a is 4.

That is, $f(a) = 4$

$$3a - 5 = 4 \Rightarrow a = 3$$

$(1, b)$ means the image of 1 is b .

That is, $f(1) = b \Rightarrow b = -2$

$$3(1) - 5 = b \Rightarrow b = -2$$

Example 1.18

time ' t ' hours is given by $S(t) = \frac{t^2 + t}{2}$. Find the distance travelled by the particle after

(i) three and half hours.

(ii) eight hours and fifteen minutes.

Solution :

The distance travelled by the particle in time t hours is given by $S(t) = \frac{t^2 + t}{2}$.

(i) $t = 3.5$ hours. Therefore,

$$S(3.5) = \frac{(3.5)^2 + 3.5}{2} = \frac{15.75}{2} = 7.875$$

The distance travelled in 3.5 hours is 7.875 kms.

(ii) $t = 8.25$ hours. Therefore, $S(8.25) =$

$$\frac{(8.25)^2 + 8.25}{2} = \frac{76.3125}{2} = 38.15625$$

The distance travelled in 8.25 hours is 38.16 kms, approximately.

Example 1.19

If the function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by


$$f(x) = \begin{cases} 2x + 7, & x < -2 \\ x^2 - 2, & -2 \leq x < 3 \\ 3x - 2, & x \geq 3 \end{cases} \text{ then find the values of}$$

(i) $f(4)$ (ii) $f(-2)$ (iii) $f(4) + 2f(1)$

(iv) $\frac{f(1) - 3f(4)}{f(-3)}$

Solution :

The function f is defined by three values in intervals I, II, III as shown below



$$f(x) = 2x + 7$$

$$f(x) = x^2 - 2$$

$$f(x) = 3x - 2$$

For a given value of $x = a$, find out the inter-

$f(a)$ using the particular value defined in that interval.

(i) First, we see that,

$x = 4$ lie in the third interval.

Therefore, $f(x) = 3x - 2$;

$$f(4) = 3(4) - 2 = 10$$

(ii) $x = -2$ lies in the second interval.

Therefore, $f(x) = x^2 - 2$;

$$f(-2) = (-2)^2 - 2 = 2$$

(iii) From (i), $f(4) = 10$.

To find $f(1)$, first we see that $x = 1$ lies in the second interval.

Therefore, $f(x) = x^2 - 2$;

$$f(1) = (1)^2 - 2 = -1$$

$$\text{So, } f(4) + 2f(1) = 10 + 2(-1) = 8$$

(iv) We know that $f(1) = -1$ and $f(4) = 10$.


For finding $f(-3)$, we see that $x = -3$, lies in the first interval.

Therefore, $f(x) = 2x + 7$; thus, $f(-3) = 2(-3) + 7 = 1$

$$\text{Hence, } \frac{f(1) - 3f(4)}{f(-3)} = \frac{-1 - 3(10)}{1} = -31$$

EXERCISE 1.4

1. Determine whether the graph given below represent functions. Give reason for your answers concerning each graph.



Solution :

(i) The curve do not represent a function since it meets y-axis at 2 points.

(ii) The curve represents a function as it meets x-axis or y-axis at only one point.

(iii) The curve do not represent a function since it meets y-axis at 2 points.

(iv) The line represents a function as it meets axes at origin.

2. Let $f: A \rightarrow B$ be a function define by f

$(x) = \frac{x}{2} - 1$, where $A = \{2, 4, 6, 10, 12\}$, $B = \{0, 1, 2, 4, 5, 9\}$. Represent f by

(i) set of ordered pairs ; (ii) a table ; (iii) an arrow diagram ; (iv) a graph

Solution :

$$\text{Given } f(x) = \frac{x}{2} - 1$$

$$x = 2 \Rightarrow f(2) = 1 - 1 = 0$$

$$x = 4 \Rightarrow f(4) = 2 - 1 = 1$$

$$x = 6 \Rightarrow f(6) = 3 - 1 = 2$$

$$x = 10 \Rightarrow f(10) = 5 - 1 = 4$$




$$x = 12 \Rightarrow f(12) = 6 - 1 = 5$$

(i) Set of order pairs :


$$f = \{(2, 0), (4, 1), (6, 2), (10, 4), (12, 5)\}$$

(ii) Table :

x	2	4	6	10	12
$f(x)$	0	1	2	4	5

(iii) Arrow diagram :**(iv) Graph****3. Represent the function $f = \{(1, 2), (2, 2), (3, 2), (4, 3), (5, 4)\}$ through****(i) an arrow diagram****(ii) a****table form****(iii) a graph****Solution :****(i) Arrow Diagram :****(ii) Table Form :**

x	1	2	3	4	5
$f(x)$	2	2	2	3	4

(iii) Graph :**4. Show that the function $f: \mathbb{N} \rightarrow \mathbb{N}$ defined $f(x) = 2x - 1$ is one-one but not onto.****Solution :**Given $f: \mathbb{N} \rightarrow \mathbb{N}$ defined by $f(x) = 2x - 1$.

$$x = 1 \Rightarrow f(1) = 2 - 1 = 1$$

$$x = 2 \Rightarrow f(2) = 4 - 1 = 3$$

$$x = 3 \Rightarrow f(3) = 6 - 1 = 5$$

$$x = 4 \Rightarrow f(4) = 8 - 1 = 7 \dots\dots$$

It is clear that f is a function from $N \rightarrow N$ and for different elements in domain, there are different images in co-domain.

$\therefore f$ is one to one function.

But co-domain is N and $\text{Range} = \{1, 3, 5, 7, \dots\}$

$\therefore \text{Range} \neq \text{Co-domain}$.

$\therefore f$ is not on-to.

5. Show that the function $f: N \rightarrow N$ defined by $f(m) = m^2 + m + 3$ is one-one function.

Solution :

Given $f: N \rightarrow N$ defined by $f(m) = m^2 + m + 3$

$$f(m) = m^2 + m + 3$$

$$m = 1 \Rightarrow f(1) = 1 + 1 + 3 = 5$$

$$m = 2 \Rightarrow f(2) = 4 + 2 + 3 = 9$$

$$m = 3 \Rightarrow f(3) = 9 + 3 + 3 = 15$$

$$m = 4 \Rightarrow f(4) = 16 + 4 + 3 = 23 \dots\dots\dots$$

For different elements of domain, there are different images in co-domain.

$\therefore f$ is one-one function.

6. Let $A = \{1, 2, 3, 4\}$ and $B = N$. Let $f: A \rightarrow B$ be defined by $f(x) = x^3$ then,

(i) find the range of f

(ii) identify the type of function

Solution :

Given $A = \{1, 2, 3, 4\}$, $B = N$

$$f(x) = x^3$$

$$x = 1 \Rightarrow f(1) = 1$$

$$x = 2 \Rightarrow f(2) = 8$$

$$x = 3 \Rightarrow f(3) = 27$$

$$x = 4 \Rightarrow f(4) = 64$$

(i) Range of $f = \{1, 8, 27, 64\}$

(ii) f is one-one (diff. elements have diff. images) and

f is into ($\text{Range} \neq \text{co-domain}$)

7. In each of the following cases state whether the function is bijective or not. Justify your answer.

(i) $f: R \rightarrow R$ defined by $f(x) = 2x + 1$

(ii) $f: R \rightarrow R$ defined by $f(x) = 3 - 4x^2$

Solution :

(i) $f: R \rightarrow R$ defined by $f(x) = 2x + 1$

$$\text{Let } f(x_1) = f(x_2)$$

$$\Rightarrow 2x_1 + 1 = 2x_2 + 1$$

$$\Rightarrow 2x_1 = 2x_2$$

$$\Rightarrow x_1 = x_2$$

$$\therefore f(x_1) = f(x_2) \Rightarrow x_1 = x_2$$

$\therefore f$ is 1 - 1 function.

$$y = 2x + 1$$

$$\therefore 2x = y - 1$$

$$\Rightarrow x = \frac{y-1}{2}$$

$$\therefore f(x) = 2\left(\frac{y-1}{2}\right) + 1$$

$$\therefore f \text{ is onto.} \quad = y$$

$\therefore f$ is one-one and onto

$\Rightarrow f$ is bijective.

(ii) $f: R \rightarrow R$ defined by $f(x) = 3 - 4x^2$.

$$\text{Let } f(x_1) = f(x_2)$$

$$3 - 4x_1^2 = 3 - 4x_2^2$$

$$x_1^2 = x_2^2$$

$$x_1 = x_2 \text{ (or) } x_1 = -x_2$$

f is not 1 - 1.

(Example : when $x = -1$, $f(x) = f(-1) = -1$

$$\text{when } x = 1, \quad f(x) = f(1) = -1$$

\therefore Two different elements in domain have same images in co-domain.

Also, any even number in the co-domain is not image of any element x in the domain.

$\therefore f$ is not onto

$\therefore f$ is not bijective.

8. Let $A = \{-1, 1\}$ and $B = \{0, 2\}$. If the function $f: A \rightarrow B$ defined by $f(x) = ax + b$ is an onto function? Find a and b .

Solution :

$$\text{Given } A = \{-1, 1\}, B = \{2, 2\}$$

$$f(x) = ax + b \text{ is onto function.}$$

$$\therefore f(-1) = 0 \Rightarrow -a + b = 0 \text{ --- (1)}$$

$$f(1) = 2 \Rightarrow a + b = 2 \text{ --- (2)}$$

Solving (1) and (2)

$$2b = 2$$

$$b = 1$$

$$\Rightarrow a = 1$$

$$\therefore a = 1, b = 1$$

9. If the function f is defined by $f(x) =$

$$\begin{cases} x+2 & \text{if } x > 1 \\ 2 & \text{if } -1 \leq x \leq 1 \\ x-1 & \text{if } -3 < x < -1 \end{cases} ; \text{ find the values of}$$

$$(i) f(3) \quad (ii) f(0) \quad (iii) f(-1, 5)$$

$$(iv) f(2) + f(-2)$$

Solution :

$$\text{Given } f(x) = \begin{cases} x+2 & \text{if } x > 1 \\ 2 & \text{if } -1 \leq x \leq 1 \\ x-1 & \text{if } -3 < x < -1 \end{cases}$$

$$(i) f(3) = 3 + 2 \quad (\because 3 \in (1 < x < \infty))$$

$$= 5$$

$$(ii) f(0) = 2 \quad (\because 0 \in (-1 \leq x \leq 1))$$

$$(iii) f(-1.5) = -1.5 - 1 \quad (\because -1.5 \in (-3 < x < -1))$$

$$= -2.5$$

$$(iv) f(2) + f(-2) \quad (\because 2 \in (1 < x < \infty))$$

$$= (2 + 2) + (-2 - 1) \quad (\because -2 \in (-3 < x < -1))$$

$$= 4 - 3$$

$$= 1$$

10. A function $f: [-5, 9] \rightarrow \mathbb{R}$ is defined as follows :

$$f(x) = \begin{cases} 6x+1 & \text{if } -5 \leq x < 2 \\ 5x^2-1 & \text{if } 2 \leq x < 6 \\ 3x-4 & \text{if } 6 \leq x < 9 \end{cases}$$

$$\text{Find (i) } f(-3) + f(2) \quad (ii) f(7) - f(1)$$

$$(iii) 2f(4) + f(8) \quad (iv) \frac{2f(-2) - f(6)}{f(4) + f(-2)}$$

Solution :

$$\text{Given } f(x) = \begin{cases} 6x+1 & \text{if } -5 \leq x < 2 \\ 5x^2-1 & \text{if } 2 \leq x < 6 \\ 3x-4 & \text{if } 6 \leq x < 9 \end{cases}$$

$$(i) f(-3) + f(2) \quad (\because -3 \in (-5 \leq x < 2))$$

$$= [6(-3) + 1] + [5(4) - 1] \quad \text{and}$$

$$= -17 + 19 \quad 2 \in (2 \leq x < 6)$$

$$(ii) f(7) - f(1) \quad (\because 7 \in (6 \leq x < 9))$$

$$= [3(7) - 4] - [6(1) + 1] \quad \text{and}$$

$$= 17 - 7 \quad 1 \in (-5 \leq x < 2)$$

$$= 10$$

$$(iii) 2f(4) + f(8) \quad (\because 4 \in (2 \leq x < 6))$$

$$= 2[5(16) - 1] + [3(8) - 4] \quad \text{and}$$

$$= 2[79] + 20 \quad 8 \in (6 \leq x < 9)$$

$$= 158 + 20$$

$$= 178$$

(iv)

$$\begin{aligned}
 & \frac{2f(-2) - f(6)}{f(4) + f(-2)} \quad (\because -2 \in (-5 \leq x < 2) \\
 & \quad \quad \quad 6 \in (6 \leq x \leq 9) \\
 & \quad \quad \quad -2 \in (-5 \leq x < 2)) \\
 &= \frac{2(-11) - 14}{79 + (-11)} \\
 &= \frac{-22 - 14}{79 - 11} \\
 &= \frac{-36}{68} \\
 &= \frac{-9}{17}
 \end{aligned}$$

11. The distance S an object travels under the influence of gravity in time t seconds is given by $S(t) = \frac{1}{2}gt^2 + at + b$ where,

a and b are constants. Check if the function $S(t)$ is one-one.

Solution :

$$\text{Given } S(t) = \frac{1}{2}gt^2 + at + b$$

To check $s(t)$ is one-one, we consider,

$$s(t_1) = s(t_2)$$

$$\Rightarrow \frac{1}{2}gt_1^2 + at_1 + b = \frac{1}{2}gt_2^2 + at_2 + b$$

$$\Rightarrow \frac{1}{2}g(t_1^2 - t_2^2) + a(t_1 - t_2) = 0$$

$$\Rightarrow (t_1 - t_2) \left[\frac{1}{2}g(t_1 + t_2) + a \right] = 0$$

$$\Rightarrow t_1 - t_2 = 0 \quad (\because \frac{1}{2}g(t_1 + t_2) + a \neq 0)$$

$$\Rightarrow t_1 = t_2$$

$\therefore s(t)$ is one-one.

12. The function 't' which maps temperature in Celsius (C) into temperature in Fahrenheit (F) is defined by $t(C) = F$ where $F = \frac{9}{5}C + 32$, Find,

(i) $t(0)$ (ii) $t(28)$ (iii) $t(-10)$

(iv) the value of C when $t(C) = 212$

(v) the temperature when the Celsius value is equal to the Fahrenheit value.

Solution :

$$\text{Given } t(C) = F = \frac{9C}{5} + 32$$

$$(i) \quad t(0) = \frac{9(0)}{5} + 32 = 32^\circ F$$

$$(ii) \quad t(28) = \frac{9(28)}{5} + 32 = 50.4 + 32 = 82.4^\circ F$$

$$(iii) \quad t(-10) = \frac{9(-10)}{5} + 32 = -18 + 32 = 14^\circ F$$

(iv) When $t(c) = 212$,

$$212 = \frac{9C}{5} + 32$$

$$\Rightarrow \frac{9C}{5} = 180$$

$$\Rightarrow C = 180 \times \frac{5}{9}$$

$$\Rightarrow C = 100^\circ C$$

(v) When Celsius value = Fahrenheit value,

$$C = \frac{9C}{5} + 32$$

$$\Rightarrow 5C = 9C + 160$$

$$\Rightarrow -4C = 160$$

$$\Rightarrow C = -40^\circ$$

V. COMPOSITION OF FUNCTIONS :

Key Points

- ✓ Let $f: A \rightarrow B$ and $g: B \rightarrow C$ be two functions. Then the **composition of f and g** denoted by $g \circ f$ is defined as the function $g \circ f(x) = g(f(x))$ for all $x \in A$.
- ✓ Generally, $f \circ g \neq g \circ f$ for any two functions f and g . So, composition of functions is not commutative.
- ✓ Composition of three functions is always associative. That is $f \circ (g \circ h) = (f \circ g) \circ h$.
- ✓ A function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined $f(x) = mx + c$, $m \neq 0$ is called a **linear function**. Geometrically this represents a straight line in the graph.
- ✓ $f: \mathbb{R} \rightarrow [0, \infty)$ defined $f(x) = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$ is called a **modulus (or) Absolute value function**.
- ✓ Modulus function is not a linear function but it is composed of two linear functions x and $-x$.
- ✓ A function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = ax^2 + bx + c$, ($a \neq 0$) is called a **quadratic function**.
- ✓ A function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = ax^3 + bx^2 + cx + d$, ($a \neq 0$) is called a **cubic function**.
- ✓ A function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = c$, for all $x \in \mathbb{R}$ is called a **constant function**.

Example 1.20

Find $f \circ g$ and $g \circ f$ when $f(x) = 2x + 1$ and $g(x) = x^2 - 2$.

Solution :

$$f(x) = 2x + 1, g(x) = x^2 - 2$$

$$f \circ g(x) = f(g(x)) = f(x^2 - 2)$$

$$= 2(x^2 - 2) + 1 = 2x^2 - 3$$

$$g \circ f(x) = g(f(x)) = g(2x + 1)$$

$$= (2x + 1)^2 - 2 = 4x^2 + 4x - 1$$

$$\text{Thus } f \circ g = 2x^2 - 3, g \circ f = 4x^2 + 4x - 1.$$

From the above, we see that $f \circ g \neq g \circ f$.

Example 1.21

Represent the function $f(x) = \sqrt{2x^2 - 5x + 3}$ as a composition of two functions.

Solution :

We set

$$f_2(x) = 2x^2 - 5x + 3 \text{ and } f_1(x) = \sqrt{x}$$

Then,

$$\begin{aligned} f(x) &= \sqrt{2x^2 - 5x + 3} = \sqrt{f_2(x)} \\ &= f_1[f_2(x)] = f_1 f_2(x) \end{aligned}$$

Example 1.22

If $f(x) = 3x - 2$, $g(x) = 2x + k$ and if $f \circ g = g \circ f$, then find the value of k .

Solution :

$$f(x) = 3x - 2, g(x) = 2x + k$$

$$f \circ g(x) = f(g(x))$$

$$= f(2x + k) = 3(2x + k) - 2 = 6x + 3k - 2$$

$$\text{Thus, } f \circ g(x) = 6x + 3k - 2.$$

$$g \circ f(x) = g(3x - 2) = 2(3x - 2) + k$$

$$\text{Thus, } g \circ f(x) = 6x - 4 + k.$$

$$\text{Given that } f \circ g = g \circ f$$

$$\text{Therefore, } 6x + 3k - 2 = 6x - 4 + k$$

$$6x - 6x + 3k - k = -4 + 2 \Rightarrow k = -1$$

Example 1.23

Find k if $f \circ f(k) = 5$ where $f(k) = 2k - 1$.

Solution :

$$\begin{aligned} f \circ f(k) &= f(f(k)) \\ &= 2(2k - 1) - 1 = 4k - 3. \end{aligned}$$

But, it is given that $f \circ f(k) = 5$

$$\text{Therefore } 4k - 3 = 5 \Rightarrow k = 2.$$

Example 1.24

$f(x) = 2x + 3$, $g(x) = 1 - 2x$ and $h(x) = 3x$. Prove that $f \circ (g \circ h) = (f \circ g) \circ h$.

Solution :

$$f(x) = 2x + 3, g(x) = 1 - 2x, h(x) = 3x$$

$$\begin{aligned} \text{Now, } (f \circ g)(x) &= f(g(x)) = f(1 - 2x) \\ &= 2(1 - 2x) + 3 = 5 - 4x \end{aligned}$$

$$\begin{aligned} \text{Then, } (f \circ g) \circ h(x) &= (f \circ g)(h(x)) = (f \circ g)(3x) \\ &= 5 - 4(3x) = 5 - 12x \quad \dots(1) \end{aligned}$$

$$(g \circ h)(x) = g(h(x)) = g(3x) = 1 - 2(3x) = 1 - 6x$$

$$\begin{aligned} \text{So, } f \circ (g \circ h)(x) &= f(1 - 6x) = 2(1 - 6x) + 3 \\ &= 5 - 12x \quad \dots(2) \end{aligned}$$

From (1) and (2), we get $f \circ (g \circ h) = (f \circ g) \circ h$.

Example 1.25

Find x if $gff(x) = fgg(x)$, given $f(x) = 3x + 1$ and $g(x) = x + 3$.

Solution :

$$\begin{aligned} gff(x) &= g[f\{f(x)\}] \text{ (This mean "g of f of f of x")} \\ &= g[f(3x + 1)] = g[3(3x + 1) + 1] = g(9x + 4) \end{aligned}$$

$$g(9x + 4) = [(9x + 4) + 3] = 9x + 7$$

$$\begin{aligned} fgg(x) &= f[g\{g(x)\}] \text{ (This means "f of g of x")} \\ &= f[g(x + 3)] = f[(x + 3) + 3] = f(x + 6) \end{aligned}$$

$$f(x + 6) = [3(x + 6) + 1] = 3x + 19$$

These two quantities being equal, we get $9x + 7 = 3x + 19$. Solving this equation we obtain $x = 2$.

EXERCISE 1.5

- Using the functions f and g given below, find $f \circ g$ and $g \circ f$. Check whether $f \circ g = g \circ f$.

$$(i) f(x) = x - 6, g(x) = x^2$$

$$(ii) f(x) = \frac{2}{x}, g(x) = 2x^2 - 1$$

$$(iii) f(x) = \frac{x+6}{3}, g(x) = 3 - x$$

$$(iv) f(x) = 3 + x, g(x) = x - 4$$

$$(v) f(x) = 4x^2 - 1, g(x) = 1 + x$$

Solution :

$$(i) f(x) = x - 6, g(x) = x^2$$

$$(f \circ g)(x) = f(g(x))$$

$$= f(x^2)$$

$$= x^2 - 6$$

$$(g \circ f)(x) = g(f(x))$$

$$= g(x - 6)$$

$$= (x - 6)^2$$

$$= x^2 - 12x + 36$$

$$\therefore f \circ g \neq g \circ f$$

$$(ii) \quad f(x) = \frac{2}{x}, g(x) = 2x^2 - 1$$

$$(f \circ g)(x) = f(g(x))$$

$$= f(2x^2 - 1)$$

$$= \frac{2}{2x^2 - 1}$$

$$(g \circ f)(x) = g(f(x))$$

$$= g\left(\frac{2}{x}\right)$$

$$= 2\left(\frac{2}{x}\right)^2 - 1$$

$$= \frac{8}{x^2} - 1$$

$$\therefore f \circ g \neq g \circ f$$

$$(iii) \quad f(x) = \frac{x}{3}, g(x) = 3 - x$$

$$(f \circ g)(x) = f(g(x))$$

$$= f(3 - x)$$

$$= \frac{(3 - x) + 6}{3}$$

$$= \frac{9 - x}{3}$$

$$(g \circ f)(x) = g(f(x))$$

$$= g\left(\frac{x + 6}{3}\right)$$

$$= 3 - \frac{x + 6}{3}$$

$$= \frac{9 - x - 6}{3}$$

$$= \frac{3 - x}{3}$$

$$\therefore f \circ g \neq g \circ f$$

$$(iv) \quad f(x) = 3 + x, g(x) = x - 4$$

$$(f \circ g)(x) = f(g(x))$$

$$= f(x - 4)$$

$$= 3 + (x - 4)$$

$$= x - 1$$

$$(g \circ f)(x) = g(f(x))$$

$$= g(3 + x)$$

$$= 3 + x - 4$$

$$= x - 1$$

$$\therefore f \circ g = g \circ f$$

$$(v) \quad f(x) = 4x^2 - 1, g(x) = 1 + x$$

$$(f \circ g)(x) = f(g(x))$$

$$= f(1 + x)$$

$$= 4(1 + x^2) - 1$$

$$= 4(1 + x^2 + 2x) - 1$$

$$= 4x^2 + 8x + 3$$

$$(g \circ f)(x) = g(f(x))$$

$$= g(4x^2 - 1)$$

$$= 1 + 4x^2 - 1$$

$$= 4x^2$$

$$\therefore f \circ g \neq g \circ f$$

2. Find the value of k , such that $f \circ g = g \circ f$.

$$(i) \quad f(x) = 3x + 2, g(x) = 6x - k$$

$$(ii) \quad f(x) = 2x - k, g(x) = 4x + 5$$

Solution :

$$(i) \quad f(x) = 3x + 2 ; g(x) = 6x - k$$

$$\text{Given } f \circ g = g \circ f$$

$$\Rightarrow (f \circ g)(x) = (g \circ f)(x)$$

$$\Rightarrow f(g(x)) = g(f(x))$$

$$\Rightarrow f(6x - k) = g(3x + 2)$$

$$\Rightarrow 3(6x - k) + 2 = 6(3x + 2) - k$$

$$\Rightarrow 18x - 3k + 2 = 18x + 12 - k$$

$$\Rightarrow -3k + 2 = 12 - k$$

$$\Rightarrow -2k = 10$$

$$\Rightarrow k = \frac{-10}{2} = -5$$

(ii) $f(x) = 2x - k$; $g(x) = 4x + 5$

$$\text{Given } f \circ g = g \circ f$$

$$\Rightarrow (f \circ g)(x) = (g \circ f)(x)$$

$$\Rightarrow f(g(x)) = g(f(x))$$

$$\Rightarrow f(4x + 5) = g(2x - k)$$

$$\Rightarrow 2(4x + 5) - k = 4(2x - k) + 5$$

$$\Rightarrow 8x + 10 - k = 8x - 4k + 5$$

$$\Rightarrow 10 - k = -4k + 5$$

$$\Rightarrow -k + 4k = 5 - 10$$

$$\Rightarrow 3k = -5$$

$$\Rightarrow k = \frac{-5}{3}$$

3. If $f(x) = 2x - 1$, $g(x) = \frac{x+1}{2}$, show that $f \circ g = g \circ f = x$.

Solution :

$$\text{Given } f(x) = 2x - 1, g(x) = \frac{x+1}{2}$$

$$\therefore f \circ g = (f \circ g)(x)$$

$$= f(g(x))$$

$$= f\left(\frac{x+1}{2}\right)$$

$$= 2\left(\frac{x+1}{2}\right) - 1$$

$$= x + 1 - 1$$

$$= x$$

$$g \circ f = (g \circ f)(x)$$

$$= g(f(x))$$

$$= g(2x - 1)$$

$$= \frac{2x - 1 + 1}{2}$$

$$= x$$

$$\therefore f \circ g = g \circ f = x$$

4. (i) If $f(x) = x^2 - 1$, $g(x) = x - 2$ find a , if $g \circ f(a) = 1$.

(ii) Find k , if $f(k) = 2k - 1$ and $f \circ f(k) = 5$.

Solution :

(i) $f(x) = x^2 - 1$, $g(x) = x - 2$

$$\text{Given } g \circ f(a) = 1$$

$$\Rightarrow g(f(a)) = 1$$

$$\Rightarrow a^2 - 1 - 2 = 1$$

$$\Rightarrow a^2 - 3 = 1$$

$$\Rightarrow a^2 = 4$$

$$\therefore a = \pm 2$$

(ii) $f(k) = 2k - 1$

$$\Rightarrow f \circ f(k) = 5$$

$$\Rightarrow f(f(k)) = 5$$

$$\Rightarrow f(2k - 1) = 5$$

$$\Rightarrow 2(2k - 1) - 1 = 5$$

$$\Rightarrow 4k - 2 = 6$$

$$\Rightarrow 4k = 8$$

$$\therefore k = 2$$

5. Let $A, B, C \subseteq \mathbb{N}$ and a function $f: A \rightarrow B$ be defined by $f(x) = 2x + 1$ and $g: B \rightarrow C$ be defined by $g(x) = x^2$. Find the range of $f \circ g$ and $g \circ f$.

Solution : $f : A \rightarrow B, g : B \rightarrow C$ where $A, B, C \subseteq \mathbb{N}$.

$$f(x) = 2x + 1, g(x) = x^2$$

Range of $f \circ g$:

$$(f \circ g)(x) = f(g(x))$$

$$= f(x^2)$$

$$= 2x^2 + 1$$

$$\therefore \text{Range of } f \circ g = \{y / y = 2x^2 + 1, x \in \mathbb{N}\}.$$

Range of $g \circ f$:

$$(g \circ f)(x) = g(f(x))$$

$$= g(2x + 1)$$

$$= (2x + 1)^2$$

$$\therefore \text{Range of } g \circ f = \{y / y = (2x + 1)^2, x \in \mathbb{N}\}.$$

 $o f$ **Solution :**

$$\text{Given } f(x) = x^2 - 1$$

$$\text{a) } f \circ f = ?$$

$$(f \circ f)(x) = f(f(x))$$

$$= f(x^2 - 1)$$

$$= (x^2 - 1)^2 - 1$$

$$= x^4 - 2x^2 + 1 - 1$$

$$= x^4 - 2x^2$$

$$\text{b) } f \circ f \circ f = ?$$

$$(f \circ f \circ f)(x) = f \circ f(f(x))$$

$$= f \circ f(x^2 - 1)$$

$$= (x^2 - 1)^4 - 2(x^2 - 1)^2$$

$$= (x^2 - 1)^2 [(x^2 - 1)^2 - 2]$$

$$= (x^4 - 2x^2 + 1)(x^4 - 2x^2 - 1)$$

$$= (x^4 - 2x^2)^2 - 1$$

$$((a + b)(a - b) = a^2 - b^2)$$

7. If $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ are defined by $f(x) = x^5$ and $g(x) = x^4$ then check if f, g are one-one and $f \circ g$ is one-one ?

Solution :

$$\text{Given } f(x) = x^5, g(x) = x^4$$

Let A be the domain.

B be the co-domain.

For every element $\in A$, there is a unique image in B. Since f is an odd function

$$\therefore f \text{ is } 1 - 1.$$

But $g(x)$ is an even function.

\therefore Two elements of domain will have the same image in co-domain.

$$\therefore g \text{ is not } 1 - 1.$$

8. Consider the functions $f(x), g(x), h(x)$ as given below. Show that $(f \circ g) \circ h = f \circ (g \circ h)$ in each case.

$$\text{(i) } f(x) = x - 1, g(x) = 3x + 1 \text{ and}$$

$$h(x) = x^2$$

$$\text{(ii) } f(x) = x^2, g(x) = 2x \text{ and } h(x) = x + 4$$

$$\text{(iii) } f(x) = x - 4, g(x) = x^2 \text{ and}$$

$$h(x) = 3x - 5$$

Solution :

$$\text{(i) Given } f(x) = x - 1, g(x) = 3x + 1, h(x) = x^2$$

$$\text{To verify : } (f \circ g) \circ h = f \circ (g \circ h)$$

$$(f \circ g)(x) = f(g(x))$$

$$= f(3x + 1)$$

$$= 3x + 1 - 1$$

$$= 3x$$

$$\therefore ((f \circ g) \circ h)(x) = (f \circ g)(h(x))$$

$$= (f \circ g)(x^2)$$

$$= 3x^2$$

$$\text{--- (1)}$$

$$(g \circ h)(x) = g(h(x))$$

$$= g(x^2)$$

$$= 3x^2 + 1$$

$$\therefore (f \circ (g \circ h))(x) = f((g \circ h)(x))$$

$$= f(3x^2 + 1)$$

$$= 3x^2 + 1 - 1$$

$$= 3x^2 \quad \text{--- (2)}$$

$$\therefore \text{From (1) and (2), } (f \circ g) \circ h = f \circ (g \circ h)$$

(ii) Given $f(x) = x^2$, $g(x) = 2x$, $h(x) = x + 4$

$$(f \circ g)(x) = f(g(x))$$

$$= f(2x)$$

$$= (2x)^2 = 4x^2$$

$$\therefore ((f \circ g) \circ h)(x) = (f \circ g)(h(x))$$

$$= (f \circ g)(x + 4)$$

$$= 4(x + 4)^2 \quad \text{--- (1)}$$

$$(g \circ h)(x) = g(h(x))$$

$$= g(x + 4)$$

$$= 2(x + 4)$$

$$= 2x + 8$$

$$\therefore (f \circ (g \circ h))(x) = f(2x + 8)$$

$$= (2x + 8)^2$$

$$= (2(x + 4))^2$$

$$= 4(x + 4)^2 \quad \text{--- (2)}$$

$$\therefore \text{From (1) and (2),}$$

$$(f \circ g) \circ h = f \circ (g \circ h).$$

(iii) Given $f(x) = x - 4$, $g(x) = x^2$, $h(x) = 3x - 5$

$$(f \circ g)(x) = f(g(x))$$

$$= f(x^2)$$

$$= x^2 - 4$$

$$\therefore ((f \circ g) \circ h)(x) = (f \circ g)(h(x))$$

$$= (f \circ g)(3x - 5)$$

$$= (3x - 5)^2 - 4 \quad \text{--- (1)}$$

$$(g \circ h)(x) = g(h(x))$$

$$= g(3x - 5)$$

$$= (3x - 5)^2$$

$$\therefore (f \circ (g \circ h))(x) = f(g \circ h)(x)$$

$$= f(3x - 5)^2$$

$$= (3x - 5)^2 - 4$$

$$\text{--- (2)}$$

$$\therefore \text{From (1) and (2),}$$

$$(f \circ g) \circ h = f \circ (g \circ h).$$

9. Let $f = \{(-1, 3), (0, -1), (2, -9)\}$ be a linear

Solution :

Given $f = \{(-1, 3), (0, -1), (2, -9)\}$ is a linear function from Z into Z .

$$\text{Let } y = ax + b$$

$$\text{When } x = -1, y = 3 \Rightarrow$$

$$3 = -a + b \quad \text{--- (1)}$$

$$\text{When } x = 0, y = -1 \Rightarrow -1 = 0 + b$$

$$\therefore b = -1$$

$$\therefore (1) \Rightarrow 3 = -a - 1$$

$$\Rightarrow a = -4$$

$$\therefore a = -4, b = -1$$

$$\therefore y = -4x - 1 \text{ is the required linear function.}$$

10. In electrical circuit theory, a circuit $C(t)$ is called a linear circuit if it satisfies the superposition principle given by $C(at_1 + bt_2) = aC(t_1) + bC(t_2)$, where a, b are constants. Show that the circuit $C(t) = 3t$ is linear.

Solution :

Given $C(t) = 3t$

To Prove : $C(t)$ is linear.

$$C(at_1) = 3at_1, C(bt_2) = 3bt_2$$

Adding,

$$C(at_1) + C(bt_2) = 3at_1 + 3bt_2 = 3(at_1 + bt_2)$$

which is the principle of super position.

$\therefore C(t) = 3t$ is a linear function.

EXERCISE 1.6**Multiple choice questions :**

1. If $n(A \times B) = 6$ and $A = \{1, 3\}$ then $n(B)$ is

(1) 1 (2) 2 (3) 3 (4) 6

Hint : **Ans : (3)**

$$n(A) = 2, n(A \times B) = 6 \Rightarrow n(B)$$

$$= n(A \times B) / n(A)$$

$$= 6/2 = 3$$

2. $A = \{a, b, p\}$, $B = \{2, 3\}$, $C = \{p, q, r, s\}$ then $n[(A \cup C) \times B]$ is

(1) 8 (2) 20 (3) 12 (4) 16

Hint : **Ans : (3)**

$$A \cup C = \{a, b, p, q, r, s\}, B = \{2, 3\}$$

$$n[(A \cup C) \times B] = 6 \times 2 = 12$$

3. If $A = \{1, 2\}$, $B = \{1, 2, 3, 4\}$, $C = \{5, 6\}$ and $D = \{5, 6, 7, 8\}$ then state which of the following statement is true ?

(1) $(A \times C) \subset (B \times D)$

(2) $(B \times D) \subset (A \times C)$

(3) $(A \times B) \subset (A \times D)$

(4) $(D \times A) \subset (B \times A)$

Hint : **Ans : (1)**

It is clearly $(A \times C) \subset (B \times D)$.

4. If there are 1024 relations from a set $A = \{1, 2, 3, 4, 5\}$ to a set B , then the number of elements in B is

(1) 3 (2) 2 (3) 4 (4) 8

Hint : **Ans : (2)**

$$n(A) = 5 = p$$

$$\therefore \text{No. of relations from } A \text{ to } B = 1024$$

$$\Rightarrow 2^{5q} = 1024$$

$$\Rightarrow (32)^q = (32)^2$$

$$\Rightarrow q = 2$$

$$\therefore n(B) = 2$$

5. The range of the relation $R = \{(x, x^2) \mid x \text{ is a prime number less than } 13\}$ is

(1) $\{2, 3, 5, 7\}$

(3) $\{4, 9, 25, 49, 121\}$

(4) $\{1, 4, 9, 25, 49, 121\}$

Hint : **Ans : (3)**

Prime no's less than 13 = $\{2, 3, 5, 7, 11\}$

$\therefore \text{Range of } R = \{4, 9, 25, 49, 121\},$

$$(\because R = \{(x, x^2)\})$$

6. If the ordered pairs $(a + 2, 4)$ and $(5, 2a + b)$ are equal then (a, b) is

(1) $(2, -2)$ (2) $(5, 1)$

(3) $(2, 3)$ (4) $(3, -2)$

Hint : **Ans : (4)**

$$a + 2 = 5, \quad 2a + b = 4$$

$$\Rightarrow a = 3 \quad 6 + b = 4$$

$$\Rightarrow b = -2$$

7. Let $n(A) = m$ and $n(B) = n$ then the total number of non-empty relations that can be defined from A to B is

(1) m^n (2) n^m (3) $2^{mn} - 1$ (4) 2^{mn}

Hint : **Ans : (4)**

Total no. of non-empty relations from A to B = $2^{n(A) \cdot n(B)} - 1 = 2^{mn} - 1$

8. If $\{(a, 8), (6, b)\}$ represents an identity function, then the value of a and b are respectively

(1) (8, 6) (2) (8, 8) (3) (6, 8) (4) (6, 6)

Hint : **Ans : (1)**

$(a, 8), (6, b) \Rightarrow$ identity function

$\therefore a = 8, b = 6$

9. Let $A = \{1, 2, 3, 4\}$ and $B = \{4, 8, 9, 10\}$. A function $f: A \rightarrow B$ given by $f = \{(1, 4), (2, 8), (3, 9), (4, 10)\}$ is a

(1) Many-one function
(2) Identity function
(3) One-to-one function
(4) Into function

Hint : **Ans : (3)**

Diff. elements of A have diff. images in B.

10. If $f(x) = 2x^2$ and $g(x) = \frac{1}{3x}$, then $f \circ g$ is

(1) $\frac{3}{2x^2}$ (2) $\frac{2}{3x^2}$ (3) $\frac{2}{9x^2}$ (4) $\frac{1}{6x^2}$

Hint : **Ans : (3)**

$(f \circ g)(x) = f(g(x))$

$$= f\left(\frac{1}{3x}\right)$$

$$= 2\left(\frac{1}{3x}\right)^2$$

$$= \frac{2}{9x^2}$$

11. If $f: A \rightarrow B$ is a bijective function and if $n(B) = 7$, then $n(A)$ is equal to

(1) 7 (2) 49 (3) 1 (4) 14

Hint : **Ans : (1)**

$f: A \rightarrow B$ is bijective (1 - 1 and onto) and $n(B) = 7 \therefore n(A) = 7$

12. Let f and g be two functions given by
 $f = \{(0, 1), (2, 0), (3, -4), (4, 2), (5, 7)\}$
 $g = \{(0, 2), (1, 0), (2, 4), (-4, 2), (7, 0)\}$
 then the range of $f \circ g$ is

(1) $\{0, 2, 3, 4, 5\}$ (2) $\{-4, 1, 0, 2, 7\}$

(3) $\{1, 2, 3, 4, 5\}$ (4) $\{0, 1, 2\}$

Hint : **Ans : (4)**

$$(f \circ g)(0) = f(g(0)) = f(2) = 0$$

$$(f \circ g)(1) = f(g(1)) = f(0) = 1$$

$$(f \circ g)(2) = f(g(2)) = f(4) = 2$$

$$(f \circ g)(7) = f(g(7)) = f(0) = 1$$

$$\therefore \text{Range} = \{0, 1, 2\}$$

13. Let $f(x) = \sqrt{1+x^2}$ then

$$(1) f(xy) = f(x) \cdot f(y)$$

$$(2) f(xy) \geq f(x) \cdot f(y)$$

$$(3) f(xy) \leq f(x) \cdot f(y)$$

$$(4) \text{None of these}$$

Hint : **Ans : (3)**

$$f(x) = \sqrt{1+x^2}$$

$$\therefore f(y) = \sqrt{1+y^2}$$

$$\therefore f(xy) = \sqrt{1+x^2y^2}$$

$$f(x) \cdot f(y) = \sqrt{(1+x^2)(1+y^2)}$$

$$= \sqrt{1+x^2+y^2+x^2y^2}$$

$$\geq \sqrt{1+x^2y^2}$$

$$\geq f(xy)$$

$$\therefore f(xy) \leq f(x) \cdot f(y)$$

14. If $g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$ is a function given by $g(x) = \alpha x + \beta$ then the values of α and β are

- (1) $(-1, 2)$ (2) $(2, -1)$
 (3) $(-1, -2)$ (4) $(1, 2)$

Hint :

Ans : (3)

$$g(x) = \alpha x + \beta$$

$$\Rightarrow 1 = \alpha + \beta, 3 = 2\alpha + \beta, 5 = 3\alpha + \beta$$

$$\text{on Subtracting, } \alpha = 2 \Rightarrow \beta = -1$$

15. $f(x) = (x+1)^3 - (x-1)^3$ represents a function which is

- (1) linear (2) cubic
 (3) reciprocal (d) quadratic

Hint :

Ans : (4)

$$\begin{aligned} f(x) &= (x+1)^3 - (x-1)^3 \\ &= (x^3 + 3x^2 + 3x + 1) - (x^3 - 3x^2 + 3x - 1) \\ &= 6x^2 + 2, \text{ a quadratic function.} \end{aligned}$$

UNIT EXERCISE - 1

1. If the ordered pairs $(x^2 - 3x, y^2 + 4y)$ and $(-2, 5)$ are equal, then find x and y .

Solution :

$$\text{Given } (x^2 - 3x, y^2 + 4y) = (-2, 5)$$

$$\begin{array}{l|l} \therefore x^2 - 3x = -2 & y^2 + 4y = 5 \\ \Rightarrow x^2 - 3x + 2 = 0 & y^2 + 4y - 5 = 0 \\ \Rightarrow (x-2)(x-1) = 0 & (y+5)(y-1) = 0 \\ \therefore x = 2, 1 & y = -5, 1 \end{array}$$

2. The Cartesian product $A \times A$ has 9 elements among which $(-1, 0)$ and $(0, 1)$ are found. Find the set A and the remaining elements of $A \times A$.

Solution :

$$\text{Given } n(A \times A) = 9 \text{ and } (-1, 0), (0, 1) \in A \times A$$

$\therefore A = \{-1, 0, 1\}$ and the remaining elements of

$$A \times A = \{(-1, -1), (-1, 1), (0, -1), (0, 0), (1, -1), (1, 0), (1, 1)\}$$

3. Given that $f(x) = \begin{cases} \sqrt{x-1} & x \geq 1 \\ 4 & x < 1 \end{cases}$. Find
 (i) $f(0)$ (ii) $f(3)$ (iii) $f(a+1)$
 in terms of a . (Given that $a \geq 0$)

Solution :

$$\text{Given } f(x) = \begin{cases} \sqrt{x-1} & x \geq 1 \\ 4 & x < 1 \end{cases}$$

$$i) f(0) = 4$$

$$ii) f(3) = \sqrt{3-1} = \sqrt{2}$$

$$\begin{aligned} iii) f(a+1) &= \sqrt{a+1-1} \quad (\because a \geq 0) \\ &= \sqrt{a} \quad \Rightarrow a+1 \geq 1 \end{aligned}$$

4. Let $A = \{9, 10, 11, 12, 13, 14, 15, 16, 17\}$ and let $f: A \rightarrow N$ be defined by $f(n) =$ the highest prime factor of $n \in A$. Write f as

of f .

Solution :

$$\text{Given } A = \{9, 10, 11, 12, 13, 14, 15, 16, 17\}$$

$f: A \rightarrow N$ defined by $f(n) =$ highest prime factor of $n \in A$.

$$f(9) = 3 (\because 3 \text{ is the highest prime factor of } 9)$$

$$f(10) = 5 (\because 10 = 5 \times 2)$$

$$f(11) = 11 (\because 11 \text{ is a prime no.)}$$

$$f(12) = 3 (\because 12 = 3 \times 2 \times 2)$$

$$f(13) = 13 (\because 13 \text{ is a prime no.)}$$

$$f(14) = 7 (\because 14 = 7 \times 2)$$

$$f(15) = 5 (\because 15 = 5 \times 3)$$

$$f(16) = 2 (\because 16 = 2 \times 2 \times 2 \times 2)$$

$$f(17) = 17 (\because 17 \text{ is a prime number})$$

$$f = \{(9, 3), (10, 5), (11, 11), (12, 3), (13, 13), (14, 7), (15, 5), (16, 2), (17, 17)\}$$

$$\text{Range of } f = \{2, 3, 5, 7, 11, 13, 17\}$$

5. Find the domain of the function

$$f(x) = \sqrt{1 + \sqrt{1 - \sqrt{1 - x^2}}}.$$

Solution :

$$\text{Given } f(x) = \sqrt{1 + \sqrt{1 - \sqrt{1 - x^2}}}$$

If $x > 1$ and $x < -1$, $f(x)$ leads to unreal

$$\therefore \text{The domain of } f(x) = \{-1, 0, 1\}$$

6. If $f(x) = x^2$, $g(x) = 3x$ and $h(x) = x - 2$. Prove that $(f \circ g) \circ h = f \circ (g \circ h)$.**Solution :**

$$\text{Given } f(x) = x^2, g(x) = 3x, \quad h(x) = x - 2$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) \\ &= f(3x) \\ &= (3x)^2 = 9x^2 \end{aligned}$$

$$\begin{aligned} ((f \circ g) \circ h)(x) &= (f \circ g)(h(x)) \\ &= (f \circ g)(x - 2) \end{aligned}$$

$$\begin{aligned} (g \circ h)(x) &= g(h(x)) \\ &= g(x - 2) \\ &= 3(x - 2) = 3x - 6 \end{aligned}$$

$$\begin{aligned} \therefore (f \circ (g \circ h))(x) &= f(3x - 6) \\ &= (3x - 6)^2 \\ &= (3(x - 2))^2 \\ &= 9(x - 2)^2 \quad \text{--- (2)} \end{aligned}$$

\therefore From (1) and (2),
 $(f \circ g) \circ h = f \circ (g \circ h)$.

7. Let $A = \{1, 2\}$ and $B = \{1, 2, 3, 4\}$, $C = \{5, 6\}$ and $D = \{5, 6, 7, 8\}$. Verify whether $A \times C$ is a subset of $B \times D$?**Solution :**

Given $A = \{1, 2\}$, $B = \{1, 2, 3, 4\}$, $C = \{5, 6\}$,
 $D = \{5, 6, 7, 8\}$

$$A \times C = \{(1, 5), (1, 6), (2, 5), (2, 6)\}$$

$$\begin{aligned} B \times D = \{ & (1, 5), (1, 6), (1, 7), (1, 8), (2, 5), \\ & (2, 6), (2, 7), (2, 8), (3, 5), (3, 6), (3, 7), (3, 8), \\ & (4, 5), (4, 6), (4, 7), (4, 8) \}. \end{aligned}$$

Clearly $A \times C$ is a subset of $B \times D$.**8. If $f(x) = \frac{x-1}{x+1}$, $x \neq -1$ show that $f(f(x)) = -\frac{1}{x}$, provided $x \neq 0$.****Solution :**

$$\text{Given } f(x) = \frac{x-1}{x+1}$$

$$f(f(x)) = f\left(\frac{x-1}{x+1}\right)$$

$$\begin{aligned} &= \frac{\frac{x-1}{x+1} - 1}{\frac{x-1}{x+1} + 1} = \frac{\frac{x-1-x-1}{x+1}}{\frac{x-1+x+1}{x+1}} \\ &= \frac{-2}{2x} = \frac{-1}{x} \end{aligned}$$

9. The functions f and g are defined by

$$f(x) = 6x + 8; g(x) = \frac{x-2}{3}$$

(i) Calculate the value of $gg\left(\frac{1}{2}\right)$ **(ii) Write an expression for $gf(x)$ in its simplest form.****Solution :**

$$\text{Given } f(x) = 6x + 8, g(x) = \frac{x-2}{3}$$

$$i) \quad gg\left(\frac{1}{2}\right) = g\left(g\left(\frac{1}{2}\right)\right)$$

$$= g\left(\frac{\frac{1}{2} - 2}{3}\right)$$

$$= g\left(\frac{-3}{6}\right)$$

$$= g\left(\frac{-1}{2}\right)$$

$$= \frac{-\frac{1}{2} - 2}{3} = \frac{-5}{6}$$

$$\begin{aligned}
 \text{ii) } gf(x) &= g(f(x)) \\
 &= g(6x+8) \\
 &= \frac{6x+8-2}{3} \\
 &= \frac{6x+6}{3} \\
 &= 2x+2 \\
 &= 2(x+1)
 \end{aligned}$$

10. Write the domain of the following real functions

$$\text{(i) } f(x) = \frac{2x+1}{x-9} \quad \text{(ii) } p(x) = \frac{-5}{4x^2+1}$$

$$\text{(iii) } g(x) = \sqrt{x-2} \quad \text{(iv) } h(x) = x+6$$

Solution :

$$\text{i) } f(x) = \frac{2x+1}{x-9}$$

$$\text{If } x = 9, f(x) \rightarrow \infty$$

The domain is $\mathbb{R} - \{9\}$

$$\text{ii) } f(x) = \frac{-5}{4x^2+1}$$

$$\text{If } x \rightarrow \infty, 4x^2+1 \text{ does not tend to } \infty.$$

\therefore The domain is \mathbb{R} .

$$\text{iii) } g(x) = \sqrt{x-2}$$

The function exists only if $x \geq 2$

\therefore The domain is $[2, \infty)$.

$$\text{iv) } h(x) = x+6$$

$h(x)$ exists for all real numbers.

\therefore The domain is \mathbb{R} .

PROBLEMS FOR PRACTICE

1. If $A \times B = \{(a, x), (a, y), (b, x), (b, y)\}$, then find A and B.

(Ans : $A = \{a, b\}, B = \{x, y\}$)

2. If $A = \{1, 2, 3\}, B = \{3, 4\}, C = \{4, 5, 6\}$, find i) $A \times (B \cap C)$ (ii) $(A \times B) \cap (A \times C)$.

(Ans : $\{(1, 4), (2, 4), (3, 4)\}$)

3. Find the Cartesian product of 3 sets.

$$A = \{1, 2\}, B = \{3, 4\}, C = \{x : x \in \mathbb{N} \text{ and } 4 \leq x \leq 6\}.$$

{Ans : $\{(1, 3, 4), (1, 3, 5), (1, 3, 6), (1, 4, 4), (1, 4, 5), (1, 4, 6), (2, 3, 4), (2, 3, 5), (2, 3, 6), (2, 4, 4), (2, 4, 5), (2, 4, 6)\}$ }

then verify $A \times (B \cup C) = (A \times B) \cup (A \times C)$.

5. If $n(P) = 3, n(Q) = 4, (r, 4), (g, 1), (w, 3), (r, 9)$ are in $P \times Q$, find P and Q and also the remaining elements of $P \times Q$.

(Ans : $(r, 1), (r, 3), (g, 3), (g, 4), (g, 9), (w, 1), (w, 4), (w, 9)$)

6. Find x and y if $(x+3, 5) = (6, 2x+y)$.

(Ans : $3, -1$)

7. If $A = \{1, 2, 3, 4, 5, 6\}, R = \{(a, b) \mid b \text{ is exactly divisible by } a\}$. Find the range of R.

(Ans : $\{1, 2, 3, 4, 5, 6\}$)

8. If the ordered pairs $(x, -1), (5, y)$ belong to the set $\{(a, b) \mid b = 2a - 3\}$, find x and y.

(Ans : $1, 7$)

9. Express $R = \{(x, y) \mid x^2 + y^2 = 25, x, y \in \mathbb{N}\}$.

(Ans : $(0, 5), (3, 4), (4, 3), (5, 0)$)

10. If $n(A) = 3$ and $B = \{2, 3, 4, 6, 7, 8\}$, find the number of relations from A to B.

(Ans : 2^{18})

11. Find the domain of the following functions:

(i) $f(x) = \frac{1}{\sqrt{x-5}}$ (ii) $g(x) = \frac{3}{2-x^2}$

(Ans : i) $(5, \infty)$ ii) $\mathbb{R} - \{-\sqrt{2}, \sqrt{2}\}$

12. If A and B are 2 sets having 3 elements in common. If $n(A) = 5$, $n(B) = 4$, find $n(A \times B)$ and $n[(A \times B) \cap (B \times A)]$.

(Ans : 20, 9)

13. Which of the following relations are functions and write them in ordered pairs also.

ii) $\{(x, y) : y > 2x+1, x \in \{1, 2\}, y \in \{2, 4, 6\}\}$.

(Ans : (i) a function (ii) not a function)

14. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = 2x^3 - 5$, show that f is bijective.

15. Prove that the function $f: \mathbb{N} \rightarrow \mathbb{N}$ defined by $f(x) = x^2 + x + 1$ is one-one but not onto.

16. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = x^2 - 3x + 2$, find $f \circ f$.

(Ans : $x^4 - 6x^3 + 10x^2 - 3x$)

17. Let $p(x) = \begin{cases} 4x-5, & x \geq 2 \\ 1-x, & x < 2 \end{cases}$,

$q(x) = \begin{cases} 3x+7 & x \leq 3 \\ 7 & x > 3 \end{cases}$

are functions from $\mathbb{R} \rightarrow \mathbb{R}$, find

(i) $(p \circ q)(3)$ (ii) $(q \circ p)(10)$.

(Ans : 59, 7)

18. If $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = (3-x^3)^{1/3}$, find $(f \circ f)(x)$.

(Ans : x)

19. If $f(x) = \sqrt{x-2}$, find $(f \circ f \circ f)(38)$.

(Ans : 0)

20. A function $f: [-3, 7) \rightarrow \mathbb{R}$ is defined by

$$f(x) = \begin{cases} 4x^2 - 1, & -3 \leq x < 2 \\ 3x - 2, & 2 \leq x \leq 4 \\ 2x - 3, & 4 < x < 7 \end{cases}$$

Find (i) $\frac{f(3)+f(-1)}{2f(6)-f(1)}$ (ii) $f(-2)-f(4)$

(Ans : $\frac{2}{3}, 5$)

21. The following table represents a function from $A = \{5, 6, 8, 10\}$ to $B = \{19, 15, 9, 11\}$ where $f(x) = 2x - 1$. Find a and b .

x	5	6	8	10
$f(x)$	a	11	6	19

(Ans : $a = 9, b = 15$)

22. A function $f: [1, 6) \rightarrow \mathbb{R}$ is defined as follows :

$$f(x) = \begin{cases} 1+x, & 1 \leq x < 2 \\ 2x-1, & 2 \leq x < 4 \\ 3x^2-10, & 4 \leq x < 6 \end{cases}$$

Find (i) $f(2)-f(4)$ (ii) $2f(5)-3f(1)$

(Ans : -35, 124)

23. If $f(x) = lx + 5$, $g(x) = 2x + 3$, find l such that $f \circ g = g \circ f$.

(Ans : $l = 8$)

24. Given $f(x) = x + \frac{1}{2}$, $g(x) = \frac{1}{2}x + 3$, $h(x) = x^2$, verify that $(f \circ g) \circ h = f \circ (g \circ h)$.

OBJECTIVE TYPE QUESTIONS

1. $X = \{8^n - 7n - 1 / n \in \mathbb{N}\}$, $Y = \{49N - 49 / n \in \mathbb{N}\}$, then

(a) $X \subset Y$ (b) $Y \subset X$
 (c) $X = Y$ (d) $X \cap Y = \phi$.

Ans : (c)

2. If $A = \{a, b\}$, $B = \{c, d\}$, $C = \{d, e\}$, then $\{(a, c), (a, d), (a, e), (b, c), (b, d), (b, e)\}$ is equal to

(a) $A \cap (B \cup C)$ (b) $A \cup (B \cap C)$
 (c) $A \times (B \cup C)$ (d) $A \times (B \cap C)$

Ans : (c)

3. If $n(A) = 4$, $n(B) = 3$, $n(A \times B \times C) = 24$, then $n(C) =$

Ans : (c)

4. If $n(A) = 4$, $n(B) = 5$, $n(A \cap B) = 3$, then $n((A \times B) \cap (B \times A))$ is

(a) 8 (b) 9 (c) 10 (d) 11

Ans : (b)

5. $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x + \sqrt{x^2}$ is

(a) 1 - 1 (b) onto
 (c) bijective (d) none of these

Ans : (d)

6. Which of the following functions from \mathbb{Z} to itself are bijections ?

(a) $f(x) = x^3$ (b) $f(x) = x + 2$
 (c) $f(x) = 2x + 1$ (d) $f(x) = x^2 + x$

Ans : (b)

7. If $f(x) = \frac{1}{x^2}$, $g(x) = 1 + x$, then $g \circ f$ is

(a) $(1 + x)^2$ (b) $\frac{x^2}{x^2 + 1}$
 (c) $\frac{1}{(1 + x)^2}$ (d) $\frac{x^2 + 1}{x^2}$

Ans : (d)

8. $f: X \rightarrow Y$ where $X = \{-1, -2, -3\}$, $Y = \{3, 4, 5\}$ is given by $f(x) = x + 6$, $x \in X$, then f is

(a) onto (b) many to one
 (c) constant function (d) bijective

Ans : (d)

9. The range of $f: \mathbb{Z} \rightarrow \mathbb{Z}$ given by $f(x) = x^2$ is

(a) \mathbb{Q} (b) \mathbb{Z} (c) \mathbb{W} (d) \mathbb{N}

Ans : (c)

10. The value of a in order that $\{(4, 3), (7, a)\}$ may represent a constant function is

(a) 7 (b) 4
 (c) 3 (d) none of these

Ans : (c)

11. The co-domain of $f: A \rightarrow \mathbb{R}$ where $A = \{2, 3, 5\}$, $f(x) = 2x - 1$ can be taken as

(a) $\{3, 5, 7\}$ (b) $\{3, 5, 9, 11\}$
 (c) $\{5, 9\}$ (d) $\{3, 5\}$

Ans : (b)

CHAPTER 2

NUMBERS AND SEQUENCES

I. EUCLID'S DIVISION LEMMA AND ALGORITHM

Key Points

- ✓ Let a and b ($a > b$) be any two positive integers. Then, there exist unique integers q and r such that $a = bq + r$, $0 \leq r < b$. (Euclid's Division Lemma)
The remainder is always less than the divisor.
If $r = 0$ then $a = bq$ so b divides a .
Similarly, if b divides a then $a = bq$.
- ✓ If a and b are any two integers then there exist unique integers q and r such that $a = bq + r$,
- ✓ If a and b are positive integers such that $a = bq + r$, then every common divisor of a and b is a common divisor of b and r and vice-versa. (Euclid's Division Algorithm)
- ✓ If a, b are two positive integers with $a > b$ then G.C.D of $(a, b) = \text{GCD of } (a - b, b)$.
- ✓ Two positive integers are said to be relatively prime or co prime if their. Highest Common Factor is 1.

Example 2.2

Find the quotient and remainder when a is divided by b in the following cases (i) $a = -12, b = 5$
(ii) $a = 17, b = -3$ (iii) $a = -19, b = -4$.

Solution :

(i) $a = -12, b = 5$

By Euclid's division lemma

$$a = bq + r, \text{ where } 0 \leq r < |b|$$

$$-12 = 5 \times (-3) + 3 \quad 0 \leq r < |5|$$

Therefore, Quotient $q = -3$, Remainder $r = 3$.

(ii) $a = 17 \quad b = -3$

By Euclid's division lemma

$$a = bq + r, \text{ where } 0 \leq r < |b|$$

$$17 = (-3) \times (-5) + 2, \quad 0 \leq r < |-3|$$

Therefore, Quotient $q = -5$, Remainder $r = 2$.

(iii) $a = -19 \quad b = -4$

By Euclid's division lemma

$$a = bq + r, \text{ where } 0 \leq r < |b|$$

$$-19 = (-4) \times (5) + 1, \quad 0 \leq r < |-4|$$

Therefore, Quotient $q = 5$, Remainder $r = 1$.

Example 2.3

Show that the square of an odd integer is of the form $4q + 1$, for some integer q .

Solution :

Let x be any odd integer. Since any odd integer is one more than an even integer, we have $x = 2k + 1$, for some integer k .

$$\begin{aligned}x^2 &= (2k + 1)^2 \\&= 4k^2 + 4k + 1 \\&= 4k(k + 1) + 1 \\&= 4q + 1,\end{aligned}$$

where $q = k(k + 1)$ is some integer.

Example 2.4

If the Highest Common Factor of 210 and 55 is expressible in the form $55x - 325$, find x .

Solution :

Using Euclid's Division Algorithm, let us find the HCF of given numbers

$$\begin{aligned}210 &= 55 \times 3 + 45 \\55 &= 45 \times 1 + 10 \\45 &= 10 \times 4 + 5 \\10 &= 5 \times 2 + 0\end{aligned}$$

The remainder is zero.

So, the last divisor 5 is the Highest Common Factor (HCF) of 210 and 55.

Since, HCF is expressible in the form $55x - 325 = 5$

$$\text{gives } 55x = 330$$

$$\text{Hence } x = 6$$

Example 2.5

Find the greatest number that will divide 445 and 572 leaving remainders 4 and 5 respectively.

Solution :

Since the remainders are 4, 5 respectively the required number is the HCF of the number $445 - 4 = 441$, $572 - 5 = 567$.

Hence, we will determine the HCF of 441 and 567. Using Euclid's Division Algorithm, we have

$$\begin{aligned}567 &= 441 \times 1 + 126 \\441 &= 125 \times 3 + 63 \\126 &= 63 \times 2 + 0\end{aligned}$$

Therefore HCF of 441, 567 = 63 and so the required number is 63.

Example 2.6

Find the HCF of 396, 504, 636.

Solution :

Using Euclid's division algorithm we get

$$504 = 396 \times 1 + 108$$

The remainder is $108 \neq 0$

Again applying Euclid's division algorithm

$$396 = 108 \times 3 + 72$$

The remainder is $72 \neq 0$,

Again applying Euclid's division algorithm

$$108 = 72 \times 1 + 36$$

The remainder is $36 \neq 0$,

Again applying Euclid's division algorithm

$$72 = 36 \times 2 + 0$$

Here the remainder is zero. Therefore HCF of 396, 504 = 36.

To find the HCF of 636 and 36.

Using Euclid's division algorithm we get

$$636 = 36 \times 17 + 24$$

The remainder is $24 \neq 0$

Again applying Euclid's division algorithm
 $36 = 24 \times 1 + 12$

The remainder is $12 \neq 0$

Again applying Euclid's division algorithm
 $24 = 12 \times 2 + 0$

Here the remainder is zero. Therefore HCF
 of 636, $36 = 12$

Therefore Highest Common Factor of 396,
 504 and 636 is 12.

EXERCISE 2.1

1. Find all positive integers, when divided
 by 3 leaves remainder 2.

Solution:

To find all positive integers which when
 divided by 3 leaves remainder 2.

$$\text{i.e.} \quad a \equiv 2 \pmod{3}$$

$$\Rightarrow a - 2 \text{ is divisible by } 3$$

$$\Rightarrow a = 3K + 2$$

$\therefore a$ takes the following values

$a = 2, 5, 8, 11, 14, \dots$ when $K = 0, 1, 2, 3, 4, \dots$

2. A man has 532 flower pots. He wants to
 arrange them in rows such that each row
 contains 21 flower pots. Find the number
 of completed rows and how many flower
 pots are left over

Solution :

No. of flower pots = 532

All pots to be arranged in rows

& each row to contain 21 flower pots.

$$\therefore 532 = 21q + r$$

$$\begin{array}{r} 25 \\ 21 \overline{) 532} \\ \underline{42} \\ 112 \\ \underline{105} \\ 7 \end{array}$$

$$\Rightarrow 532 = 21 \times 25 + 7$$

\therefore Number of completed rows = 25

Number of flower pots left out = 7

3. Prove that the product of two consecu-
 tive positive integers is divisible by 2.

Solution :

Let the 2 consecutive positive integers be

$$x, x + 1$$

$$= x^2 + x$$

Case (i)

If x is an even number

$$\text{Let } x = 2k$$

$$\therefore x^2 + x = (2k)^2 + 2k$$

$$= 2k(2k + 1), \text{ divisible by } 2$$

Case (ii)

If x is an odd number,

$$\text{Let } x = 2k + 1$$

$$x^2 + x = (2k + 1)^2 + (2k + 1)$$

$$= 4k^2 + 4k + 1 + 2k + 1$$

$$= 4k^2 + 6k + 2$$

$$= 2(2k^2 + 3k + 2) \text{ divisible by } 2$$

\therefore Product of 2 consecutive positive inte-
 gers is divisible by 2.

4. When the positive integers a , b and c are divided by 13, the respective remainders are 9, 7 and 10. Show that $a + b + c$ is divisible by 13.

Solution :

When a is divided by 13, remainder is 9

$$\text{i.e., } a = 13q + 9 \quad \dots\dots (1)$$

When b is divided by 13, remainder is 7

$$\text{i.e., } b = 13q + 7 \quad \dots\dots (2)$$

When c is divided by 13, remainder is 11

$$\text{i.e., } c = 13q + 11 \quad \dots\dots (3)$$

Adding (1), (2) & (3)

$$\begin{aligned} a + b + c &= 39q + 26 \\ &= 13(2q + 2) \end{aligned}$$

Hence proved.

5. Prove that square of any integer leaves the remainder either 0 or 1 when divided by 4.

Solution :

Case (i)

Let x be the even number

$$\text{Let } x = 2n$$

$$x^2 = (2n)^2$$

$= 4n^2$, which leaves remainder 0 when divided by 4.

Case (ii)

Let x be the odd number

$$\therefore \text{Let } x = 2n + 1$$

$$x^2 = (2n + 1)^2$$

$$= 4n^2 + 4n + 1$$

$= 4(n^2 + n) + 1$ leaves remainder 1 when divided by 4.

Hence proved.

6. Use Euclid's Division Algorithm to find the Highest Common Factor (HCF) of

(i) 340 and 412

(ii) 867 and 255

(iii) 10224 and 9648

(iv) 84, 90 and 120

Solution :

- i) HCF of 340, 412 by Euclid's algorithm.

First we should divide 412 by 340.

$$412 = 340 \times 1 + 72$$

$$340 = 72 \times 4 + 52$$

$$72 = 52 \times 1 + 20$$

$$52 = 20 \times 2 + 12$$

$$20 = 12 \times 1 + 8$$

$$8 = (4) \times 2 + 0$$

\therefore The last divisor "4" is the HCF.

- ii) HCF of 867 and 255.

$$867 = 255 \times 3 + 102$$

$$255 = 102 \times 2 + 51$$

$$102 = (51) \times 2 + 0$$

\therefore The last divisor "51" is the HCF.

- iii) HCF of 10224 and 9648

$$10224 = 9648 \times 1 + 576$$

$$9648 = 576 \times 16 + 432$$

$$576 = 432 \times 1 + 144$$

$$432 = (144) \times 3 + 0$$

\therefore The last divisor "144" is the HCF.

- iv) HCF of 84, 90, and 120

First we find HCF of 84 & 90

$$90 = 84 \times 1 + 6$$

$$84 = (6) \times 14 + 0$$

\therefore HCF of 84, 90 is 6

Next, we find HCF of 120 and 6.

$$120 = 6 \times 20 + 0$$

\therefore HCF of 84, 90 and 120 = 6

- 7. Find the largest number which divides 1230 and 1926 leaving remainder 12 in each case.**

Solution :

To find the largest number which divides 1230 and 1926 leaving remainder 12

i.e., HCF of 1230 - 12 and 1926 - 12

i.e., HCF of 1218 and 1914

First we divide 1914 by 1218

$$1218 = 696 \times 1 + 522$$

$$696 = 522 \times 1 + 174$$

$$522 = (174) \times 3 + 0$$

\therefore HCF = 174

\therefore The required largest number = 174.

- 8. If d is the Highest Common Factor of 32 and 60, find x and y satisfying $d = 32x + 60y$.**

Solution :

Given d is the HCF of 32 and 60

$$\therefore 60 = 32 \times 1 + 28 \quad \dots\dots\dots (1)$$

$$32 = 28 \times 1 + 4 \quad \dots\dots\dots (2)$$

$$28 = (4) \times 7 + 0$$

$\therefore d = 4$

$$= 32 - (28 \times 1) \quad [\text{From (2)}]$$

$$= 32 \times 1 - (60 - 32) \quad [\text{From (1)}]$$

$$= 32 - (1 \times 60 - 1 \times 32)$$

$$= 32 - 1 \times 60 + 1 \times 32$$

$$= 32(1 + 1) + 60(-1)$$

$$d = 32(2) + 60(-1)$$

$$\Rightarrow 32x + 60y = 32(2) + 60(-1)$$

$$\therefore x = 2, y = -1$$

- 9. A positive integer when divided by 88 gives the remainder 61. What will be the remainder when the same number is divided by 11?**

Solution :

The standard form is $a = bq + r$

$$\Rightarrow a = 88q + 61$$

$$a = (11 \times 8q) + (55 + 6)$$

$$a = 11[8q + 5] + 6$$

\therefore When the same positive integer is divided by 11 the remainder is 6.

- 10. Prove that two consecutive positive integers are always coprime.**

Solution :

Let $x, x + 1$ be two consecutive integers.

Let G.C.D. of $(x, x + 1)$ be ' n '

Then ' n ' divides $x + 1 - x$

i.e., $n = 1$

G.C.D. of $(x, x + 1) = 1$

x & $x + 1$ are Co-prime.

II. FUNDAMENTAL THEOREM OF ARITHMETIC :


Key Points

- ✓ Every natural number except 1 can be factorized as a product of primes and this factorization is unique except for the order in which the prime factors are written.
- ✓ If a prime number p divides ab then either p divides a or p divides b . That is p divides at least one of them.
- ✓ If a composite number n divides ab , then n neither divide a nor b . For example, 6 divides 4×3 but 6 neither divide 4 nor 3.

Example 2.7

In the given factor tree, find the numbers m and n .

Solution :



Value of the first box from bottom = $5 \times 2 = 10$

Value of $n = 5 \times 10 = 50$

Value of the second box from bottom

$$= 3 \times 50 = 150$$

Value of $m = 2 \times 150 = 300$

Thus, the required numbers are $m = 300$, $n = 50$.

Example 2.8

Can the number 6^n , n being a natural number end with the digit 5 ? Give reason for your answer.

Solution :

$$\text{Since } 6^n = (2 \times 3)^n = 2^n \times 3^n,$$

2 is a factor of 6^n . So, 6^n is always even.

But any number whose last digit is 5 is always odd.

Hence, 6^n cannot end with the digit 5.

Example 2.9

Is $7 \times 5 \times 3 \times 2 + 3$ a composite number ? Justify your answer.

Solution :

Yes, the given number is a composite num-

$$7 \times 5 \times 3 \times 2 + 3 = 3 \times (7 \times 5 \times 2 + 1) = 3 \times 71$$

Since the given number can be factorized in terms of two primes, it is a composite number.

Example 2.10

' a ' and ' b ' are two positive integers such that $a^b \times b^a = 800$. Find ' a ' and ' b '.

Solution :

The number 800 can be factorized as

$$800 = 2 \times 2 \times 2 \times 2 \times 2 \times 5 \times 5 = 2^5 \times 5^2$$

$$\text{Hence, } a^b \times b^a = 2^5 \times 5^2$$

This implies that $a = 2$ and $b = 5$ (or) $a = 5$ and $b = 2$.

EXERCISE 2.2

1. For what values of natural number n , 4^n can end with the digit 6?

Solution :

$$\begin{aligned} 4^n &= (2 \times 2)^n \\ &= 2^n \times 2^n \end{aligned}$$

Since 2 is a factor of 4, 4^n is always even.

\therefore If 4^n is to be end with 6, n should be even.

\therefore For the even powers of ' n ', 4^n will be ended with even no.

2. If m, n are natural numbers, for what values of m , does $2^n \times 5^m$ ends in 5?

Solution :

$n \quad m$

Since m and n are natural, numbers and 2^n is even.

for any value of m , $2^n \times 5^m$ will not be ended in 5.

\therefore No value of m will make x true.

3. Find the HCF of 252525 and 363636.

Solution :

Given numbers are 252525, 363636

By using prime factorization

5	252525	2	363636
5	50505	2	181818
3	10101	3	90909
7	3367	3	30303
	481	3	10101
		7	3367
			481

$$\therefore 252525 = 5 \times 5 \times \underline{3} \times \underline{7} \times \underline{481}$$

$$363636 = 2 \times 2 \times \underline{3} \times 3 \times 3 \times \underline{7} \times \underline{481}$$

$$\begin{aligned} \therefore \text{HCF} &= 3 \times 7 \times 481 \\ &= 10,101 \end{aligned}$$

4. If $13824 = 2^a \times 3^b$ then find a and b .

Solution :

Given $2^a \times 3^b = 13824$

$$\Rightarrow 2^a \times 3^b = 2^9 \times 3^2$$

$$\therefore a = 9, b = 2$$

2	13824
2	6912
2	3456
2	1728
2	864
2	432
2	216
2	54
3	27
3	3

5. If $p_1^{x_1} \times p_2^{x_2} \times p_3^{x_3} \times p_4^{x_4} = 113400$ where p_1, p_2, p_3, p_4 are primes in ascending order and x_1, x_2, x_3, x_4 are integers, find the value of p_1, p_2, p_3, p_4 and x_1, x_2, x_3, x_4 .

Solution :

$$p_1^{x_1} \times p_2^{x_2} \times p_3^{x_3} \times p_4^{x_4} = 113400$$

2	113400
2	56700
2	28350
3	14175
3	4725
3	1575
3	525
5	175
5	35
	7

$$\therefore 113400 = 2^3 \times 3^4 \times 5^2 \times 7^1$$

$$\therefore p_1 = 2, p_2 = 3, p_3 = 5, p_4 = 7 \text{ and}$$

$$x_1 = 3, x_2 = 4, x_3 = 2, x_4 = 1$$

- 6. Find the LCM and HCF of 408 and 170 by applying the fundamental theorem of arithmetic.**

Solution :

Given no's are 408, 170

$\begin{array}{r l} 2 & 408 \\ 2 & 204 \\ 2 & 102 \\ 3 & 51 \\ \hline & 17 \end{array}$	$\begin{array}{r l} 2 & 170 \\ 5 & 85 \\ \hline & 17 \end{array}$
---	---

$$\therefore 408 = 2^3 \times 3 \times 17$$

$$170 = 2 \times 5 \times 17$$

$$\therefore \text{H.C.F} = 2 \times 17 = 34$$

$$\text{L.C.M} = 2^3 \times 17 \times 5 \times 3 = 2040$$

- 7. Find the greatest number consisting of 6 digits which is exactly divisible by 24, 15, 36?**

Solution :

First, we find the L.C.M of 24, 15, 36

$$\begin{array}{r|l} 3 & 24, 15, 36 \\ 2 & 8, 5, 12 \\ 2 & 4, 5, 6 \\ \hline & 2, 5, 3 \end{array}$$

$$\text{L.C.M} = 5 \times 3^2 \times 2^3$$

$$= 5 \times 9 \times 8$$

$$= 360$$

The greatest 6 digit no. is 999999

$$\begin{array}{r} 360 \overline{) 999999} \quad (277 \\ \underline{720} \\ 2799 \\ \underline{2520} \\ 2799 \\ \underline{2520} \\ 279 \\ \underline{279} \\ 0 \end{array}$$

\therefore Required greatest number

$$= 999999 - 279 \text{ (remainder)}$$

$$= 999720$$

- 8. What is the smallest number that when divided by three numbers such as 35, 56 and 91 leaves remainder 7 in each case?**

Solution :

91) + remainder 7

$$35 = 7 \times 5$$

$$56 = 7 \times 2 \times 2 \times 2$$

$$91 = 7 \times 13$$

$$\therefore \text{L.C.M} = 7 \times 5 \times 13 \times 8$$

$$= 3640$$

\therefore The required number is $3640 + 7$

$$= 3647$$

- 9. Find the least number that is divisible by the first ten natural numbers.**

Solution :

The required number is the LCM of

$$(1, 2, 3, \dots, 10)$$

$$2 = \underline{2} \times 1$$

$$4 = \underline{2} \times 2$$

$$6 = 3 \times \underline{2}$$

$$8 = 2 \times 2 \times \underline{2}$$

$$9 = 3 \times 3$$

$$10 = 5 \times \underline{2} \text{ and } 1, 3, 5, 7$$

$$\text{L.C.M} = 2 \times 2 \times 2 \times 3 \times 3 \times 5 \times 7$$

$$= 2520$$

III. MODULAR ARITHMETIC :

Key Points

- ✓ Two integers a and b are congruence modulo n if they differ by an integer multiple of n . That $b - a = kn$ for some integer k . This can also be written as $a \equiv b \pmod{n}$.
- ✓ Here the number n is called modulus. In other words, $a \equiv b \pmod{n}$ means $a - b$ is divisible by n .
- ✓ The equation $n = mq + r$ through Euclid's Division lemma can also be written as $n \equiv r \pmod{m}$.
- ✓ Two integers a and b are congruent modulo m , written as $a \equiv b \pmod{m}$, if they leave the same remainder when divided by m .
- ✓ a, b, c and d are integers and m is a positive integer such that if $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$ then
 - (i) $(a + c) \equiv (b + d) \pmod{m}$ (ii) $(a - c) \equiv (b - d) \pmod{m}$ (iii) $(a \times c) \equiv (b \times d) \pmod{m}$
 - (i) $ac \equiv bc \pmod{m}$ (ii) $a \pm c \equiv b \pm c \pmod{m}$ for any integer c

Example 2.11

Find the remainders when 70004 and 778 is divided by 7.

Solution :

Since 70000 is divisible by 7

$$70000 \equiv 0 \pmod{7}$$

$$70000 + 4 \equiv 0 + 4 \pmod{7}$$

$$70004 \equiv 4 \pmod{7}$$

Therefore, the remainder when 70004 is divided by 7 is 4.

Since 777 is divisible by 7

$$777 \equiv 0 \pmod{7}$$

$$777 + 1 \equiv 0 + 1 \pmod{7}$$

$$778 \equiv 1 \pmod{7}$$

Therefore, the remainder when 778 is divided by 7 is 1.

Example 2.12

Determine the value of d such that $15 \equiv 3 \pmod{d}$.

Solution :

$15 \equiv 3 \pmod{d}$ means $15 - 3 = kd$, for some integer k .

$$12 = kd. \text{ gives } d \text{ divides } 12.$$

The divisors of 12 are 1, 2, 3, 4, 6, 12. But d should be larger than 3 and so the possible values for d are 4, 6, 12.

Example 2.13

Find the least positive value of x such that

$$(i) 67 + x \equiv 1 \pmod{4} \quad (ii) 98 \equiv (x + 4) \pmod{5}$$

Solution :

$$(i) 67 + x \equiv 1 \pmod{4}$$

$$67 + x - 1 = 4n, \text{ for some integer } n$$

$$66 + x = 4n$$

$66 + x$ is a multiple of 4.

Therefore, the least positive value of x must be 2, since 68 is the nearest multiple of 4 more than 66.

$$(ii) \quad 98 \equiv (x + 4) \pmod{5}$$

$$98 - (x + 4) = 5n, \text{ for some integer } n.$$

$$94 - x = 5n$$

$94 - x$ is a multiple of 5.

Therefore, the least positive value of x must be 4

Since $94 - 4 = 90$ is the nearest multiple of 5 less than 94.

Example 2.14

Solve $8x \equiv 1 \pmod{11}$

Solution :

$11k$, for some integer k .

$$x = \frac{11k + 1}{8}$$

When we put $k = 5, 13, 21, 29, \dots$ then $11k + 1$ is divisible by 8.

$$x = \frac{11 \times 5 + 1}{8} = 7$$

$$x = \frac{11 \times 13 + 1}{8} = 18$$

Therefore, the solutions are 7, 18, 29, 40,

Example 2.15

Compute x , such that $10^4 \equiv x \pmod{19}$

Solution :

$$10^2 = 100 \equiv 5 \pmod{19}$$

$$10^4 = (10^2)^2 \equiv 5^2 \pmod{19}$$

$$10^4 \equiv 25 \pmod{19}$$

$$10^4 \equiv 6 \pmod{19} \quad (\text{since } 25 \equiv 6 \pmod{19})$$

Therefore, $x = 6$.

Example 2.16

Find the number of integer solutions of $3x \equiv 1 \pmod{15}$.

Solution : $3x \equiv 1 \pmod{15}$ can be written as

$$3x - 1 = 15k \text{ for some integer } k$$

$$3x = 15k + 1$$

$$x = \frac{15k + 1}{3}$$

$$x = 5k + \frac{1}{3}$$

Since $5k$ is an integer, $5k + \frac{1}{3}$ cannot be an integer.

So there is no integer solution.

Example 2.17

A man starts his journey from Chennai to Delhi by train. He starts at 22.30 hours on Wednesday. If it takes 32 hours of travelling time and assuming that the train is not late, when will he reach Delhi ?

Solution :

Starting time 22.30, Travelling time 32 hours. Here we use modulo 24.

The reaching time is

$$22.30 + 32 \pmod{24} \equiv 54.30 \pmod{24}$$

$$\equiv 6.30 \pmod{24}$$

$$(\text{Since } 32 = (1 \times 24) + 8)$$

Thursday Friday

Example 2.18

Kala and Vani are friends, Kala says, "Today is my birthday" and she asks Vani, "When will you celebrate your birthday?" Vani replies, "Today is Monday and I celebrated my birthday 75 days ago". Find the day when Vani celebrated her birthday.

Solution :

Let us associate the numbers 0, 1, 2, 3, 4, 5, 6 to represent the weekdays from Sunday to Saturday respectively.

Vani says today is Monday. So the number for Monday is 1. Since Vani's birthday was 75 days ago, we have to subtract 75 from 1 and take the modulo 7, since a week contains 7 days.

$$-74 \pmod{7} \equiv -4 \pmod{7} \equiv 7 - 4 \pmod{7} \equiv 3 \pmod{7}$$

(Since, $-74 - 3 = -77$ is divisible by 7)

$$\text{Thus, } 1 - 75 \equiv 3 \pmod{7}$$

The day for the number 3 is Wednesday.

Therefore, Vani's birthday must be on Wednesday.

1. Find the least positive value of x such that

(i) $71 \equiv x \pmod{8}$ (ii) $78 + x \equiv 3 \pmod{5}$

(iii) $89 \equiv (x + 3) \pmod{4}$

(iv) $96 \equiv \frac{x}{7} \pmod{5}$ (v) $5x \equiv 4 \pmod{6}$

Solution :

i) $71 \equiv x \pmod{8}$

$$71 - x = 8n$$

$\therefore 71 - x$ is a multiple of 8

\therefore The least +ve value of x is 7

($\because 71 - 7 = 64$ is the nearest multiple of 8) less than 71.

ii) $78 + x \equiv 3 \pmod{5}$

$$\Rightarrow 78 + x - 3 = 5n$$

$\Rightarrow 75 + x$ is a multiple of 5

\therefore The least +ve value of x is 5

($\because 75 + 5 = 80$, is the nearest multiple of 5 above 75)

iii) $89 \equiv (x + 3) \pmod{4}$

$$\Rightarrow 89 - x - 3 = 4n$$

$\Rightarrow 86 - x$ is a multiple of 4

\therefore The least +ve value is 2

($\because 86 - 2 = 84$ is the nearest multiple of 4 less than 86)

iv) $96 \equiv \frac{x}{7} \pmod{5}$

$$\Rightarrow 96 - \frac{x}{7} = 5n$$

$\Rightarrow 96 - \frac{x}{7}$ is a multiple of 5

\therefore The least +ve value of x is 7

($\because 96 - 1 = 95$ is a multiple of 5)

v) $5x \equiv 4 \pmod{6}$

$$\Rightarrow 5x - 4 = 6n$$

$\therefore 5x - 4$ is a multiple of 6

\therefore The least +ve value of x is 2

($\because 5(2) - 4 = 6$ is a multiple of 6)

2. If x is congruent to 13 modulo 17 then $7x - 3$ is congruent to which number modulo 17?

Solution :

Given $x \equiv 13 \pmod{17}$

$$\Rightarrow x - 13 \text{ is a multiple of } 17$$

\therefore The least +ve value of x is 30

$$\therefore 7x - 3 \equiv y \pmod{17}$$

$$\Rightarrow 7(30) - 3 \equiv y \pmod{17}$$

$$\Rightarrow 207 \equiv y \pmod{17}$$

$$\therefore y = 3 \quad (\because 207 - 3 = 204 \text{ is divisible by } 17)$$

3. Solve $5x \equiv 4 \pmod{6}$

Solution :

Given $5x \equiv 4 \pmod{6}$

$$\Rightarrow 5x - 4 \text{ is divisible by } 6$$

$$\therefore x = 2, 8, 14, \dots \quad (\text{by assumption})$$

4. Solve $3x - 2 \equiv 0 \pmod{11}$ **Solution :**

$$\text{Given } 3x - 2 \equiv 0 \pmod{11}$$

$$\therefore 3x - 2 \text{ is divisible by } 11$$

\therefore The possible values of x are 8, 19, 30,

5. What is the time 100 hours after 7 a.m.?**Solution :**

Formula :

$$t + n = f \pmod{24}$$

 $t \rightarrow$ current time $n \rightarrow$ no. of hrs. $f \rightarrow$ future time

$$100 + 7 = f \pmod{24}$$

$$\Rightarrow 107 - f \text{ is divisible by } 24$$

$\therefore f = 11$ so that $107 - 11 = 96$ is divisible by 24.

\therefore The time is 11 A.M.

6. What is the time 15 hours before 11 p.m.?**Solution :****Formula :**

$$\begin{array}{l|l} t - n \equiv p \pmod{24} & t \rightarrow \text{Current time} \\ \Rightarrow 11 - 15 = -4 \equiv -1 \times 24 + 20 & n \rightarrow \text{no of hrs} \\ \equiv 20 \pmod{24} & p \rightarrow \text{past time} \end{array}$$

\therefore The time 15 hours in the past was 8 p.m.

7. Today is Tuesday. My uncle will come after 45 days. In which day my uncle will be coming?**Solution :**

Today is Tuesday

Day after 45 days = ?

When we divide 45 by 7, remainder is 3.

\therefore The 3rd day from Tuesday is Friday

8. Prove that $2^n + 6 \times 9^n$ is always divisible by 7 for any positive integer n .**Solution :**

$$\text{When } n = 1, 2n + 6 \times 9^n$$

$$= 2 + (6 \times 9) = 56, \text{ divisible by } 7.$$

9. Find the remainder when 2^{81} is divided by 17.**Solution :**

To find the remainder when 2^{81} is divided by 17.

$$2^4 = 16 \equiv -1 \pmod{17}$$

$$\Rightarrow 2^{80} = (2^4)^{20} = (-1)^{20} = 1$$

$$\therefore 2^{81} = 2^{80} \times 2^1$$

$$= 1 \times 2$$

$$= 2$$

nai to London through British Airlines is approximately 11 hours. The airplane begins its journey on Sunday at 23:30 hours. If the time at Chennai is four and half hours ahead to that of London's time, then find the time at London, when will the flight lands at London Airport.

Solution :**Formula :**

$$\begin{array}{l|l} t + n \equiv f \pmod{24} & t \rightarrow \text{present time} \\ \Rightarrow 23.30 + 11 = f \pmod{24} & n \rightarrow \text{no of hours} \\ \Rightarrow 34.30 = f \pmod{24} & f \rightarrow \text{future time} \\ \therefore 34.30 - f \text{ is divisible by } 24 & \\ \Rightarrow f = 10.30 \text{ (a.m.)} & \end{array}$$

But the time difference between London & Chennai is 4.30 hrs.

\therefore Flight reaches London Airport at

$$= 10.30 \text{ hrs} - 4.30 \text{ hrs}$$

$$= 6 \text{ AM next day i.e. 6 AM on Monday}$$

IV. SEQUENCES :

Key Points

- ✓ A real valued sequence is a function defined on the set of natural numbers and taking real values.
- ✓ A sequence can be considered as a function defined on the set of natural numbers.
- ✓ Though all the sequences are functions, not all the functions are sequences.

Example 2.19

Find the next three terms of the sequences.

- (i) $\frac{1}{2}, \frac{1}{6}, \frac{1}{14}, \dots$ (ii) $5, 2, -1, -4, \dots$
 (iii) $1, 0.1, 0.01, \dots$

Solution :

$$(i) \quad \frac{1}{2}, \quad \frac{1}{6}, \quad \frac{1}{10}, \quad \frac{1}{14}, \quad \dots$$

$\xrightarrow{+4} \quad \xrightarrow{+4} \quad \xrightarrow{+4}$

In the above sequence the numerators are same and the denominator is increased by 4.

So the next three terms are

$$a_5 = \frac{1}{14+4} = \frac{1}{18}$$

$$a_6 = \frac{1}{18+4} = \frac{1}{22}$$

$$a_7 = \frac{1}{22+4} = \frac{1}{26}$$

$$(ii) \quad 5, \quad 2, \quad -1, \quad -4, \quad \dots$$

$\xrightarrow{-3} \quad \xrightarrow{-3} \quad \xrightarrow{-3}$

Here each term is decreased by 3. So the next three terms are $-7, -10, -13$.

$$(iii) \quad 1, \quad 0.1, \quad 0.01, \quad \dots$$

$\xrightarrow{+10} \quad \xrightarrow{+10}$

Here each term is divided by 10. Hence, the next three terms are

$$a_4 = \frac{0.01}{10} = 0.001$$

$$a_5 = \frac{0.001}{10} = 0.0001$$

$$a_6 = \frac{0.0001}{10} = 0.00001$$

Example 2.20

Find the general term for the following sequences

- (i) $3, 6, 9, \dots$ (ii) $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots$
 (iii) $5, -25, 125, \dots$

Solution :

- (i) $3, 6, 9, \dots$

Here the terms are multiples of 3. So the general term is

$$a_n = 3n,$$

- (ii) $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots$

$$a_1 = \frac{1}{2}; a_2 = \frac{2}{3}; a_3 = \frac{3}{4}$$

We see that the numerator of n^{th} terms is n , and the denominator is one more than the numerator. Hence, $a_n = \frac{n}{n+1}, n \in \mathbb{N}$

- (iii) $5, -25, 125, \dots$

The terms of the sequence have + and - sign alternatively and also they are in powers of 5.

So the general terms $a_n = (-1)^{n+1} 5^n, n \in \mathbb{N}$

Example 2.21

The general term of a sequence is defined as

$$a_n = \begin{cases} n(n+3); & n \in N \text{ is odd} \\ n^2 + 1 & ; n \in N \text{ is even} \end{cases}$$

Find the eleventh and eighteenth terms.

Solution :

To find a_{11} , since 11 is odd, we put

$$n = 11 \text{ in } a_n = n(n+3)$$

Thus, the eleventh term

$$a_{11} = 11(11+3) = 154,$$

To find a_{18} , since 18 is even we put

$$n = 18 \text{ in } a_n = n^2 + 1$$

Thus, the eighteenth term

$$a = 18^2 + 1 = 325.$$

Example 2.22

Find the first five terms of the following sequence.

$$a_1 = 1, a_2 = 1, a_n = \frac{a_{n-1}}{a_{n-2} + 3}; n \geq 3, n \in N$$

Solution :

The first two terms of this sequence are given by $a_1 = 1$, $a_2 = 1$. The third term a_3 depends on the first and second terms.

$$a_3 = \frac{a_{3-1}}{a_{3-2} + 3} = \frac{a_2}{a_1 + 3} = \frac{1}{1+3} = \frac{1}{4}$$

Similarly the fourth term a_4 depends upon a_2 and a_3 .

$$a_4 = \frac{a_{4-1}}{a_{4-2} + 3} = \frac{a_3}{a_2 + 3} = \frac{\frac{1}{4}}{1+3} = \frac{\frac{1}{4}}{4} = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$$

In the same way, the fifth term a_5 can be calculated as

$$a_5 = \frac{a_{5-1}}{a_{5-2} + 3} = \frac{a_4}{a_3 + 3} = \frac{\frac{1}{16}}{\frac{1}{4} + 3} = \frac{1}{16} \times \frac{4}{13} = \frac{1}{52}$$

Therefore, the first five terms of the sequence are $1, 1, \frac{1}{4}, \frac{1}{16}, \frac{1}{52}$

EXERCISE 2.4

1. Find the next three terms of the following sequence.

(i) 8, 24, 72, ...

(ii) 5, 1, -3, ...

(iii) $\frac{1}{4}, \frac{2}{9}, \frac{3}{16}, \dots$

Solution :

i) Given sequence is 8, 24, 72,

Each number is multiplied by 3

\therefore The next 3 terms in the sequence are

$$72 \times 3 = 216$$

$$216 \times 3 = 648$$

$$648 \times 3 = 1944$$

\therefore 216, 648, 1944

ii) Given sequence is 5, 1, -3,

Each number is subtracted by 4

$$-3 - 4 = -7$$

$$-7 - 4 = -11$$

$$-11 - 4 = -15$$

\therefore The next 3 terms in the sequence are -7, -11, -15

iii) Given sequence is $\frac{1}{4}, \frac{2}{9}, \frac{3}{16}, \dots$

Each no. in Numerator is increased by 1 & all nos in denominator are consecutive square no's

\therefore The next 3 terms are

$$\frac{4}{25}, \frac{5}{36}, \frac{6}{49}, \dots$$

2. Find the first four terms of the sequences whose n^{th} terms are given by

(i) $a_n = n^3 - 2$

(ii) $a_n = (-1)^{n+1}$

$n(n+1)$

(iii) $a_n = 2n^2 - 6$

Solution :

i) Given $a_n = n^3 - 2$

$$n = 1 \Rightarrow a_1 = 1^3 - 2 = 1 - 2 = -1$$

$$n = 2 \Rightarrow a_2 = 2^3 - 2 = 8 - 2 = 6$$

$$n = 3 \Rightarrow a_3 = 3^3 - 2 = 27 - 2 = 25$$

$$n = 4 \Rightarrow a_4 = 4^3 - 2 = 64 - 2 = 62$$

\therefore The first 4 terms are -1, 6, 25, 62

ii) Given $a_n = (-1)^{n+1} n(n+1)$

$$n = 1 \Rightarrow a_1 = (-1)^2 \cdot 1(2) = 2$$

$$n = 2 \Rightarrow a_2 = (-1)^3 \cdot 2(3) = -6$$

$$n = 3 \Rightarrow a_3 = (-1)^4 \cdot 3(4) = 12$$

$$n = 4 \Rightarrow a_4 = (-1)^5 \cdot 4(5) = -20$$

iii) Given $a_n = 2n^2 - 6$

$$n = 1 \Rightarrow a_1 = 2(1) - 6 = -4$$

$$n = 2 \Rightarrow a_2 = 2(4) - 6 = 2$$

$$n = 3 \Rightarrow a_3 = 2(9) - 6 = 12$$

$$n = 4 \Rightarrow a_4 = 2(16) - 6 = 26$$

\therefore The first 4 terms are -4, 2, 12, 26

3. Find the n^{th} term of the following sequences

(i) 2, 5, 10, 17, ... (ii) 0, $\frac{1}{2}$, $\frac{2}{3}$,

(iii) 3, 8, 13, 18, ...

Solution :

i) Given sequence is 2, 5, 10, 17,

$$\Rightarrow 1^2 + 1, 2^2 + 1, 3^2 + 1, 4^2 + 1, \dots$$

\therefore The n^{th} term of the sequence is

$$a_n = n^2 + 1$$

ii) Given sequence is 0, $\frac{1}{2}$, $\frac{1}{3}$,

$$\Rightarrow \frac{0}{1}, \frac{1}{2}, \frac{2}{3}, \dots$$

$$\Rightarrow \frac{1-1}{1}, \frac{2-1}{2}, \frac{3-1}{3}, \dots$$

\therefore The n^{th} term of the sequence is

$$a_n = \frac{n-1}{n}$$

iii) Given sequence is 3, 8, 13, 18,

$$\Rightarrow 5 - 2, 10 - 2, 15 - 2, 20 - 2, \dots$$

$$\Rightarrow 5(1) - 2, 5(2) - 2, 5(3) - 2, 5(4) - 2$$

\therefore The n^{th} term of the sequence is

$$a_n = 5n - 2$$

4. Find the indicated terms of the sequences whose n^{th} terms are given by

(i) $a_n = \frac{5n}{n+2}$; a_6 and a_{13}

(ii) $a_n = -(n^2 - 4)$; a_4 and a_{11}

Solution :

i) Given $a_n = \frac{5n}{n+2}$, a_6 , $a_{13} = ?$

$$a_6 = \frac{5(6)}{6+2} = \frac{30}{8} = \frac{15}{4}$$

$$a_{13} = \frac{5(13)}{13+2} = \frac{65}{15} = \frac{13}{3}$$

ii) $a_n = -(n^2 - 4)$; a_4 , $a_{11} = ?$

$$a_4 = -(16 - 4) = -12$$

$$a_{11} = -(121 - 4) = -117$$

5. Find a_8 and a_{15} whose n^{th} term is $a_n =$

$$\begin{cases} \frac{n^2 - 1}{n + 3} & ; n \text{ is even, } n \in N \\ \frac{n^2}{2n + 1} & ; n \text{ is odd,} \end{cases}$$

Solution :

Given

$$a_n = \begin{cases} \frac{n^2 - 1}{n + 3} ; n \text{ is even, } n \in \mathbb{N} \\ \frac{n^2}{2n + 1} ; n \text{ is odd,} \end{cases}$$

$$a_8 = \frac{8^2 - 1}{8 + 3} = \frac{63}{11}$$

$$a_{15} = \frac{15^2}{2(15) + 1} = \frac{225}{31}$$

6. If $a_1 = 1$, $a_2 = 1$ and $a_n = 2a_{n-1} + a_{n-2}$, $n \geq 3$, $n \in \mathbb{N}$, then find the first six terms of the sequence.

Solution :Given $a_1 = 1, a_2 = 1$

$$a_n = 2a_{n-1} + a_{n-2}$$

$$a_3 = 2a_2 + a_1$$

$$= 2(1) + 1$$

$$= 3$$

$$a_4 = 2a_3 + a_2$$

$$= 2(3) + 1$$

$$= 7$$

$$a_5 = 2a_4 + a_3$$

$$= 2(7) + 3$$

$$= 17$$

$$a_6 = 2a_5 + a_4$$

$$= 2(17) + 7$$

$$= 41$$

\therefore The first 6 terms are 1, 1, 3, 7, 17, 41

V. ARITHMETIC PROGRESSION :**Key Points**

- ✓ An Arithmetic Progression is a sequence whose successive terms differ by a constant number.
- ✓ Let a and d be real numbers. Then the numbers of the form $a, a + d, a + 2d, a + 3d, a + 4d, \dots$ is said to form Arithmetic Progression denoted by A.P. The number ' a ' is called the first term and ' d ' is called the common difference.
- ✓ If there are finite numbers of terms in an A.P. then it is called Finite Arithmetic Progression. If there are infinitely many terms in an A.P. then it is called Infinite Arithmetic Progression.
- ✓ The n^{th} term denoted by t_n can be written as $t_n = a + (n - 1)d$.
- ✓ The common difference of an A.P. can be positive, negative or zero.
- ✓ An Arithmetic progression having a common difference of zero is called a constant arithmetic progression.
- ✓ In a finite A.P. whose first term is a and last term l , then the number of terms in the A.P. is given by $l = a + (n - 1)d$ gives $n = \left(\frac{l - a}{d} \right) + 1$.
- ✓ If every term is added or subtracted by a constant, then the resulting sequence is also an A.P.

- ✓ In every term is multiplied or divided by a non-zero number, then the resulting sequence is also an A.P.
- ✓ If the sum of three consecutive terms of an A.P. is given, then they can be taken as $a - d$, a and $a + d$. Here the common difference is d .
- ✓ If the sum of four consecutive terms of an A.P. is given then, they can be taken as $a - 3d$, $a - d$, $a + d$ and $a + 3d$. Here common difference is $2d$.
- ✓ Three non-zero numbers a , b , c are in A.P. if and only if $\boxed{2b = a + c}$.

Example 2.23

Check whether the following sequences are in A.P. or not ?

(i) $x + 2, 2x + 3, 3x + 4, \dots$

(ii) $2, 4, 8, 16, \dots$

(iii) $3\sqrt{2}, 5\sqrt{2}, 7\sqrt{2}, 9\sqrt{2}, \dots$

Solution :

it is enough to check if the differences between the consecutive terms are equal or not.

(i) $t_2 - t_1 = (2x + 3) - (x + 2) = x + 1$

$$t_3 - t_2 = (3x + 4) - (2x + 3) = x + 1$$

$$t_2 - t_1 = t_3 - t_2$$

Thus, the differences between consecutive terms are equal.

Hence the sequence $x + 2, 2x + 3, 3x + 4, \dots$ is in A.P.

(ii) $t_2 - t_1 = 4 - 2 = 2$

$$t_3 - t_2 = 8 - 4 = 4$$

$$t_2 - t_1 \neq t_3 - t_2$$

Thus, the differences between consecutive terms are not equal. Hence the terms of the sequence $2, 4, 8, 16, \dots$ are not in A.P.

(iii) $t_2 - t_1 = 5\sqrt{2} - 3\sqrt{2} = 2\sqrt{2}$

$$t_3 - t_2 = 7\sqrt{2} - 5\sqrt{2} = 2\sqrt{2}$$

$$t_4 - t_3 = 9\sqrt{2} - 7\sqrt{2} = 2\sqrt{2}$$

Thus, the differences between consecutive terms are equal. Hence the terms of the sequence $3\sqrt{2}, 5\sqrt{2}, 7\sqrt{2}, 9\sqrt{2}, \dots$ are in A.P.

Example 2.24

Write an A.P. whose first term is 20 and common difference is 8.

Solution :

First term = $a = 20$; common difference = $d = 8$

Arithmetic Progression is $a, a + d, a + 2d, a + 3d, \dots$

In this case, we get $20, 20 + 8, 20 + 2(8), 20 + 3(8) \dots$

So, the required A.P. is $20, 28, 36, 44, \dots$

Example 2.25

Find the 15th, 24th and n^{th} term (general term) of an A.P. given by $3, 15, 27, 39, \dots$

Solution :

We have, first term = $a = 3$ and common difference = $d = 15 - 3 = 12$.

We know that n^{th} term (general term) of an A.P. with first term a and common difference d is given by $t_n = a + (n - 1)d$

$$t_{15} = a + (15 - 1)d = a + 14d$$

$$= 3 + 14(12) = 171$$

$$t_{24} = a + 23d = 3 + 23(12) = 279$$

(Here $a = 3$ and $d = 12$)

The n^{th} (general term) term is given by

$$t_n = a + (n - 1)d$$

$$\text{Thus, } t_n = 3 + (n - 1) 12$$

$$t_n = 12n - 9$$

Example 2.26

Find the number of terms in the A.P. 3, 6, 9, 12, 111.

Solution :

First term $a = 3$; common difference $d = 6 - 3 = 3$; last term $l = 111$

$$\text{We know that, } n = \left(\frac{l - a}{d} \right) + 1$$

$$n = \left(\frac{111 - 3}{3} \right) + 1 = 37$$

Thus the A.P. contain 37 terms.

Example 2.27

Determine the general term of an A.P. whose 7 term is -1 and 16th term is 17.

Solution :

Let the A.P. be $t_1, t_2, t_3, t_4, \dots$

It is given that $t_7 = -1$ and $t_{16} = 17$

$$a + (7 - 1)d = -1 \text{ and } a + (16 - 1)d = 17$$

$$a + 6d = -1 \quad \dots(1)$$

$$a + 15d = 17 \quad \dots(2)$$

Subtracting equation (1) from equation (2), we get $9d = 18$ gives $d = 2$

Putting $d = 2$ in equation (1), we get $a + 12 = -1$ so $a = -13$

$$\begin{aligned} \text{Hence, general term } t_n &= a + (n - 1)d \\ &= -13 + (n - 1) \times 2 \\ &= 2n - 15 \end{aligned}$$

Example 2.28

If $l^{\text{th}}, m^{\text{th}}$ and n^{th} terms of an A.P. are x, y, z respectively, then show that

$$(i) x(m - n) + y(n - l) + z(l - m) = 0$$

$$(ii) (x - y)n + (y - z)l + (z - x)m = 0$$

Solution :

(i) Let a be the first term and d be the common difference. It is given that

$$t_l = x, t_m = y, t_n = z$$

Using the general term formula

$$a + (l - 1)d = x \quad \dots(1)$$

$$a + (m - 1)d = y \quad \dots(2)$$

$$a + (n - 1)d = z \quad \dots(3)$$

We have, $x(m - n) + y(n - l) + z(l - m)$

$$= a[(m - n) + (n - l) + (l - m)] + d[(m - n)(l - 1) + (n - l)(m - 1) + (l - m)(n - 1)]$$

$$= a[0] + d[lm - ln - m + n + mn - lm - n + l + ln - mn - l + m]$$

$$= a(0) + d(0) = 0$$

(ii) On subtracting equation (2) from equation (1), equation (3) from equation (2) and equation (1) from equation (3), we get

$$x - y = (l - m)d$$

$$y - z = (m - n)d$$

$$z - x = (n - l)d$$

$$(x - y)n + (y - z)l + (z - x)m = [(l - m)n + (m - n)l + (n - l)m]d$$

$$= [ln - mn + lm - nl + nm - lm]d = 0$$

Example 2.29

In an A.P., sum of four consecutive terms is 28 and their sum of their squares is 276. Find the four numbers.

Solution :

Let us take the four terms in the form $(a - 3d)$, $(a - d)$, $(a + d)$ and $(a + 3d)$.

Since sum of the four terms is 28,

$$a - 3d + a - d + a + d + a + 3d = 28$$

$$4a = 28 \text{ gives } a = 7$$

Similarly, since sum of their squares is 276,

$$(a - 3d)^2 + (a - d)^2 + (a + d)^2 + (a + 3d)^2 = 276.$$

$$a^2 - 6ad + 9d^2 + a^2 - 2ad + d^2 + a^2 + 2ad + d^2 + a^2 + 6ad + 9d^2 = 276$$

$$4a^2 + 20d^2 = 276 \Rightarrow 4(7)^2 + 20d^2 = 276.$$

$$d^2 = 4 \text{ gives } d = \pm 2$$

If $d = 2$ then the four numbers are $7 - 3(2)$, $7 - 2$, $7 + 2$, $7 + 3(2)$

That is the four numbers are 1, 5, 9 and 13.

If $a = 7$, $d = -2$ then the four numbers are 13, 9, 5 and 1

Therefore, the four consecutive terms of the A.P. are 1, 5, 9 and 13.

Example 2.30

A mother divides ₹207 into three parts such that the amount are in A.P. and gives it to her three children. The product of the two least amounts that the children had ₹4623. Find the amount received by each child.

Solution :

Let the amount received by the three children be in the form of A.P. is given by

$a - d$, a , $a + d$. Since, sum of the amount is ₹207, we have

$$(a - d) + a + (a + d) = 307$$

$$3a = 207 \text{ gives } a = 69$$

It is given that product of the two least amounts is 4623

$$(a - d)a = 4623$$

$$(69 - d)69 = 4623$$

$$d = 2$$

Therefore, amount given by the mother to her three children are

₹(69 - 2), ₹69, ₹(69 + 2). That is, ₹67, ₹69 and ₹71.

EXERCISE 2.5

1. Check whether the following sequences are in A.P.

(i) $a - 3, a - 5, a - 7, \dots$

(ii) $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$

(iii) 9, 13, 17, 21, 25,

(iv) $\frac{-1}{3}, 0, \frac{1}{3}, \frac{2}{3}, \dots$

(v) 1, -1, 1, -1, 1, -1,

Solution :

i) $a - 3, a - 5, a - 7, \dots$

$$t_2 - t_1 = (a - 5) - (a - 3)$$

$$= a - 5 - a + 3$$

$$= -2$$

$$t_3 - t_2 = (a - 7) - (a - 5)$$

$$= a - 7 - a + 5$$

$$= -2$$

$$\therefore t_2 - t_1 = t_3 - t_2$$

\therefore The given sequence is in A.P.

ii) $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$

$$\begin{array}{l|l} t_2 - t_1 = \frac{1}{3} - \frac{1}{2} & t_3 - t_2 = \frac{1}{4} - \frac{1}{3} \\ = \frac{2-3}{6} & = \frac{3-4}{12} \\ = -\frac{1}{6} & = -\frac{1}{12} \end{array}$$

$$\therefore t_2 - t_1 \neq t_3 - t_2$$

\therefore The given sequence is not in A.P.

iii) 9, 13, 17, 21, 25,

Each term of the sequence is increased by a constant number 4.

\therefore The sequence is in A.P.

iv) $-\frac{1}{3}, 0, \frac{1}{3}, \frac{2}{3}, \dots$

$$\begin{array}{l} t_2 - t_1 = 0 - \left(-\frac{1}{3}\right) = \frac{1}{3} \\ t_3 - t_2 = \frac{1}{3} - 0 = \frac{1}{3} \\ t_4 - t_3 = \frac{2}{3} - \frac{1}{3} = \frac{1}{3} \end{array}$$

\therefore The sequence is in A.P.

v) 1, -1, 1, -1, 1, -1

$$t_2 - t_1 = -1 - 1 = -2$$

$$t_3 - t_2 = 1 - (-1) = 1 + 1 = 2$$

\therefore The sequence is not in A.P.

2. First term a and common difference d are given below. Find the corresponding A.P.

(i) $a = 5, d = 6$ (ii) $a = 7, d = -5$

(iii) $a = \frac{3}{4}, d = \frac{1}{2}$

Solution :

i) $a = 5, d = 6$

\therefore The A.P. is $a, a + d, a + 2d, a + 3d, \dots$

$$= 5, 5 + 6, 5 + 2(6), 5 + 3(6), \dots$$

$$= 5, 11, 17, 23, \dots$$

ii) $a = 7, d = -5$

\therefore The A.P. is

$$= 7, 7 + (-5), 7 + 2(-5), 7 + 3(-5), \dots$$

$$= 7, 2, -3, -8, \dots$$

iii) $a = \frac{3}{4}, d = \frac{1}{2}$

\therefore The A.P. is

$$= \frac{3}{4}, \frac{3}{4} + \frac{1}{2}, \frac{3}{4} + 2\left(\frac{1}{2}\right), \frac{3}{4} + 3\left(\frac{1}{2}\right), \dots$$

$$= \frac{3}{4}, \frac{5}{4}, \frac{7}{4}, \frac{9}{4}, \dots$$

3. Find the first term and common difference of the Arithmetic Progressions whose n^{th} terms are given below

"

"

Solution :

i) $t_n = -3 + 2n$

$$n = 1 \Rightarrow t_1 = -3 + 2(1) = -1$$

$$n = 2 \Rightarrow t_2 = -3 + 2(2) = 1$$

$$n = 3 \Rightarrow t_3 = -3 + 2(3) = 3 \dots$$

$$\therefore a = -1, \quad d = t_2 - t_1 = 1 - (-1)$$

$$d = 2$$

ii) $t_n = 4 - 7n$

$$n = 1 \Rightarrow t_1 = 4 - 7(1) = -3$$

$$n = 2 \Rightarrow t_2 = 4 - 7(2) = -10$$

$$n = 3 \Rightarrow t_3 = 4 - 7(3) = -17$$

$$\therefore a = -3, \quad d = t_2 - t_1 = -10 - (-3)$$

$$= -10 + 3$$

$$d = -7$$

4. Find the 19th term of an A.P. -11, -15, -19,.....

Solution :

Given A.P is -11, -15, -19,

$$\begin{aligned} a &= -11, \quad d = -15 - (-11) \\ &= -15 + 11 \\ &= -4 \end{aligned}$$

\therefore The 19th term is

$$\begin{aligned} t_{19} &= a + (19 - 1) d \\ &= a + 18 d \\ &= (-11) + 18 (-4) \\ &= -11 - 72 \\ &= -83 \end{aligned}$$

5. Which term of an A.P. 16, 11, 6, 1,... is -54?

Solution :

$$\begin{aligned} a &= 16, d = -5, t_n = -54 \\ \Rightarrow a + (n - 1) d &= -54 \\ \Rightarrow 16 + (n - 1) (-5) &= -54 \\ \Rightarrow 16 - 5n + 5 &= -54 \\ \Rightarrow -5n + 21 &= -54 \\ \Rightarrow -5n + 21 &= -54 \\ \Rightarrow -5n &= -54 - 21 \\ \Rightarrow -5n &= -75 \\ \therefore n &= 15 \end{aligned}$$

\therefore 15th term of A.P. is -54

6. Find the middle term(s) of an A.P. 9, 15, 21, 27, ..., 183.

Solution :

Given A.P is 9, 15, 21, 27, 183

$$a = 9, d = 6, l = 183$$

$$\begin{aligned} n &= \frac{l - a}{d} + 1 \\ &= \frac{183 - 9}{6} + 1 \\ &= \frac{174}{6} + 1 \\ &= 29 + 1 \end{aligned}$$

$$n = 30$$

$$\begin{aligned} \therefore \text{Middle terms are } \frac{30}{2}, \frac{30}{2} + 1 \\ = 15^{\text{th}}, 16^{\text{th}} \end{aligned}$$

$$\begin{array}{l|l} t_{15} = a + 14d & t_{16} = a + 15d \\ = 9 + 14(6) & = 9 + 15(6) \\ = 9 + 84 & = 9 + 90 \\ = 93 & = 99 \end{array}$$

\therefore The 2 middle terms are 93, 99.

7. If nine times ninth term is equal to the times twenty fourth term is zero.

Solution :

$$\text{Given } 9(t_9) = 15(t_{15})$$

$$\text{To Prove : } 6(t_{24}) = 0$$

$$\begin{aligned} \Rightarrow 9(t_9) &= 15(t_{15}) \\ \Rightarrow 9(a + 8d) &= 15(a + 14d) \\ \Rightarrow 3(a + 8d) &= 5(a + 14d) \\ \Rightarrow 3a + 24d &= 5a + 70d \\ \Rightarrow 2a + 46d &= 0 \\ \Rightarrow 2(a + 23d) &= 0 \end{aligned}$$

Multiplying 3 on both sides,

$$\begin{aligned} \Rightarrow 6(a + 23d) &= 0 \\ \Rightarrow 6(t_{24}) &= 0 \end{aligned}$$

Hence proved.

8. If $3 + k$, $18 - k$, $5k + 1$ are in A.P. then find k .

Solution :

Given $3 + k$, $18 - k$, $5k + 1$ are in A.P.

$$\Rightarrow (18 - k) - (3 + k) = (5k + 1) - (18 - k)$$

$$\Rightarrow 15 - 2k = 6k - 17$$

$$\Rightarrow -8k = -32$$

$$\Rightarrow k = 4$$

9. Find x , y and z , given that the numbers x , 10 , y , 24 , z are in A.P.

Solution :

Given that x , 10 , y , 24 , z are in A.P.

$\therefore y$ is the arithmetic mean of 10 & 24

$$\Rightarrow y = \frac{10 + 24}{2} = \frac{34}{2} = 17$$

$\therefore x$, 10 , y , 24 , z are in A.P.

Clearly $d = 7$

$$\therefore x = 10 - 7 = 3 \quad \& \quad z = 24 + 7 = 31$$

$$\therefore x = 3, y = 17, z = 31$$

10. In a theatre, there are 20 seats in the front row and 30 rows were allotted. Each successive row contains two additional seats than its front row. How many seats are there in the last row?

Solution :

By the data given,

$$a = 20, d = 2, n = 30$$

$$\begin{aligned} t_{30} &= a + 29d \\ &= 20 + 29(2) \\ &= 20 + 58 \\ &= 78 \end{aligned}$$

\therefore The no. of seats in 30th row = 78

11. The sum of three consecutive terms that are in A.P. is 27 and their product is 288. Find the three terms.

Solution :

Let the 3 consecutive terms in an A.P. be

$$a - d, a, a + d$$

i) Sum of 3 terms = 27

$$\Rightarrow a - d + a + a + d = 27$$

$$\Rightarrow 3a = 27$$

$$a = 9$$

ii) Product of 3 terms = 288

$$\Rightarrow (a - d) \cdot a \cdot (a + d) = 288$$

$$\Rightarrow a^2 (a^2 - d^2) = 288$$

$$\Rightarrow 9 (81 - d^2) = 288$$

$$\Rightarrow 81 - d^2 = 32$$

$$\Rightarrow d^2 = 49$$

$$\Rightarrow d = \pm 7$$

$a = 9, d = 7 \Rightarrow$ the 3 terms are 2, 9, 16

$a = 9, d = -7 \Rightarrow$ the 2 terms are 16, 9, 2

12. The ratio of 6th and 8th term of an A.P. is 7 : 9. Find the ratio of 9th term to 13th term.

Solution :

$$\text{Given } \frac{t_6}{t_8} = \frac{7}{9}$$

$$\Rightarrow \frac{a + 5d}{a + 7d} = \frac{7}{9}$$

$$\Rightarrow 9a + 45d = 7a + 49d$$

$$\Rightarrow 2a = 4d$$

$$\Rightarrow a = 2d \quad \dots(1)$$

$$\therefore \frac{t_9}{t_{13}} = \frac{a + 8d}{a + 12d}$$

$$= \frac{2d + 8d}{2d + 12d} \quad (\text{from (1)})$$

$$= \frac{10d}{14d}$$

$$= \frac{5}{7}$$

$$\therefore t_9 : t_{13} = 5 : 7$$

13. In a winter season let us take the temperature of Ooty from Monday to Friday to be in A.P. The sum of temperatures from Monday to Wednesday is 0°C and the sum of the temperatures from Wednesday to Friday is 18°C . Find the temperature on each of the five days.

Solution :

Let the temperature from Monday to Friday respectively be

$$a, a + d, a + 2d, a + 3d, a + 4d$$

i) Given $a + (a + d) + (a + 2d) = 0$

$$3a + 3d = 0$$

$$a + d = 0$$

$$a = -d$$

ii) Given $(a + 2d) + (a + 3d) + (a + 4d) = 18$

$$\Rightarrow 3a + 9d = 18$$

$$\Rightarrow 6d = 18$$

$$\Rightarrow d = 3$$

$$\therefore a = -3$$

The temperature of each of the 5 days

$$-3^\circ \text{C}, 0^\circ \text{C}, 3^\circ \text{C}, 6^\circ \text{C}, 9^\circ \text{C}$$

14. Priya earned ₹ 15,000 in the first year. Thereafter her salary increased by ₹ 1500 per year. Her expenses are ₹ 13,000 during the first year and the expenses increase by ₹ 900 per year. How long will it take for her to save ₹ 20,000.

Solution :

	1 st year	2 nd year
Salary :	₹15,000	₹16,500
Expense :	₹13,000	₹13,900
Savings :	₹2,000	₹2,600
\therefore the yearly savings are		
₹2,000, ₹2,600, ₹3,200, form an A.P		
with $a = 2,000$, $d = 600$, $t_n = 20,000$		
$a + (n - 1)d = 20,000$		
$\Rightarrow 2,000 + (n - 1)600 = 20,000$		
$\Rightarrow 600n - 600 = 18,000$		
$\Rightarrow 600n = 18,600$		
$\Rightarrow n = \frac{186}{6}$		
$n = 31$		

After 31 years, her savings will be ₹20,000.

VI. ARITHMETIC SERIES :

Key Points

- ✓ The sum of terms of a sequence is called series.
- ✓ Let $a_1, a_2, a_3, \dots, a_n, \dots$ be the sequence of real numbers. Then the real numbers $a_1 + a_2 + a_3 + \dots$ is defined as the series of real numbers.
- ✓ If a series has finite number of terms then it is called a Finite series. If a series has infinite number of terms then it is called Infinite series.
- ✓ Sum to n terms of an A.P. $S_n = \frac{n}{2}[2a + (n - 1)d]$
- ✓ If the first term a , and the last term l (n^{th} term) are given then $S_n = \frac{n}{2}[a + l]$.

Example 2.31

Find the sum of first 15 terms of the A.P.

$$8, 7\frac{1}{4}, 6\frac{1}{2}, 5\frac{3}{4}, \dots$$

Solution :

Here the first term $a = 8$, common difference $d = 7\frac{1}{4} - 8 = -\frac{3}{4}$,

Sum of first n terms of an A.P.

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$S_{15} = \frac{15}{2}\left[2 \times 8 + (15-1)\left(-\frac{3}{4}\right)\right]$$

$$S_{15} = \frac{15}{2}\left[16 - \frac{21}{2}\right] = \frac{165}{4}$$

Example 2.32

Find the sum of $0.40 + 0.43 + 0.46 + \dots + 1$.

Solution :

Here the value of n is not given. But the last term is given. From this, we can find the value of n .

Given $a = 0.40$ and $l = 1$, we find $d = 0.43 - 0.40 = 0.03$.

$$\text{Therefore, } n = \left(\frac{l-a}{d}\right) + 1$$

$$= \left(\frac{1-0.40}{0.03}\right) + 1 = 21$$

Sum of first n terms of an A.P.

$$S_n = \frac{n}{2}[a + l]$$

Here, $n = 21$.

$$\text{Therefore, } S_{21} = \frac{21}{2}[0.40 + 1] = 14.7$$

So, the sum of 21 terms of the given series is 14.7.

Example 2.33

How many terms of the series $1 + 5 + 9 + \dots$ must be taken so that their sum is 190 ?

Solution :

Here we have to find the value of n , such that $S_n = 190$.

First term $a = 1$, common difference

$$d = 5 - 1 = 4.$$

Sum of first n terms of an A.P.

$$S_n = \frac{n}{2}[2a + (n-1)d] = 190$$

$$\frac{n}{2}[2 \times 1 + (n-1) \times 4] = 190$$

$$n[4n - 2] = 380$$

$$2n^2 - n - 190 = 0$$

$$(n-10)(2n+19) = 0$$

But $n = 10$ as $n = -\frac{19}{2}$ is impossible. Therefore, $n = 10$.

Example 2.34

The 13th term of an A.P. is 3 and the sum of first 13 terms is 234. Find the common difference and the sum of first 21 terms.

Solution :

Given the 13th term = 3 so,

$$t_{13} = a + 12d = 3 \quad \dots(1)$$

Sum of first 13 terms = 234 gives

$$S_{13} = \frac{13}{2}[2a + 12d] = 234$$

$$2a + 12d = 36 \quad \dots(2)$$

Solving (1) and (2) we get, $a = 33$, $d = \frac{-5}{2}$

Therefore, common difference is $\frac{-5}{2}$.

Sum of first 21 terms

$$\begin{aligned} S_{21} &= \frac{21}{2} \left[2 \times 33 + (21-1) \times \left(-\frac{5}{2} \right) \right] \\ &= \frac{21}{2} [66 - 50] = 168 \end{aligned}$$

Example 2.35

In an AP, the sum of first n terms is $\frac{5n^2}{2} + \frac{3n}{2}$. Find the 17th term.

Solution :

The 17th term can be obtained by subtracting the sum of first 16 terms from the sum of first 17 terms.

$$\begin{aligned} S_{17} &= \frac{5 \times (17)^2}{2} + \frac{3 \times 17}{2} = \frac{1445}{2} + \frac{51}{2} = 748 \\ {}^{16} \quad \frac{5 \times (16)^2}{2} + \frac{3 \times 16}{2} &= \frac{1280}{2} + \frac{48}{2} \end{aligned}$$

$$\text{Now, } t_{17} = S_{17} - S_{16} = 748 - 664 = 84$$

Example 2.36

Find the sum of all natural numbers between 300 and 600 which are divisible by 7.

Solution :

The natural numbers between 300 and 600 which are divisible by 7 are 301, 308, 315, ..., 595.

The sum of all natural numbers between 300 and 600 is $301 + 308 + 315 + \dots + 595$.

The terms of the above series are in A.P.

First term $a = 301$; common difference $d = 7$; Last term $l = 595$.

$$n = \left(\frac{l - a}{d} \right) + 1 = \left(\frac{595 - 301}{7} \right) + 1 = 43$$

$$\text{Since, } S_n = \frac{n}{2} [a + l],$$


$$\text{we have } S_{43} = \frac{43}{2} [301 + 595] = 19264.$$

Example 2.37

A mosaic is designed in the shape of an equilateral triangle, 12 ft on each side. Each tile in the mosaic is in the shape of an equilateral triangle of 12 inch side. The tiles are alternate in colour as shown in the figure. Find the number of tiles of each colour and total number of tiles in the mosaic.

Solution :

Since the mosaic is in the shape of an equilateral triangle of 12 ft, and the tile is in the shape of an equilateral triangle of 12 inch (1 ft), there will be 12 rows in the mosaic.



From the figure, it is clear that number of white tiles in each row are 1, 2, 3, 4, ..., 12 which clearly forms an Arithmetic Progression.

Similarly the number of blue tiles in each row are 0, 1, 2, 3, ..., 11 which is also an Arithmetic Progression.

Number of white tiles

$$= 1 + 2 + 3 + \dots + 12 = \frac{12}{2} [1 + 12] = 78$$

Number of blue tiles

$$= 0 + 1 + 2 + 3 + \dots + 11 = \frac{12}{2} [0 + 11] = 66$$

$$\begin{aligned} \text{The total number of tiles in the mosaic} \\ = 78 + 66 = 144 \end{aligned}$$

Example 2.38

The houses of a street are numbered from 1 to 49. Senthil's house is numbered such that the sum of numbers of the houses prior to Senthil's house is equal to the sum of numbers of the houses following Senthil's house. Find Senthil's house number ?

Solution :

Let Senthil's house number be x .

It is given that $1 + 2 + 3 + \dots + (x - 1)$

$$= (x - 1) + (x + 2) + \dots + 49$$

$$1 + 2 + 3 + \dots + (x - 1)$$

$$= [1 + 2 + 3 + \dots + 49] - [1 + 2 + 3 + \dots + x]$$

$$\frac{x-1}{2}[1 + (x-1)] = \frac{49}{2}[1 + 49] - \frac{x}{2}[1 + x]$$

$$\frac{x(x-1)}{2} = \frac{49 \times 50}{2} - \frac{x(x+1)}{2}$$

$$x^2 - x = 2450 - x^2 - x \Rightarrow 2x^2 = 2450$$

$$x^2 = 1225 \text{ gives } x = 35$$

Therefore, Senthil's house number is 35.

Example 2.39

The sum of first, n , $2n$ and $3n$ terms of an A.P. are S_1 , S_2 and S_3 respectively. Prove that $S_3 = 3(S_2 - S_1)$.

Solution :

If S_1 , S_2 and S_3 are sum of first n , $2n$ and $3n$ terms of an A.P. respectively then

$$S_1 = \frac{n}{2}[2a + (n-1)d], \quad S_2 = \frac{2n}{2}[2a + (2n-1)d],$$

$$S_3 = \frac{3n}{2}[2a + (3n-1)d]$$

Consider

$$S_2 - S_1 = \frac{2n}{2}[2a + (2n-1)d] - \frac{n}{2}[2a + (n-1)d]$$

$$= \frac{n}{2}[(4a + 2(2n-1)d] - [2a + (n-1)d]$$

$$S_2 - S_1 = \frac{n}{2} \times [2a + (3n-1)d]$$

$$3(S_2 - S_1) = \frac{3n}{2}[2a + (3n-1)d]$$

$$3(S_2 - S_1) = S_3$$

EXERCISE 2.6**1. Find the sum of the following**

(i) 3, 7, 11,... up to 40 terms.

(ii) 102, 97, 92,... up to 27 terms.

(iii) 6 + 13 + 20 + + 97

Solution :

i) Given A.P is 3, 7, 11, up to 40 terms

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$S_{40} = \frac{40}{2}[6 + 39(4)]$$

$$= 20[6 + 156]$$

$$= 20 \times 162$$

$$= 3240$$

ii) Given A.P is 102, 97, 92, up to 27 terms

$$a = 102, d = -5, n = 27$$

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$S_{27} = \frac{27}{2}[204 + 26(-5)]$$

$$= \frac{27}{2}[204 - 130]$$

$$= \frac{27}{2} \times 74$$

$$= 27 \times 37$$

$$= 999$$

iii) Given $6 + 13 + 20 + \dots + 97$

$$a = 6, d = 7, l = 97$$

$$\therefore n = \frac{l - a}{d} + 1$$

$$= \frac{97 - 6}{7} + 1$$

$$= \frac{91}{7} + 1$$

$$= 13 + 1$$

$$= 14$$

$$\therefore S_n = \frac{n}{2}(a + l)$$

$$S_{14} = \frac{14}{2}(6 + 97)$$

$$= 7 \times 103$$

$$= 721$$

2. How many consecutive odd integers beginning with 5 will sum to 480?

Solution :

By the data given,

The series is $5 + 7 + 9 + \dots + n = 480$

$$\therefore a = 5, d = 2, S_n = 480$$

$$\Rightarrow \frac{n}{2} [2a + (n-1)d] = 480$$

$$\Rightarrow \frac{n}{2} [10 + (n-1)2] = 480$$

$$\Rightarrow \frac{n}{2} [5 + (n-1)] = 480$$

$$\Rightarrow n[n+4] = 480$$

$$\Rightarrow n^2 + 4n - 480 = 0$$

$$\Rightarrow (n+24)(n-20) = 0$$

$$\Rightarrow n = -24, n = 20$$

$$\therefore n = 20$$

3. Find the sum of first 28 terms of an A.P. whose n^{th} term is $4n - 3$.

Solution :

$$\text{Given } t_n = 4n - 3$$

$$n = 1 \Rightarrow t_1 = 4 - 3 = 1$$

$$n = 2 \Rightarrow t_2 = 8 - 3 = 5$$

$$n = 3 \Rightarrow t_3 = 12 - 3 = 9$$

$$\therefore a = 1, d = 5 - 1 = 4$$

$$\therefore S_n = \frac{n}{2} [2a + (n-1)d]$$

$$S_{28} = \frac{28}{2} [2 + 27(4)] = 14 [2 + 108]$$

$$= 14 \times 110 = 1540$$

4. The sum of first n terms of a certain series is given as $2n^2 - 3n$. Show that the series is an A.P.

Solution :

$$\text{Given } S_n = 2n^2 - 3n$$

$$n = 1 \Rightarrow S_1 = 2 - 3 = -1$$

$$n = 2 \Rightarrow S_2 = 2(4) - 3(2) = 8 - 6 = 2$$

$$\therefore t_1 = a = -1, S_2 = 2$$

$$\Rightarrow t_2 - t_1 = 2 - (-1) = 3$$

$$\Rightarrow t_2 = 3$$

$$\therefore t_1 = -1, t_2 = 3, d = t_2 - t_1 = 4$$

$$\therefore a = -1, d = 4$$

\therefore The series is $-1 + 3 + 7 + \dots$ is an A.P.

5. The 104th term and 4th term of an A.P. are 125 and 0. Find the sum of first 35 terms.

Solution :

$$\text{Given } t_{104} = 125, \quad t_4 = 0$$

To find : S_{35}

$$a + 103d = 125$$

$$a + 3d = 0$$

$$\hline 100d = 125$$

$$d = \frac{5}{4}$$

$$a + 3\left(\frac{5}{4}\right) = 0$$

$$\Rightarrow a + \frac{15}{4} = 0$$

$$\Rightarrow a = -\frac{15}{4}$$

$$\therefore S_n = \frac{n}{2} [2a + (n-1)d]$$

$$S_{35} = \frac{35}{2} \left[-\frac{15}{2} + (34)\left(\frac{5}{4}\right) \right]$$

$$= \frac{35}{2} \left[-\frac{15}{2} + \frac{85}{2} \right]$$

$$= \frac{35}{2} \times 35$$

$$= \frac{1225}{2}$$

$$= 612.5$$

6. Find the sum of all odd positive integers less than 450.

Solution :

To find the sum :

$$1 + 3 + 5 + 7 + \dots + 449$$

$$a = 1, d = 2, l = 449$$

$$\therefore n = \frac{l-a}{d} + 1$$

$$= \frac{449-1}{2} + 1$$

$$= \frac{448}{2} + 1$$

$$= 224 + 1$$

$$= 225$$

$$\therefore S_n = \frac{n}{2} [a + l]$$

$$S_{225} = \frac{225}{2} [450]$$

$$= 225 \times 225$$

$$= 50,625$$

7. Find the sum of all natural numbers between 602 and 902 which are not divisible by 4.

Solution :

First we take the sum of the numbers from 603 to 901

$$a = 603, d = 1, l = 901$$

$$\therefore n = \frac{l-a}{d} + 1$$

$$= \frac{901-603}{1} + 1$$

$$= 298 + 1$$

$$= 299$$

$$\therefore S_n = \frac{n}{2} [a + l]$$

$$= \frac{299}{2} \times 1504$$

$$= 299 \times 752$$

$$= 224848$$

Next we take sum of all the no's between 602 & 902 which are divi. by 4

$$a = 604, d = 4, l = 900$$

$$\therefore n = \frac{l-a}{d} + 1$$

$$= \frac{900-604}{4} + 1$$

$$= \frac{296}{4} + 1$$

$$= 74 + 1$$

$$= 75$$

$$\therefore S_n = \frac{n}{2} [a + l]$$

$$S_{75} = \frac{75}{2} \times 1504$$

$$= 75 \times 752$$

$$= 56,400$$

\therefore Sum of no's which are not div. by 4

$$= 224848 - 56400 = 168448$$

8. Raghu wish to buy a laptop. He can buy it by paying ₹ 40,000 cash or by giving it in 10 installments as ₹ 4800 in the first month, ₹ 4750 in the second month, ₹ 4700 in the third month and so on. If he pays the money in this fashion, find
- total amount paid in 10 installments.
 - how much extra amount that he has to pay than the cost?

Solution :

Installment in 1st month = Rs. 4800

Installment in 2nd month = Rs. 4750

Installment in 3rd month = Rs. 4700

i.e., 4800, 4750, 4700, forms an A.P. with

$$a = 4800, d = -50, n = 10$$

$$\therefore S_n = \frac{n}{2} [2a + (n-1)d]$$

$$\therefore S_{10} = 5 [9600 + 9(-50)]$$

$$= 5 [9600 - 450]$$

$$= 5 \times 9150$$

$$= \text{Rs. } 45,750/-$$

ii) Amount he paid extra in installments

$$= 45,750 - 40,000$$

$$= \text{Rs. } 5,750/-$$

9. A man repays a loan of ₹ 65,000 by paying ₹ 400 in the first month and then increasing the payment by ₹ 300 every month. How long will it take for him to clear the loan?

Solution :

Amounts of repayment in successive months

$$400 + 700 + 1000 + \dots n \text{ months} = ₹ 65,000$$

$$a = 400, d = 300, S_n = 65,000$$

$$\begin{aligned} \frac{n}{2} [2a + (n-1)d] &= 65,000 \\ \Rightarrow \frac{n}{2} [800 + (n-1)300] &= 65,000 \\ \Rightarrow n [400 + (n-1)150] &= 65,000 \\ \Rightarrow n [150n + 250] &= 65,000 \\ \Rightarrow n [3n + 5] &= 1,300 \\ \Rightarrow 3n^2 + 5n - 1300 &= 0 \\ \Rightarrow n &= 20, -\frac{65}{3} \end{aligned}$$

5	-3900
-65	-60
3	3

$$\frac{65}{3}, -20$$

$\therefore n = 20$

10. A brick staircase has a total of 30 steps. The bottom step requires 100 bricks. Each successive step requires two bricks less than the previous step.

(i) How many bricks are required for the

(ii) How many bricks are required to build the stair case?

Solution :

No. of bricks in bottom step = 100

No. of bricks in successive steps are

$$98, 96, 94, \dots$$

\therefore 100, 98, 96, 94, for 30 steps form an A.P. with

$$a = 100, d = -2, n = 30$$

i) No. of bricks used in the top most step

$$t_{30} = a + 29d$$

$$= 100 + 29(-2)$$

$$= 100 - 58$$

$$= 42$$

ii) Total no. of bricks used to build the stair case

$$\begin{aligned} S_{20} &= \frac{30}{2} (100 + 42) \\ &= 15 \times 142 \\ &= 2130 \end{aligned}$$

11. If $S_1, S_2, S_3, \dots, S_m$ are the sums of n terms of m A.P.'s whose first terms are 1, 2, 3, ..., m and whose common differences are 1, 3, 5, ..., $(2m - 1)$ respectively, then show that

$$S_1 + S_2 + S_3 + \dots + S_m = \frac{1}{2} mn(mn + 1)$$

Solution :

$$1^{st} \text{ A.P.} \Rightarrow a = 1, d = 1$$

$$\begin{aligned} \Rightarrow S_1 &= \frac{n}{2} [2 + (n - 1) 1] \\ &= \frac{n}{2} [n + 1] \end{aligned}$$

$$2^{nd} \text{ A.P.} \Rightarrow a = 2, d = 3$$

$$\begin{aligned} \Rightarrow S_2 &= \frac{n}{2} [4 + (n - 1) 3] \\ &= \frac{n}{2} [3n + 1] \end{aligned}$$

$$m^{th} \text{ A.P.} \Rightarrow a = m, d = 2m - 1$$

$$\begin{aligned} \Rightarrow S_m &= \frac{n}{2} [2m + (n - 1) (2m - 1)] \\ &= \frac{n}{2} [2m + 2mn - 2m - n + 1] \\ &= \frac{n}{2} [2mn - n + 1] \\ &= \frac{n}{2} [(2m - 1)n + 1] \end{aligned}$$

$$\begin{aligned} S_1 + S_2 + \dots + S_m &= \frac{n}{2} (n + 1) + \frac{n}{2} (3n + 1) + \dots + \frac{n}{2} ((2m - 1)n + 1) \\ &= \frac{n}{2} [(n + 3n + \dots + (2m - 1)n) + (1 + 1 + \dots m \text{ terms})] \\ &= \frac{n}{2} [(n(1 + 3 + 5 + \dots + (2m - 1)) + m)] \\ &= \frac{n}{2} [(n(m^2) + m)] \\ &= \frac{n}{2} [m(mn + 1)] \\ &= \frac{1}{2} mn(mn + 1) \\ &= \text{RHS} \end{aligned}$$

Hence Proved

12. Find the sum $\left[\frac{a-b}{a+b} + \frac{3a-2b}{a+b} + \frac{5a-3b}{a+b} + \dots \text{to 12 terms} \right]$

Solution :

Given series is

$$\frac{a-b}{a+b} \quad \frac{3a-2b}{a+b} \quad \frac{5a-3b}{a+b}$$

$$1^{st} \text{ term} = \frac{a-b}{a+b}, \quad \text{Common diff} = \frac{2a-b}{a+b}$$

$$\begin{aligned} S_{12} &= \frac{12}{2} \left[2 \left(\frac{a-b}{a+b} \right) + 11 \left(\frac{2a-b}{a+b} \right) \right] \left(\because S_n = \frac{n}{2} [2a + (n-1)d] \right) \\ &= 6 \left[\frac{2a-2b+22a-11b}{a+b} \right] \\ &= 6 \left[\frac{24a-13b}{a+b} \right] \\ &= \frac{6}{a+b} [24a-13b] \end{aligned}$$

Hence proved.

Key Points

- ✓ A Geometric Progression is a sequence in which each term is obtained by multiplying a fixed non-zero number to the preceding term except the first term. The fixed number is called common ratio. The common ratio is usually denoted by r .
- ✓ Let a and $r \neq 0$ be real numbers. Then the numbers of the form $a, ar, ar^2, \dots, ar^{n-1} \dots$ is called a Geometric Progression. The number ' a ' is called the first term and number ' r ' is called the common ratio.
- ✓ The general term or n^{th} term of a G.P. is $t_n = ar^{n-1}$.
- ✓ When the product of three consecutive terms of a G.P. are given, we can take the three terms as $\frac{a}{r}, a, ar$.
- ✓ When the products of four consecutive terms are given for a G.P. then we can take the four terms as $\frac{a}{r^3}, \frac{a}{r}, ar, ar^3$.
- ✓ When each term of a Geometric Progression is multiplied or divided by a non-zero constant then the resulting sequence is also a Geometric Progression.
- ✓ Three non-zero numbers a, b, c are in G.P. if and only if $b^2 = ac$.

Which of the following sequences form a Geometric Progression ?

- (i) 7, 14, 21, 28, (ii) $\frac{1}{2}, 1, 2, 4, \dots$
 (iii) 5, 25, 50, 75, ...

Solution :

To check if a given sequence form a G.P. we have to see if the ratio between successive terms are equal.

- (i) 7, 14, 21, 28,

$$\frac{t_2}{t_1} = \frac{14}{7} = 2; \quad \frac{t_3}{t_2} = \frac{21}{14} = \frac{3}{2}; \quad \frac{t_4}{t_3} = \frac{28}{21} = \frac{4}{3}$$

Since the ratios between successive terms are not equal, the sequence 7, 14, 21, 28, is not a Geometric Progression.

- (ii) $\frac{1}{2}, 1, 2, 4, \dots$

$$\frac{t_2}{t_1} = \frac{1}{\frac{1}{2}} = 2; \quad \frac{t_3}{t_2} = \frac{2}{1} = 2; \quad \frac{t_4}{t_3} = \frac{4}{2} = 2$$

Here the ratios between successive terms are equal. Therefore the sequence $\frac{1}{2}, 1, 2, 4, \dots$ is a Geometric Progression with common ratio $r = 2$.

- (iii) 5, 25, 50, 75, ...

$$\frac{t_2}{t_1} = \frac{25}{5} = 5; \quad \frac{t_3}{t_2} = \frac{50}{25} = 2; \quad \frac{t_4}{t_3} = \frac{75}{50} = \frac{3}{2}$$

Since the ratios between successive terms are not equal, the sequence 5, 25, 50, 75, ... is not a Geometric Progression.

Example 2.41

Find the geometric progression whose first term and common ratios are given by (i) $a = -7, r = 6$
 (ii) $a = 256, r = 0.5$

Solution :

(i) The general form of Geometric progression is a, ar, ar^2, \dots

$$a = -7, ar = -7 \times 6 = -42, ar^2 = -7 \times 6^2 = -252$$

Therefore the required Geometric Progression is $-7, -42, -252, \dots$

(ii) The general form of Geometric progression is a, ar, ar^2, \dots

$$a = 256, ar = 256 \times 0.5 = 128, ar^2 = 256 \times (0.5)^2 = 64$$

Example 2.42

Find the 8th term of the G.P. 9, 3, 1, ...

Solution :

To find the 8th term we have to use the n^{th} term formula $t_n = ar^{n-1}$

$$\frac{t_2}{t_1} = \frac{3}{9} = \frac{1}{3}$$

$$t_8 = 9 \times \left(\frac{1}{3}\right)^{8-1} = 9 \times \left(\frac{1}{3}\right)^7 = \frac{1}{243}$$

Therefore the 8th term of the G.P. is $\frac{1}{243}$.

Example 2.43

In a Geometric progression, the 4th term is $\frac{8}{9}$ and the 7th term is $\frac{64}{243}$. Find the Geometric Progression.

Solution :

$$4^{\text{th}} \text{ term, } t_4 = \frac{8}{9} \text{ gives } ar^3 = \frac{8}{9} \quad \dots(1)$$

$$7^{\text{th}} \text{ term, } t_7 = \frac{64}{243} \text{ gives } ar^6 = \frac{64}{243} \quad \dots(2)$$

$$\text{Dividing (2) by (1) we get, } \frac{ar^6}{ar^3} = \frac{\frac{64}{243}}{\frac{8}{9}}$$

$$r^3 = \frac{8}{27} \text{ gives } r = \frac{2}{3}$$

Substituting the value of r in (1), we get

$$a \times \left[\frac{2}{3}\right]^3 = \frac{8}{9} \Rightarrow a = 3$$

Therefore the Geometric Progression is a, ar, ar^2, \dots That is, $3, 2, \frac{4}{3}, \dots$

Example 2.44

The product of three consecutive terms of a Geometric Progression is 343 and their sum is $\frac{91}{3}$. Find the three terms.

Solution :

Since the product of 3 consecutive terms is given.

we can take them as $\frac{a}{r}, a, ar$.

$$\text{Product of the terms} = 343$$

$$\frac{a}{r}$$

$$a^3 = 7^3 \text{ gives } a = 7$$

$$\text{Sum of the terms} = \frac{91}{3}$$

$$\text{gives } 7 \left(\frac{1+r+r^2}{r} \right) = \frac{91}{3}$$

$$3 + 3r + 3r^2 = 13r \text{ gives } 3r^2 - 10r + 3 = 0$$

$$(3r - 1)(r - 3) = 0 \text{ gives } r = 3 \text{ or } r = \frac{1}{3}$$

If $a = 7, r = 3$ then the three terms are $\frac{7}{3}, 7, 21$.

If $a = 7, r = \frac{1}{3}$ then the three terms are $21, 7, \frac{7}{3}$.

Example 2.45

The present value of a machine is ₹40,000 and its value depreciates each year by 10%. Find the estimated value of the machine in the 6th year.

Solution :

The value of the machine at present is

₹40,000. Since it is depreciated at the rate of 10% after one year the value of the machine is 90% of the initial value.

That is the value of the machine at the end of the first year is $40,000 \times \frac{90}{100}$

After two years, the value of the machine is 90% of the value in the first year.

Value of the machine at the end of the 2nd year is $40,000 \times \left(\frac{90}{100}\right)^2$

Continuing this way, the value of the machine depreciates in the following way as

$$40000, 40000 \times \frac{90}{100}, 40000 \times \left(\frac{90}{100}\right)^2 \dots$$

This sequence is in the form of G.P. with first term 40,000 and common ratio $\frac{90}{100}$.

For finding the value of the machine at the end of 5th year (i.e. in 6th year), we need to find the sixth term of this G.P.

$$\text{Thus, } n = 6, a = 40,000, r = \frac{90}{100}.$$

$$\text{Using } t_n = ar^{n-1}, \text{ we have } t_6 = 40,000 \times \left(\frac{90}{100}\right)^{n-1}$$

$$= 40000 \times \left(\frac{90}{100}\right)^5$$

$$t_6 = 40,000 \times \frac{9}{10} \times \frac{9}{10} \times \frac{9}{10} \times \frac{9}{10} \times \frac{9}{10} \\ = 23619.6$$

Therefore the value of the machine in 6th year = ₹23619.60.

EXERCISE 2.7

1. Which of the following sequences are in G.P.?

(i) 3, 9, 27, 81, ... (ii) 4, 44, 444, 4444, ...

(iii) 0.5, 0.05, 0.005, ...

iv) $\frac{1}{3}, \frac{1}{6}, \frac{1}{12}, \dots$ v) 1, -5, 25, -125 ...

vi) 120, 60, 30, 18, ... vii) 16, 4, 1, $\frac{1}{4}, \dots$

Solution :

i) Given sequence is 3, 9, 27, 81, ...

$$\frac{t_2}{t_1} = \frac{9}{3} = 3$$

$$\frac{t_3}{t_2} = \frac{27}{9} = 3$$

$$\frac{t_4}{t_3} = \frac{81}{27} = 3$$

∴ The sequence is a G.P.

ii) Given sequence is 4, 44, 444, ...

$$\frac{t_2}{t_1} = \frac{44}{4} = 11$$

$$\frac{t_3}{t_2} = \frac{444}{44} = 11\frac{1}{11} \neq 11$$

$$\therefore \frac{t_2}{t_1} \neq \frac{t_3}{t_2}$$

∴ The sequence is not a G.P.

iii) Given sequence is 0.5, 0.05, 0.005, ...

$$\frac{t_2}{t_1} = \frac{0.05}{0.5} = \frac{5}{50} = \frac{1}{10}$$

$$\frac{t_3}{t_2} = \frac{0.005}{0.05} = \frac{5}{50} = \frac{1}{10}$$

∴ The sequence is a G.P.

iv) $\frac{1}{3}, \frac{1}{6}, \frac{1}{12}, \dots$

$$\frac{t_2}{t_1} = \frac{\frac{1}{6}}{\frac{1}{3}} = \frac{3}{6} = \frac{1}{2}$$

$$\frac{t_3}{t_2} = \frac{\frac{1}{12}}{\frac{1}{6}} = \frac{6}{12} = \frac{1}{2}$$

\therefore The sequence is a G.P.

v) $1, -5, 25, -125, \dots$

$$\frac{t_2}{t_1} = -5$$

$$\frac{t_3}{t_2} = \frac{25}{-5} = -5$$

$$\frac{t_4}{t_3} = \frac{-125}{25} = -5$$

\therefore The sequence is a G.P.

vi) $120, 60, 30, 18, \dots$

$$\frac{t_2}{t_1} = \frac{60}{120} = \frac{1}{2}$$

$$\frac{t_3}{t_2} = \frac{30}{60} = \frac{1}{2}$$

$$\frac{t_4}{t_3} = \frac{18}{30} \neq \frac{1}{2}$$

\therefore The sequence is not a G.P.

vii) $16, 4, 1, \frac{1}{4}, \dots$

$$\frac{t_2}{t_1} = \frac{4}{16} = \frac{1}{4}$$

$$\frac{t_3}{t_2} = \frac{1}{4}$$

$$\frac{t_4}{t_3} = \frac{1}{4}$$

\therefore The sequence is a G.P.

2. Write the first three terms of the G.P. whose first term and the common ratio are given below.

(i) $a = 6, r = 3$ (ii) $a = \sqrt{2}, r = \sqrt{2}$

(iii) $a = 1000, r = \frac{2}{5}$

Solution :

i) Given $a = 6, r = 3$

\therefore The first 3 terms of the G.P. are

$$6, 18, 54, \dots$$

ii) Given $a = \sqrt{2}, r = \sqrt{2}$

\therefore The first 3 terms of the G.P. are

$$\sqrt{2}, 2, 2\sqrt{2}, \dots$$

iii) $a = 1000, r = \frac{2}{5}$

$$1000, 1000 \times \frac{2}{5}, 1000 \times \frac{2}{5} \times \frac{2}{5} \\ = 1000, 400, 160, \dots$$

3. In a G.P. 729, 243, 81, ... find t_7 .

Solution :

Given G.P is 729, 243, 81, ...

$$a = 729, r = \frac{81}{243} = \frac{1}{3}$$

$$\therefore t_n = a \cdot r^{n-1}$$

$$\Rightarrow t_7 = a \cdot r^6$$

$$= 729 \times \left(\frac{1}{3}\right)^6$$

$$= 3^6 \times \frac{1}{3^6} = 1$$

4. Find x so that $x + 6, x + 12$ and $x + 15$ are consecutive terms of a Geometric Progression.

Solution :

Given $x + 6, x + 12, x + 15$ are consecutive terms of a G.P.

$$\begin{aligned} \Rightarrow \quad \frac{x+12}{x+6} &= \frac{x+15}{x+12} \\ \Rightarrow \quad (x+12)^2 &= (x+15)(x+6) \\ \Rightarrow \quad x^2 + 24x + 144 &= x^2 + 21x + 90 \\ \Rightarrow \quad 3x &= -54 \\ \Rightarrow \quad x &= -18 \end{aligned}$$

5. Find the number of terms in the following G.P.

i) 4, 8, 16, ..., 8192 ?

ii) $\frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \dots, \frac{1}{2187}$?

Solution :

i) Given G.P is 4, 8, 16, 8192

$$\begin{aligned} \Rightarrow \quad a &= 4, r = 2, t_n = 8192 \\ \Rightarrow \quad a \cdot r^{n-1} &= 8192 \\ \Rightarrow \quad 4 \times 2^{n-1} &= 8192 \\ \Rightarrow \quad 2 &= 2048 \\ \Rightarrow \quad 2^{n-1} &= 2^{11} \\ \Rightarrow \quad n-1 &= 11 \\ \therefore n &= 12 \end{aligned}$$

ii) Given G.P is $\frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \dots, \frac{1}{2187}$

$$\begin{aligned} a &= \frac{1}{3}, r = \frac{1}{3}, t_n = \frac{1}{2187} \\ \Rightarrow \quad a \cdot r^{n-1} &= \frac{1}{2187} \\ \Rightarrow \quad \frac{1}{3} \cdot \left(\frac{1}{3}\right)^{n-1} &= \frac{1}{2187} \\ \Rightarrow \quad \left(\frac{1}{3}\right)^{n-1} &= \frac{3}{2187} \\ \Rightarrow \quad \left(\frac{1}{3}\right)^{n-1} &= \frac{1}{729} \\ \Rightarrow \quad \left(\frac{1}{3}\right)^{n-1} &= \left(\frac{1}{3}\right)^6 \\ \therefore n-1 &= 6 \end{aligned}$$

6. In a G.P. the 9th term is 32805 and 6th term is 1215. Find the 12th term.

Solution :

Given $t_9 = 32805, t_6 = 1215, t_{12} = ?$

$$a \cdot r^8 = 32805 \quad \dots\dots\dots (1)$$

$$a \cdot r^5 = 1215 \quad \dots\dots\dots (2)$$

$$(1) \div (2) \Rightarrow r^3 = \frac{32805}{1215}$$

$$\Rightarrow r^3 = 27$$

$$\Rightarrow r = 3$$

Sub. $r = 3$ in (2)

$$a \times 3^5 = 1215$$

$$\Rightarrow a \times 243 = 1215$$

$$\Rightarrow a = \frac{1215}{243}$$

$$\Rightarrow a = 5$$

$$\begin{aligned} \Rightarrow \therefore t_{12} &= a \cdot r^{11} \\ &= 5 \times 3^{11} \end{aligned}$$

7. Find the 10th term of a G.P. whose 8th term is 768 and the common ratio is 2.

Solution :

Given $t_8 = 768, r = 2$

$$\Rightarrow a \cdot r^7 = 768$$

$$\Rightarrow a \times 2^7 = 768$$

$$\Rightarrow a \times 128 = 768$$

$$a = 6$$

$$\therefore t_{10} = a \cdot r^9$$

$$= 6 \times 2^9$$

$$= 6 \times 512$$

$$= 3072$$

8. If a, b, c are in A.P. then show that $3a, 3b, 3c$ are in G.P.

Solution :

Given a, b, c are in A.P.

$$\Rightarrow b = \frac{a+c}{2} \quad \dots\dots(1)$$

To Prove : $3^a, 3^b, 3^c$ are in G.P.

i.e. TP : $(3^b)^2 = 3^a \cdot 3^c$

$$\begin{aligned} \text{LHS : } & (3^b)^2 \\ &= 3^{2b} \\ &= 3^{a+c} \quad (\text{from (1)}) \\ &= 3^a \cdot 3^c \\ &= \text{RHS} \\ \therefore 3^a, 3^b, 3^c &\text{ are in G.P.} \end{aligned}$$

9. In a G.P. the product of three consecutive terms is 27 and the sum of the product of two terms taken at a time is $\frac{57}{2}$. Find the three terms.

Solution :

Let the 3 consecutive terms of a G.P be

$$\frac{a}{r}, a, ar$$

i) Product of 3 terms = 27

$$\Rightarrow \frac{a}{r} \times a \times ar = 27$$

$$\begin{aligned} \Rightarrow a^3 &= 27 \\ \therefore a &= 3 \end{aligned}$$

ii) Sum of product of terms taken 2 at a

$$\text{time} = \frac{57}{2}$$

$$\text{i.e., } \frac{a}{r} \times a + a \times ar + ar \times \frac{a}{r} = \frac{57}{2}$$

$$\Rightarrow a^2 \left[\frac{1}{r} + r + 1 \right] = \frac{57}{2}$$

$$\Rightarrow 9 \left[\frac{1+r^2+r}{r} \right] = \frac{57}{2}$$

$$\Rightarrow \frac{1+r+r^2}{r} = \frac{19}{2}$$

$$\Rightarrow \frac{1+r+r^2}{r} = \frac{19}{6}$$

$$\Rightarrow 6r^2 + 6r + 6 = 19r$$

$$\Rightarrow 6r^2 - 13r + 6 = 0$$

$$\Rightarrow r = \frac{3}{2}, \frac{2}{3}$$

$$\therefore a = 3, r = \frac{3}{2} \Rightarrow 3 \text{ terms are } 2, 3, \frac{9}{2}$$

&

$$\therefore a = 3, r = \frac{3}{2} \Rightarrow 3 \text{ terms are } \frac{9}{2}, 3, 2$$

Manager. The company gave him a starting salary of ₹ 60,000 and agreed to increase his salary 5% annually. What will be his salary after 5 years?

Solution :

Given, initial salary = Rs. 60,000

Annual increment = 5%

Salary increment at the end of 1 year =

$$60,000 \times \frac{5}{100} = 3000$$

\therefore Continuing this way, we want to find the total salary after 5 years.

$$\begin{aligned} A &= P \left(1 + \frac{r}{100} \right)^n \\ &= 60,000 \left(1 + \frac{5}{100} \right)^5 \\ &= 60,000 \times \left(\frac{105}{100} \right)^5 \\ &= 60,000 \times (1.05)^5 \\ &= \text{Rs. } 76,600 \end{aligned}$$

$$\log 60,000 = 4.7782$$

$$5 \log (1.05) = 0.1060$$

$$5.8842$$

$$\text{Antilog } 76,600$$

11. Sivamani is attending an interview for a job and the company gave two offers to him.

Offer A: ₹ 20,000 to start with followed by a guaranteed annual increase of 6% for the first 5 years.

Offer B: ₹ 22,000 to start with followed by a guaranteed annual increase of 3% for the first 5 years.

What is his salary in the 4th year with respect to the offers A and B?

Offer A :

$$P = ₹ 20,000 \quad r = 6\%$$

$$n = 3 \text{ (in the 4th year)}$$

$$A = P \left(1 + \frac{r}{100}\right)^3$$

$$\log 20,000 = 4.3010$$

$$3 \log (1.06) = 0.0759$$

$$4.3769$$

$$\text{Antilog } 23820$$

$$= 20,000 \left(1 + \frac{6}{100}\right)^3$$

$$= 20,000 \left(\frac{106}{100}\right)^3$$

$$= 20,000 (1.06)^3$$

$$= 23,820$$

Offer B :

$$P = ₹ 22,000 \quad r = 3\%$$

$$n = 3 \text{ (in the 4th year)}$$

$$A = P \left(1 + \frac{r}{100}\right)^3$$

$$\log 22,000 = 4.3424$$

$$3 \log (1.03) = 0.0384$$

$$= 22,000 \times (1.03)^3 \quad \frac{4.3808}{4.3808}$$

$$= ₹ 24040$$

$$\text{Antilog } 24040$$

12. If a, b, c are three consecutive terms of an A.P. and x, y, z are three consecutive terms of a G.P. then prove that $x^{b-c} \times y^{c-a} \times z^{a-b} = 1$.

Solution :

Given a, b, c are consecutive terms of A.P.

$$\Rightarrow a, a + d, a + 2d, \dots$$

x, y, z are consecutive terms of G.P.

$$\Rightarrow x, xr, xr^2, \dots$$

$$\text{T.P : } x^{b-c} \times y^{c-a} \times z^{a-b} = 1$$

$$\begin{aligned} \text{LHS : } x^{b-c} \times y^{c-a} \times z^{a-b} &= x^{-d} \times (xr)^{2d} \times (xr^2)^{-d} \\ &= x^0 \times r^{2d} \times r^{-2d} = x^0 \times r^0 = 1 \\ &= \text{RHS. Hence proved.} \end{aligned}$$

VIII. GEOMETRIC SERIES :

Key Points

- ✓ A series whose terms are in Geometric progression is called Geometric series.
- ✓ The sum to n terms is $S_n = \frac{a(r^n - 1)}{r - 1}$, $r \neq 1$.
- ✓ If $r = 1$, then $S_n = a + a + a + \dots + a = na$.
- ✓ The sum of infinite terms of a G.P. is given by $a + ar + ar^2 + ar^3 + \dots = \frac{a}{1 - r}$, $-1 < r < 1$.

Example 2.46

Find the sum of 8 terms of the G.P.

1, -3, 9, -27, ...

Solution :

Here the first term $a = 1$,
 $\frac{-3}{1}$

Sum to n terms of a G.P. is

$$S_n = \frac{a(r^n - 1)}{r - 1} \text{ if } r \neq 1$$

$$\text{Hence, } S_8 = \frac{1((-3)^8 - 1)}{(-3) - 1} = \frac{6561 - 1}{-4} = -1640$$

Example 2.47

Find the first term of G.P. in which $S_6 = 4095$ and $r = 4$.

Solution :

Common ratio $= 4 > 1$, Sum of first 6 terms
 $S_6 = 4095$

$$\text{Hence, } S_n = \frac{a(r^n - 1)}{r - 1} = 4095$$

$$\text{Since, } r = 4, \frac{a(4^6 - 1)}{4 - 1} = 4095 \text{ gives}$$

$$a \times \frac{4095}{3} = 4095$$

First term $a = 3$.

Example 2.48

How many terms of the series $1 + 4 + 16 + \dots$ make the sum 1365 ?

Solution :

Let n be the number of terms to be added to get the sum 1365

$$a = 1, r = \frac{4}{1} = 4 > 1$$

$$S_n = 1365 \text{ gives } \frac{a(r^n - 1)}{r - 1} = 1365$$

$$\frac{1(4^n - 1)}{4 - 1} = 1365 \text{ so, } (4^n - 1) = 4095$$

$$4^n = 4096 \text{ then } 4^n = 4^6$$

$$n = 6$$

Example 2.49

Find the sum $3 + 1 + \frac{1}{3} + \dots \infty$

Solution :

$$\text{Here } a = 3, r = \frac{t_2}{t_1} = \frac{1}{3}$$

$$\text{Sum of infinite terms} = \frac{a}{1 - r} = \frac{3}{1 - \frac{1}{3}} = \frac{9}{2}$$

Example 2.50

Find the rational form of the number 0.6666 ...

Solution :

We can express the number 0.6666 ... as follows

$$0.6666... = 0.6 + 0.06 + 0.006 + 0.0006 + ...$$

We now see that numbers 0.6, 0.06, 0.006 ... forms an G.P. whose first term $a = 0.6$ and common ratio $r = \frac{0.06}{0.6} = 0.1$. Also $-1 < r = 0.1 < 1$

Using the infinite G.P. formula, we have

$$\begin{aligned} 0.6666... &= 0.6 + 0.06 + 0.006 + 0.0006 + ... \\ &= \frac{0.6}{1-0.1} = \frac{0.6}{0.9} = \frac{2}{3} \end{aligned}$$

Thus the rational number equivalent of $\frac{2}{3}$

Example 2.51

Find the sum to n terms of the series $5 + 55 + 555 + ...$

Solution :

The series is neither Arithmetic nor Geometric series. So it can be split into two series and then find the sum.

$$\begin{aligned} 5 + 55 + 555 + ... + n \text{ terms} &= 5 [1 + 11 + 111 + ... + n \text{ terms}] \\ &= \frac{5}{9} [9 + 99 + 999 + ... + n \text{ terms}] \\ &= \frac{5}{9} [(10-1) + (100-1) + (1000-1) + ... + n \text{ terms}] \\ &= \frac{5}{9} [(10 + 100 + 1000 + ... + n \text{ terms}) - n] \\ &= \frac{5}{9} \left[\frac{10(10^n - 1)}{(10 - 1)} - n \right] = \frac{50(10^n - 1)}{81} - \frac{5n}{9} \end{aligned}$$

Example 2.52

Find the least positive integer n such that $1 + 6 + 6^2 + ... + 6^n > 5000$.

Solution :

We have to find the least number of terms for which the sum must be greater than 5000.

That is, to find the least value of n , such that $S_n > 5000$

$$\text{We have } S_n = \frac{a(r^n - 1)}{r - 1} = \frac{1(6^n - 1)}{6 - 1} = \frac{6^n - 1}{5}$$

$$S_n > 5000 \text{ gives } \frac{6^n - 1}{5} > 5000$$

$$6^n - 1 > 25000 \text{ gives } 6^n > 25001$$

$$\text{Since, } 6^5 = 7776 \text{ and } 6^6 = 46656$$

The least positive value of n is 6 such that $1 + 6 + 6^2 + ... + 6^n > 5000$.

Example 2.53

A person saved money every year, half as much as he could in the previous year. If he had totally saved ₹ 7875 in 6 years then how much did he save in the first year ?

Solution :

Total amount saved in 6 years is $S_6 = 7875$

Since he saved half as much money as every year he saved in the previous year.

$$\text{We have } r = \frac{1}{2} < 1$$

$$\frac{a(1 - r^n)}{1 - r} = \frac{a \left(1 - \left(\frac{1}{2} \right)^6 \right)}{1 - \frac{1}{2}} = 7875$$

$$\frac{a \left(1 - \frac{1}{64} \right)}{\frac{1}{2}} = 7875 \text{ gives } a \times \frac{63}{32} = 7875$$

$$a = \frac{7875 \times 32}{63} \text{ so, } a = 4000$$

The amount saved in the first year is ₹4000.

EXERCISE 2.8**1. Find the sum of first n terms of the G.P.**

(i) $5, -3, \frac{9}{5}, -\frac{27}{25}, \dots$ (ii) 256, 64, 16, \dots

Solution :

i) Given G.P is $5, -3, \frac{9}{5}, -\frac{27}{25}, \dots$

$$a = 5, r = -\frac{3}{5} < 1$$

$$\therefore S_n = a \cdot \frac{1-r^n}{1-r}$$

$$= (5) \times \left(\frac{1 - \left(-\frac{3}{5}\right)^n}{1 - \left(-\frac{3}{5}\right)} \right)$$

$$= (5) \times \left(\frac{1 - \left(-\frac{3}{5}\right)^n}{\frac{8}{5}} \right)$$

$$\frac{25}{8} \left(\left(-\frac{3}{5}\right)^n \right)$$

ii) Given G.P is 256, 64, 16, \dots

$$a = 256, r = \frac{1}{4} < 1$$

$$\therefore S_n = a \cdot \frac{1-r^n}{1-r}$$

$$= 256 \times \frac{1 - \left(\frac{1}{4}\right)^n}{1 - \frac{1}{4}}$$

$$= 256 \times \frac{1 - \left(\frac{1}{4}\right)^n}{\frac{3}{4}}$$

$$= \frac{1024}{3} \left(1 - \left(\frac{1}{4}\right)^n \right)$$

2. Find the sum of first six terms of the G.P.
5, 15, 45, ...**Solution :**Given G.P is 5, 15, 45, \dots

$$a = 5, r = 3 > 1$$

$$S_n = a \cdot \frac{r^n - 1}{r - 1}$$

$$\therefore S_6 = 5 \cdot \frac{3^6 - 1}{3 - 1}$$

$$= \frac{5}{2} \times 728$$

$$= 5 \times 364$$

$$= 1820$$

3. Find the first term of the G.P. whose common ratio 5 and whose sum to first 6 terms is 46872.**Solution :**

$$\text{Given } r = 5, S_6 = 46872$$

$$S_n = a \cdot \frac{r^n - 1}{r - 1}$$

$$\Rightarrow a \times \frac{5^6 - 1}{4} = 46872$$

$$\Rightarrow a(5^6 - 1) = 46872 \times 4$$

$$\Rightarrow a(15624) = 46872 \times 4$$

$$\therefore a = \frac{46872 \times 4}{15624}$$

$$= 3 \times 4$$

$$a = 12$$

4. Find the sum to infinity of

(i) $9 + 3 + 1 + \dots$

(ii) $21 + 14 + \frac{28}{3} + \dots$

Solution :

i) $9 + 3 + 1 + \dots$ is a geometric series

$$\text{with } a = 9, r = \frac{1}{3} < 1$$

$$S_\infty = \frac{a}{1-r} = \frac{9}{1 - \frac{1}{3}}$$

$$= \frac{9}{\frac{2}{3}}$$

$$= \frac{27}{2}$$

ii) $21 + 14 + \frac{28}{3}, \dots$ is geo. series

$$\text{with } a = 21, r = \frac{14}{21} = \frac{2}{3} < 1$$

$$\begin{aligned} \therefore S_{\infty} &= \frac{a}{1-r} = \frac{21}{1-\frac{2}{3}} \\ &= \frac{21}{\frac{1}{3}} \\ &= \frac{21}{1/3} \\ &= 63 \end{aligned}$$

5. If the first term of an infinite G.P. is 8 and its sum to infinity is $\frac{32}{3}$ then find the common ratio.

Solution :

$$\begin{aligned} \frac{\infty}{3} \\ \Rightarrow \frac{a}{1-r} &= \frac{32}{3} \\ \Rightarrow \frac{8}{1-r} &= \frac{32}{3} \\ \Rightarrow 3 &= 4 - 4r \\ \Rightarrow 4r &= 1 \\ \therefore r &= \frac{1}{4} \end{aligned}$$

6. Find the sum to n terms of the series

(i) $0.4 + 0.44 + 0.444 + \dots$ to n terms

(ii) $3 + 33 + 333 + \dots$ to n terms

Solution :

i) $0.4 + 0.44 + 0.444 + \dots$ to n terms

$$\begin{aligned} &= \frac{4}{10} + \frac{44}{100} + \frac{444}{1000} + \dots \text{ to } n \text{ terms} \\ &= 4 \left[\frac{1}{10} + \frac{11}{100} + \frac{111}{1000} + \dots n \text{ terms} \right] \end{aligned}$$

$$\begin{aligned} &= \frac{4}{9} \left[\frac{9}{10} + \frac{99}{100} + \frac{999}{1000} + \dots n \text{ terms} \right] \\ &= \frac{4}{9} \left[\left(1 - \frac{1}{10}\right) + \left(1 - \frac{1}{100}\right) + \left(1 - \frac{1}{1000}\right) + \dots n \text{ terms} \right] \\ &= \frac{4}{9} \left[(1 + 1 + 1 + \dots n \text{ terms}) - \left(\frac{1}{10} + \frac{1}{100} + \frac{1}{1000} + \dots n \text{ terms}\right) \right] \\ &= \frac{4}{9} \left[n - \frac{1}{10} \left(\frac{1 - \left(\frac{1}{10}\right)^n}{1 - \frac{1}{10}} \right) \right] \\ &= \frac{4}{9} \left[n - \frac{1}{9} \left(1 - \left(\frac{1}{10}\right)^n \right) \right] \end{aligned}$$

ii) $3 + 33 + 333 + \dots$ upto n terms

$$\begin{aligned} &= 3(1 + 11 + 111 + \dots + n \text{ terms}) \\ &= \frac{3}{9}(9 + 99 + 999 + \dots + n \text{ terms}) \\ &= \frac{3}{9}[(10 - 1) + (100 - 1) + (1000 - 1) + \dots n \text{ terms}] \\ &= \frac{3}{9}[(10 + 100 + 1000 + \dots n \text{ terms}) - [(1 + 1 + 1 + \dots n \text{ terms})]] \\ &= \frac{3}{9} \left[10 \cdot \left(\frac{10^n - 1}{n} \right) - n \right] \\ &= \frac{30}{81}(10^n - 1) - \frac{3n}{9} \\ &= \frac{10}{27}(10^n - 1) - \frac{n}{3} \end{aligned}$$

7. Find the sum of the Geometric series $3 + 6 + 12 + \dots + 1536$.

Solution :

Given $3 + 6 + 12 + \dots + 1536$ is a geometric series

$$a = 3, r = 2, t_n = 1536$$

$$\Rightarrow a \cdot r^{n-1} = 1536$$

$$\Rightarrow 3 \cdot 2^{n-1} = 1536$$

$$\Rightarrow 2^{n-1} = \frac{1536}{3}$$

$$\Rightarrow 2^{n-1} = 512 = 2^9$$

$$\therefore n-1 = 9$$

$$n = 10$$

$$\therefore S_n = a \cdot \frac{r^n - 1}{r - 1}$$

$$S_{10} = 3 \times \frac{2^{10} - 1}{2 - 1}$$

$$= 3 (1023)$$

$$= 3069$$

8. Kumar writes a letter to four of his friends. He asks each one of them to copy

sons with the instruction that they continue the process similarly. Assuming that the process is unaltered and it costs ₹ 2 to mail one letter, find the amount spent on postage when 8th set of letters is mailed.

Solution :

By the data given,

The number of mails delivered are

$$4, 4 \times 4, 4 \times 4 \times 4, \dots$$

i.e., 4, 16, 64, 8th set of letters.

Each mail costs ₹ 2

\therefore The total cost is

$$(4 \times 2) + (16 \times 2) + (64 \times 2) + \dots \text{8th set}$$

$$= 8 + 32 + 128 + \dots \text{8th set (which forms)}$$

form a geometric series with $a = 8, r = 4, n = 8$

$$\therefore S_n = a \cdot \frac{r^n - 1}{r - 1}$$

$$S_8 = 8 \cdot \frac{4^8 - 1}{3}$$

$$= 8 \times \frac{65535}{3}$$

$$= 8 \times 21845$$

$$= ₹ 174760$$

9. Find the rational form of the number $0.\overline{123}$.

Solution :

$$\text{Let } x = 0.\overline{123}$$

$$x = 0.123123123 \dots (1)$$

$$\Rightarrow 1000x = 123.123123 \dots$$

$$\Rightarrow 1000x = 123.\overline{123} \dots (2)$$

$$\therefore (2) - (1) \Rightarrow 999x = 123$$

$$\Rightarrow x = \frac{123}{999}$$

$$\therefore x = \frac{41}{333}$$

10. If $S_n = (x + y) + (x^2 + xy + y^2) + (x^3 + x^2y + xy^2 + y^3) + \dots$ n terms then prove that

$$(x - y) S_n = \left| \frac{x^2 (x^n - 1)}{x - 1} - \frac{y^2 (y^n - 1)}{y - 1} \right|$$

Solution :

Given

$$S_n = (x + y) + (x^2 + xy + y^2) + (x^3 + x^2y + xy^2 + y^3) + \dots + n \text{ terms}$$

$$\Rightarrow (x - y) S_n = (x - y) (x + y) + (x - y)$$

$$(x^2 + xy + y^2) + (x - y) (x^3 + x^2y + xy^2 + y^3) + \dots n \text{ terms}$$

$$\Rightarrow (x - y) S_n = (x^2 - y^2) + (x^3 + y^3) + (x^4 - y^4) + \dots n \text{ terms}$$

$$(x^2 + x^3 + x^4 + \dots n \text{ terms})$$

$$- (y^2 + y^3 + y^4 + \dots n \text{ terms}),$$

both of them are geometric series

$$a = x^2, r = x \text{ \& } a = y^2, r = y$$

$$\therefore (x - y) S_n = \frac{x^2(x^n - 1)}{x - 1} - \frac{y^2(y^n - 1)}{y - 1}$$

$$\left(\because S_n = a \cdot \frac{r^n - 1}{r - 1} \right)$$

Hence proved.

IX. SPECIAL SERIES :

Key Points

✓ The sum of first n natural numbers $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$

✓ The sum of squares of first n natural numbers

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

✓ The sum of cubes of first n natural numbers $1^3 + 2^3 + 3^3 + \dots + n^3 = \left[\frac{n(n+1)}{2} \right]^2$

✓ The sum of first n odd natural numbers $1 + 3 + 5 + \dots + (2n - 1) = n^2$.

Example 2.54

Find the value of (i) $1 + 2 + 3 + \dots + 50$ (ii) $16 + 17 + 18 + \dots + 75$

Solution :

(i) $1 + 2 + 3 + \dots + 50$

$$\text{Using, } 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

$$1 + 2 + 3 + \dots + 50 = \frac{50 \times (50+1)}{2} = 1275$$

(ii) $16 + 17 + 18 + \dots + 75$

$$= (1+2+3 + \dots + 75) - (1 + 2 + 3 + \dots + 15)$$

$$= \frac{75(75+1)}{2} - \frac{15(15+1)}{2}$$

$$= 2850 - 120 = 2730$$

Example 2.55

Find the sum of (i) $1 + 3 + 5 + \dots$ + to 40 terms
(ii) $2 + 4 + 6 + \dots + 80$ (iii) $1 + 3 + 5 + \dots + 55$

Solution :

(i) $1 + 3 + 5 + \dots + 40 \text{ terms} = 40^2 = 1600$

(ii) $2 + 4 + 6 + \dots + 80$

$$= 2(1 + 2 + 3 + \dots + 40)$$

$$= 2 \times \frac{40 \times (40+1)}{2} = 1640$$

(iii) $1 + 3 + 5 + \dots + 55$

Here the number of terms is not given. Now we have to find the number of terms using the formula, $n = \frac{(l-a)}{d} + 1$ gives $n = \frac{(55-1)}{2} + 1 = 28$.

Therefore, $1 + 3 + 5 + \dots + 55$

$$= (28)^2 = 784.$$

Example 2.56Find the sum of (i) $1^2 + 2^2 + \dots + 19^2$

(ii) $5^2 + 10^2 + 15^2 + \dots + 105^2$

(iii) $15^2 + 16^2 + 17^2 + \dots + 28^2$

Solution :

$$\begin{aligned} \text{(i)} \quad 1^2 + 2^2 + \dots + 19^2 &= \frac{19 \times (19+1) (2 \times 19 + 1)}{6} \\ &= \frac{19 \times 20 \times 39}{6} = 2470 \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad 5^2 + 10^2 + 15^2 + \dots + 105^2 \\ &= 5^2 (1^2 + 2^2 + 3^2 + \dots + 21^2) \\ &= 25 \times \frac{21 \times (21+1) (2 \times 21 + 1)}{6} \\ &= \frac{25 \times 21 \times 22 \times 43}{6} = 82775 \end{aligned}$$

$$\begin{aligned} &= (1^2 + 2^2 + 3^2 + \dots + 28^2) \\ &\quad - (1^2 + 2^2 + 3^2 + \dots + 14^2) \\ &= \frac{28 \times 29 \times 57}{6} - \frac{14 \times 15 \times 29}{6} \\ &= 7714 - 1015 = 6699 \end{aligned}$$

Example 2.57Find the sum of (i) $1^3 + 2^3 + 3^3 + \dots + 16^3$

(ii) $9^3 + 10^3 + \dots + 21^3$

Solution :

$$\begin{aligned} \text{(i)} \quad 1^3 + 2^3 + 3^3 + \dots + 16^3 &= \left[\frac{16 \times (16+1)}{2} \right]^2 \\ &= (136)^2 = 18496 \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad 9^3 + 10^3 + \dots + 21^3 &= (1^3 + 2^3 + 3^3 + \dots + 21^3) \\ &\quad - (1^3 + 2^3 + 3^3 + \dots + 8^3) \\ &= \left[\frac{21 \times (21+1)}{2} \right]^2 - \left[\frac{8 \times (8+1)}{2} \right]^2 \\ &= (231)^2 - (36)^2 = 52065 \end{aligned}$$

Example 2.58If $1 + 2 + 3 + \dots + n = 666$ then find n .**Solution :**

$$\text{Since, } 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2},$$

$$\text{we have } \frac{n(n+1)}{2} = 666$$

$$n^2 + n - 1332 = 0 \text{ gives } (n + 37) (n - 36) = 0$$

$$\text{So, } n = -37 \text{ or } n = 36$$

But $n \neq -37$ (Since n is a natural number) ;
Hence $n = 36$.

EXERCISE 2.9**1. Find the sum of the following series**

i) $1 + 2 + 3 + \dots + 60$

ii) $3 + 6 + 9 + \dots + 96$

iii) $51 + 52 + 53 + \dots + 92$

iv) $1 + 4 + 9 + 16 + \dots + 225$

v) $6^2 + 7^2 + 8^2 + \dots + 21^2$

vi) $10^3 + 11^3 + 12^3 + \dots + 20^3$

vii) $1 + 3 + 5 + \dots + 71$

Solution :

i) $1 + 2 + 3 + \dots + 60$

$$\begin{aligned} \sum_{k=1}^n K &= \frac{n(n+1)}{2} \\ &= \frac{60 \times 61}{2} \\ &= 30 \times 61 \\ &= 1830 \end{aligned}$$

ii) $3 + 6 + 9 + \dots + 96$

$$= 3 (1 + 2 + 3 + \dots + 32)$$

$$\begin{aligned} &= 3 \left(\frac{32 \times 33}{2} \right) \\ &= 3 \times 16 \times 33 \\ &= 1584 \end{aligned}$$

$$\begin{aligned}
 \text{iii)} \quad & 51 + 52 + 53 + \dots + 92 \\
 &= (1 + 2 + 3 + \dots + 92) \\
 &\quad - (1 + 2 + 3 + \dots + 50) \\
 &= \frac{92 \times 93}{2} - \frac{50 \times 51}{2} \\
 &= 46 \times 93 - 25 \times 51 \\
 &= 4278 - 1275 \\
 &= 3003
 \end{aligned}$$

$$\begin{aligned}
 \text{iv)} \quad & 1 + 4 + 9 + 16 + \dots + 225 \\
 &= 1^2 + 2^2 + 3^2 + \dots + 15^2 \\
 \sum_{k=1}^n K^2 &= \frac{n(n+1)(2n+1)}{6} \\
 &= \frac{15 \times 16 \times 31}{6} \\
 &= 1240
 \end{aligned}$$

$$\begin{aligned}
 \text{v)} \quad & 6^2 + 7^2 + 8^2 + \dots + 21^2 \\
 &= (1^2 + 2^2 + 3^2 + \dots + 21^2) \\
 &\quad - (1^2 + 2^2 + \dots + 5^2) \\
 &= \frac{21 \times 22 \times 43}{6} - \frac{5 \times 6 \times 11}{6} \\
 &= 3311 - 55 \\
 &= 3256
 \end{aligned}$$

$$\begin{aligned}
 \text{vi)} \quad & 10^3 + 11^3 + 12^3 + \dots + 20^3 \\
 &= (1^3 + 2^3 + \dots + 20^3) \\
 &\quad - (1^3 + 2^3 + \dots + 9^3) \\
 \sum_{k=1}^n K^3 &= \left(\frac{n(n+1)}{2} \right)^2 \\
 &= \left(\frac{20 \times 21}{2} \right)^2 - \left(\frac{9 \times 10}{2} \right)^2 \\
 &= (210)^2 - (45)^2 \\
 &= 44100 - 2025 \\
 &= 42075
 \end{aligned}$$

$$\begin{aligned}
 \text{vii)} \quad & 1 + 3 + 5 + \dots + 71 \\
 & a = 1, d = 2, l = 71
 \end{aligned}$$

$$\begin{aligned}
 \therefore n &= \frac{l-a}{d} + 1 \\
 &= \frac{71-1}{2} + 1 \\
 &= 36
 \end{aligned}$$

$$\therefore 1 + 3 + 5 + \dots + 71 = (36)^2$$

$$\begin{aligned}
 (\because 1 + 3 + 5 + \dots + n \text{ terms} &= n^2) \\
 &= 1296
 \end{aligned}$$

2. If $1 + 2 + 3 + \dots + k = 325$, then find $1^3 + 2^3 + 3^3 + \dots + k^3$

Solution :

$$\text{Given } 1 + 2 + 3 + \dots + k = 325$$

$$\Rightarrow \frac{k(k+1)}{2} = 325$$

$$\therefore 1^3 + 2^3 + 3^3 + \dots + k^3$$

$$= \left(\frac{k(k+1)}{2} \right)^2$$

$$= (325)^2$$

$$= 105625$$

3. If $1^3 + 2^3 + 3^3 + \dots + k^3 = 44100$, then find $1 + 2 + 3 + \dots + k$.

Solution :

$$\text{Given } 1^3 + 2^3 + 3^3 + \dots + k^3 = 44100$$

$$\Rightarrow \left(\frac{k(k+1)}{2} \right)^2 = 44100$$

$$\Rightarrow \frac{k(k+1)}{2} = 210$$

$$\Rightarrow 1 + 2 + 3 + \dots + k = 210$$

4. How many terms of the series $1^3 + 2^3 + 3^3 + \dots$ should be taken to get the sum 14400?

Solution :

$$\text{Given } 1^3 + 2^3 + 3^3 + \dots + k^3 = 14400$$

$$\Rightarrow \left(\frac{k(k+1)}{2} \right)^2 = 14400$$

$$\Rightarrow \frac{k(k+1)}{2} = 120$$

$$\Rightarrow k^2 + k - 240 = 0$$

$$\Rightarrow (k+16)(k-15) = 0$$

$$\therefore k = -16, k = 15$$

$$\text{But } k \neq -16$$

$$\therefore k = 15$$

5. The sum of the squares of the first n natural numbers is 285, while the sum of their

Solution :

Give sum of the squares of first ' n ' natural numbers = 285

$$\text{i.e., } \frac{n(n+1)(2n+1)}{6} = 285 \quad \dots\dots\dots (1)$$

and Sum of their cubes = 2025

$$\text{i.e., } \left(\frac{n(n+1)}{2} \right)^2 = 2025$$

$$\Rightarrow n \left(\frac{n+1}{2} \right) = 45 \quad \dots\dots\dots (2)$$

Sub (2) in (1)

$$(1) \Rightarrow \frac{n(n+1)}{2} \times \frac{2n+1}{3} = 285$$

$$\Rightarrow 45 \times \frac{2n+1}{3} = 285$$

$$\Rightarrow 2n+1 = \frac{285}{15} = 19$$

$$\Rightarrow 2n = 18$$

$$\therefore n = 9$$

6. Rekha has 15 square colour papers of sizes 10 cm, 11 cm, 12 cm, ..., 24 cm. How much area can be decorated with these colour papers?

Solution :

Given sides of 15 square Colour papers are

10 cm, 11 cm, 12 cm, 24 cm

$$\begin{aligned} \therefore \text{its area} &= 10^2 + 11^2 + 12^2 + \dots + 24^2 \\ &= (1^2 + 2^2 + 3^2 + \dots + 24^2) - (1^2 + 2^2 + \dots + 9^2) \end{aligned}$$

$$= \frac{24 \times 25 \times 49}{6} - \frac{9 \times 10 \times 19}{6}$$

$$= 4900 - 285$$

$$= 4615 \text{ cm}^2$$

7. Find the sum of the series to $(2^3 - 1^3) + (4^3 - 3^3) + (6^3 - 5^3) + \dots$ to
(i) n terms (ii) 8 terms

Solution :

To find the sum of the series :

$$\begin{aligned} \text{i) } &(2^3 - 1^3) + (4^3 - 3^3) + (6^3 - 5^3) + \dots n \text{ terms} \\ &= (2^3 + 4^3 + 6^3 + \dots n \text{ terms}) \\ &\quad - (1^3 + 3^3 + 5^3 + \dots n \text{ terms}) \end{aligned}$$

$$\sum_1^n (2n)^3 - \sum_1^n (2n-1)^3$$

$$= \sum_1^n [(2n)^3 - (2n-1)^3]$$

$$(\because a^3 - b^3 = (a-b)(a^2 + ab + b^2))$$

$$= \sum_1^n [(2n - 2n + 1)(4n^2 + 2n(2n-1) + (2n-1)^2)]$$

$$= \sum_1^n [4n^2 + 4n^2 - 2n + 4n^2 - 4n + 1]$$

$$= \sum_1^n [12n^2 - 6n + 1]$$

$$\begin{aligned}
 &= 12 \sum n^2 - 6 \sum n + \sum 1 \\
 &= 12 \left[\frac{n(n+1)(2n+1)}{6} \right] - 6 \left[\frac{n(n+1)}{2} \right] + n \\
 &= n(n+1)[4n+2-3] + n \\
 &= (n^2 + n)(4n-1) + n \\
 &= 4n^3 + 4n^2 - n^2 - n + n \\
 &= 4n^3 + 3n^2
 \end{aligned}$$

ii) When $n = 8$,

$$\begin{aligned}
 S_8 &= 4(8^3) + 3(8^2) \\
 &= 4(512) + 3(64) \\
 &= 2048 + 192 \\
 &= 2240
 \end{aligned}$$

EXERCISE 2.10

1. Euclid's division lemma states that for positive integers a and b , there exist unique integers q and r such that $a = bq + r$, where r must satisfy.

- (1) $1 < r < b$ (2) $0 < r < b$
 (3) $0 \leq r < b$ (4) $0 < r \leq b$

Hint : Ans : (3)

By definition of Euclid's lemma $0 \leq r < b$

2. Using Euclid's division lemma, if the cube of any positive integer is divided by 9 then the possible remainders are

- (1) 0, 1, 8 (2) 1, 4, 8
 (3) 0, 1, 3 (4) 1, 3, 5

Hint : Ans : (1)

$$x^3 \equiv y \pmod{9}$$

when $x = 3$, $y = 0$ (27 is div. by 9)

when $x = 4$, $y = 1$ (64 is div. by 9)

when $x = 5$, $y = 8$ (125 is div. by 9)

\therefore The remainders are 0, 1, 8, ...

3. If the HCF of 65 and 117 is expressible in the form of $65m - 117$, then the value of m is

- (1) 4 (2) 2 (3) 1 (4) 3

Hint : Ans : (2)

HCF of 65, 117 is 13

$$65m - 117 = 13$$

$$\Rightarrow 65m = 130$$

$$\Rightarrow m = 2$$

4. The sum of the exponents of the prime factors in the prime factorization of 1729 is

- (1) 1 (2) 2 (3) 3 (4) 4

Hint : Ans : (3)

$$1729 = 7 \times 13 \times 19$$

$$= 7 \times 13 \times 19$$

$$\therefore \text{Sum of the exponents} = 1 + 1 + 1 = 3$$

5. The least number that is divisible by all the numbers from 1 to 10 (both inclusive) is

- (1) 2025 (2) 5220 (3) 5025 (d) 2520

Hint : Ans : (4)

Refer 9th sum in Ex. 2.2

6. $7^{4k} \equiv \underline{\hspace{2cm}} \pmod{100}$

- (1) 1 (2) 2 (3) 3 (4) 4

Hint : Ans : (1)

If $k = 1$, 7^4 leaves remainder 1 modulo 100.

7. Given $F_1 = 1$, $F_2 = 3$ and $F_n = F_{n-1} + F_{n-2}$ then F_5 is

- (1) 3 (2) 5 (3) 8 (4) 11

Hint : Ans : (4)

$$F_3 = F_2 + F_1 = 4$$

$$F_4 = F_3 + F_2 = 7$$

$$F_5 = F_4 + F_3 = 4 + 7 = 11$$

8. The first term of an arithmetic progression is unity and the common difference is 4. Which of the following will be a term of this A.P.

- (1) 4551 (2) 10091
(3) 7881 (4) 13531

Hint : **Ans : (3)**

$$a = 1, d = 4$$

\therefore The A.P is 1, 5, 9, 13, leaves remainder 1 when divided by 4.

\therefore 7881 leaves remainder 1 when divided by 4.

9. If 6 times 6th term of an A.P. is equal to 7 times the 7th term, then the 13th term of the A.P. is

- (1) 0 (2) 6 (3) 7 (4) 13

Hint : **Ans : (1)**

$$6(t_6) = 7(t_7) \Rightarrow 6(a + 5d) = 7(a + 6d)$$

$$\Rightarrow 6a + 30d = 7a + 42d$$

$$\Rightarrow a + 12d = 0$$

$$\Rightarrow t_{13} = 0$$

10. An A.P. consists of 31 terms. If its 16th term is m , then the sum of all the terms of this A.P. is

- (1) 16m (2) 62m (3) 31m (d) $\frac{31}{2}m$

Hint : **Ans : (3)**

$$n = 31, a + 15d = m \quad S_{31} = \frac{31}{2} [2a + 30d]$$

$$= 31(a + 15d)$$

$$= 31m$$

11. In an A.P., the first term is 1 and the common difference is 4. How many terms of the A.P. must be taken for their sum to be equal to 120?

- (1) 6 (2) 7 (3) 8 (4) 9

Hint :

Ans : (3)

$$a = 1, d = 4, S_n = 120$$

$$\Rightarrow \frac{n}{2}(2a + (n-1)d) = 120$$

$$\Rightarrow \frac{n}{2}(2 + (n-1)4) = 120$$

$$\Rightarrow n(1 + 2n - 2) = 120$$

$$\Rightarrow n(2n - 1) = 120$$

$$\Rightarrow 2n^2 - n - 120 = 0$$

$$\Rightarrow n = 8$$

-1	-240
-16	15
2	2
-8,	15
	2

12. If $A = 2^{65}$ and $B = 2^{64} + 2^{63} + 2^{62} + \dots + 2^0$ which of the following is true ?

- (1) B is 2^{64} more than A
(2) A and B are equal
(3) B is larger than A by 1

Hint :

Ans : (4)

2^4 is greater than $2^0 + 2^1 + 2^2 + 2^3$ by 1

2^5 is greater than $2^0 + 2^1 + 2^2 + 2^3 + 2^4$ by 1

$\therefore 2^{65}$ is greater than $2^0 + 2^1 + \dots + 2^{64}$ by 1

\therefore A is larger than B by 1.

13. The next term of the sequence

$$\frac{3}{16}, \frac{1}{8}, \frac{1}{12}, \frac{1}{18}, \dots \text{ is}$$

- (1) $\frac{1}{24}$ (2) $\frac{1}{27}$ (3) $\frac{2}{3}$ (4) $\frac{1}{81}$

Hint :

Ans : (2)

$$r = \frac{\frac{1}{8}}{\frac{3}{16}} = \frac{1}{8} \times \frac{16}{3} = \frac{2}{3}$$

$$\therefore \text{Next term of the sequence} = \frac{1}{18} \times \frac{2}{3} = \frac{1}{27}$$

14. If the sequence t_1, t_2, t_3, \dots are in A.P. then the sequence $t_6, t_{12}, t_{18}, \dots$ is

- (1) a Geometric Progression
- (2) an Arithmetic Progression
- (3) neither an Arithmetic Progression nor a Geometric Progression
- (4) a constant sequence

Hint : **Ans : (3)**

Obviously they should be in A.P.

15. The value of $(1^3 + 2^3 + 3^3 + \dots + 15^3) - (1 + 2 + 3 + \dots + 15)$ is

- (1) 14400
- (2) 14200
- (3) 14280
- (4) 14520

Hint : **Ans : (3)**

$$\left(\frac{15 \times 16}{2}\right)^2 - \frac{15 \times 16}{2}$$

$$= 14400 - 120$$

$$= 14280$$

UNIT EXERCISE - 2

1. Prove that $n^2 - n$ divisible by 2 for every positive integer n .

Solution :

Any positive integer is of the form $2q$ (or) $2q + 1$ for some integer q .

- i) $n^2 - n = (2q)^2 - 2q$

$$= 2q(2q - 1)$$

which is divisible by 2.
- ii) $n^2 - n = (2q + 1)^2 - (2q + 1)$

$$= (2q + 1)(2q + 1 - 1)$$

$$= 2q(2q + 1), \text{ which is divisible by 2.}$$

Hence proved.

2. A milk man has 175 litres of cow's milk and 105 litres of buffalow's milk. He wishes to sell the milk by filling the two types of milk in cans of equal capacity. Calculate the following (i) Capacity of a can (ii) Number of cans of cow's milk (iii) Number of cans of buffalow's milk.

Solution :

Cow's milk = 175 lrs.

Buffalow's milk = 105 lrs.

Since he wish to sell the milk by filling the 2 types of milk in cans of equal capacity,

- i) Capacity of a can = HCF of 175 and 105

$$= 35 \text{ litres}$$

- ii) Number of cans of Cow's milk = $\frac{175}{35} = 5$

- iii) Number of cans of buffalow's milk

$$= \frac{105}{35} = 3$$

3. When the positive integers a, b and c are divided by 13 the respective remainders are 9, 7 and 10. Find the remainder when $a + 2b + 3c$ is divided by 13.

Solution :

$$\text{Let } a = 13q + 9$$

$$b = 13q + 7 \Rightarrow 2b = 26q + 14$$

$$c = 13q + 10 \Rightarrow 3c = 39q + 30$$

$$a + 2b + c = (13q + 9) + (26q + 14) + (39q + 30)$$

$$= 78q + 53$$

$$= 13(6q) + 13(4) + 1$$

\therefore When $a + 2b + 3c$ is divided by 13, the remainder is 1.

4. Show that 107 is of the form $4q + 3$ for any integer q .

Solution :

When 107 is divided by 4,

$$107 = 4(26) + 3$$

This is of the form

$$107 = 4q + 3 \text{ for } q = 26.$$

5. If $(m + 1)^{\text{th}}$ term of an A.P. is twice the $(n + 1)^{\text{th}}$ term, then prove that $(3m + 1)^{\text{th}}$ term is twice the $(m + n + 1)^{\text{th}}$ term.

Solution :

$$\text{Given } t_{m+1} = 2(t_{n+1})$$

$$a + (m + 1 - 1)d = 2(a + (n + 1 - 1)d)$$

$$a + md = 2(a + nd)$$

$$a + md = 2a + 2nd \quad \text{--- (1)}$$

$$\text{To Prove : } t_{3m+1} = 2(t_{m+n+1})$$

$$\text{LHS : } t$$

$$= a + (3m + 1 - 1)d$$

$$= a + 3md$$

$$= (a + md) + 2md$$

$$= 2a + 2nd + 2md \quad (\text{from (1)})$$

$$= 2[a + (m + n)d]$$

$$= 2[t_{m+n+1}]$$

$$= \text{RHS}$$

Hence proved.

6. Find the 12th term from the last term of the A.P. $-2, -4, -6, \dots -100$.

Solution :

Given A.P. is $-2, -4, -6, \dots -100$

To find : t_{12} from the last term

$$a = -100, d = 2$$

$$t_{12} = a + 11d$$

$$= -100 + 11(2)$$

$$= -100 + 22$$

$$= -78$$

7. Two A.P.'s have the same common difference. The first term of one A.P. is 2 and that of the other is 7. Show that the difference between their 10th terms is the same as the difference between their 21st terms, which is the same as the difference between any two corresponding terms.

Solution :

1st A.P.

2nd A.P.

$$a = 2, d = d$$

$$a = 7, d = d$$

$$t_{10} = a + 9d$$

$$T_{10} = a + 9d$$

$$= 2 + 9d$$

$$= 7 + 9d$$

$$t_{21} = a + 20d$$

$$T_{21} = a + 20d$$

$$= 2 + 20d$$

$$= 7 + 20d$$

$$\therefore T_{10} - t_{10} = 5 \text{ and } T_{21} - t_{21} = 5 = T_n - t_n = 5$$

8. A man saved ₹16500 in ten years. In each

than he did in the preceding year. How much did he save in the first year ?

Solution :

Given $S_n = ₹ 16500$, $d = ₹ 100$, $n = 10$ in A.P.

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$\Rightarrow \frac{10}{2}[2a + 9(100)] = 16500$$

$$\Rightarrow 2a + 900 = \frac{16500}{5}$$

$$\Rightarrow 2a + 900 = 3300$$

$$\Rightarrow 2a = 2400$$

$$\therefore a = 1200$$

\therefore He saved Rs. 1200 in 1st year.

9. Find the G.P. in which the 2nd term is $\sqrt{6}$ and the 6th term is $9\sqrt{6}$.

Solution :

Given $t_2 = \sqrt{6}$, $t_6 = 9\sqrt{6}$ in G.P.

$$\Rightarrow a.r = \sqrt{6}, \quad a.r^5 = 9\sqrt{6}$$

$$\therefore r^4 = 9 \quad (\text{when divide})$$

$$\Rightarrow r = \sqrt{3}$$

$$\therefore a \times \sqrt{3} = \sqrt{6}$$

$$\therefore a = \sqrt{2}$$

\therefore The G.P is

$$\sqrt{2}, \sqrt{6}, \sqrt{18}, \dots$$

(or)

$$\sqrt{2}, \sqrt{6}, 3\sqrt{2}, \dots$$

10. The value of a motor cycle depreciates at

the value of the motor cycle 3 year hence, which is now purchased for ₹ 45,000 ?

Solution :

$P = ₹ 45000$, $n = 3$, $r = 15\%$ (depreciation)

$$A = P \left(1 - \frac{r}{100} \right)^n$$

$$= 45,000 \left(1 - \frac{15}{100} \right)^3$$

$$= 45,000 \times \frac{85}{100} \times \frac{85}{100} \times \frac{85}{100}$$

$$= 27,635.625$$

$$= 27636$$

PROBLEMS FOR PRACTICE

1. Express the number $0.\overline{3178}$ in the form of $\frac{a}{b}$.

(Ans: $\frac{3178}{999}$)

2. A class of 20 boys and 15 girls is divided into n groups so that each group has x boys and y girls. Find x , y and n .

(Ans : $n = 7, x = 4, y = 3$)

3. Show that 7^n cannot end with digit zero for any natural number.

4. Find the HCF of the following numbers by using Euclid's division algorithm.

i) 867, 255

ii) 1656, 4025

iii) 180, 252, 324

iv) 92690, 7378

v) 134791, 6341, 6339

(Ans : i) 51 ii) 23 iii) 36 iv) 31 v) 1)

5. Use Euclid's lemma, show that the square of any positive integer is either of the form $3m$ (or) $3m+1$ for same integer m .

6. Find the largest number which divides 70 and 125 leaving remainder 5 and 8 respectively.

(Ans : 13)

7. Find the largest positive integer that will divide 398, 436 and 542 that leaves remainders 7, 11, 15 respectively.

(Ans : 17)

8. If HCF of 144 and 180 is expressed in the form $13m - 3$, find m .

(Ans : 3)

9. If d is the HCF of 56 and 72, find x and y satisfying $d = 56x + 72y$. Also show that x and y is not unique.

(Ans : 4 and -3, -68, 53)

10. Find the largest number of four digits exactly divisible by 12, 15, 18 and 27.

(Ans : 9720)

11. Write the first 6 terms of the sequence whose n^{th} term is

$$\text{i) } a_n = \begin{cases} n & , \text{ if } n = 1, 2, 3 \\ a_{n-1} + a_{n-2} + a_{n-3}, & \text{ if } n > 3 \end{cases}$$

$$\text{ii) } a = \frac{3n-2}{3^{n-1}}$$

(Ans : i) 1, 2, 3, 5, 8, 13 ii) $1, \frac{4}{3}, \frac{7}{9}, \frac{10}{27}, \dots$)

following :

$$\text{i) } a_n = (-1)^n \cdot 2^{n+3} (n+1) ; a_5, a_8$$

$$\text{ii) } a_n = (-1)^n, (1 - n + n^2) ; a_2, a_9$$

(Ans : i) -1536, 18432 ii) 3, -73)

13. How many terms are there in the A.P.

$$-1, \frac{-5}{6}, \frac{-2}{3}, \dots, \frac{10}{3} ?$$

(Ans : 27)

14. Find the 40th term of an A.P whose 5th term is 41 and 11th term is 71.

(Ans : 216)

15. If 7th term of an A.P is $\frac{1}{9}$ and 9th term is $\frac{1}{7}$.

(Ans : 1)

16. Find the middle term of the A.P 213, 205, 197, ...37

(Ans :125)

17. The first term of an A.P is 5, the last term is 45. Sum of all its terms is 400. Find the number of terms and the common difference of A.P.

(Ans : $n = 16, d = \frac{8}{3}$)

18. The 24th term of an A.P is twice its 10th term. Show that its 72nd term is 4 times its 15th term.

19. Find the sum of all two digit odd positive numbers.

(Ans : 2475)

20. Which term of the sequence $20, 19\frac{1}{4}, 18\frac{1}{2}, \dots$ is the first negative term ?

(Ans : 28)

21. Find the 18th term of the A.P from right end 3, 7, 11, 407.

(Ans : 339)

22. How many consecutive integers beginning with 10 must be taken for their sum to be 2035 ?

(Ans : 55)

23. Sum of 3 numbers in an A.P is 54 and their product is 5670. Find the 3 numbers.

(Ans : 15, 18, 21)

24. Find the sum of all natural numbers between 201 and 399 that are divisible by 5.

25. Find : $\left(4 - \frac{1}{n}\right) + \left(7 - \frac{2}{n}\right) + \left(10 - \frac{3}{n}\right) + \dots$ up to n terms.

(Ans : $\frac{3n^2 + 4n - 1}{2}$)

26. The sum of first 'n' terms of an A.P is $5n - n^2$. Find the n^{th} term of the A.P.

(Ans : $-2(n - 3)$)

27. If 7 times the 7^{th} term of an A.P is equal to 11 times its 11^{th} term, show that its 18^{th} term is 0.

28. Find the sum of 22 terms of the A.P $x + y, x - y, x - 3y, \dots$

(Ans : $22(x - 20y)$)

29. If the ratio between the sums of n terms of AP's is $(7n + 1) : (4n + 27)$, find the ratio of their 11^{th} terms.

(Ans : $148 : 111$)

30. A man gets the initial salary of 5200 p.m. and receive an automatic increase of 320 in the very next month and each month hereafter.

earings during the 1st year.

(Ans : 8080, 83520)

31. Sum of 3 terms of a G.P is $\frac{39}{10}$ and their product is 1. Find the G.P.

(Ans : $\frac{5}{2}, 1, \frac{2}{5}$)

32. If 4^{th} and 7^{th} terms of a G.P are 54 and 1458 respectively, find G.P.

(Ans : 2, 6, 18, 54,)

33. In the series 18, -12, 8,, which term is $\frac{512}{729}$?

(Ans : 9^{th} term)

34. Find the sum : $0.2 + 0.92 + 0.992 + \dots$ to n terms.

35. Find the sum of the series :

$$9^{\frac{1}{3}} 9^{\frac{1}{9}} 9^{\frac{1}{27}} \dots \infty.$$

(Ans : 3)

36. A farmer buys a used tractor for Rs. 12000. He pays Rs. 6000 and agrees to pay the balance in annual instalment of Rs. 500 plus 12% interest on unpaid amount. How much will the tractor cost him ?

(Ans : Rs. 16,680)

37. The sum of an infinite geometric series is 15 and the sum of the squares of these terms is 45. Find the series.

(Ans : $5 + \frac{10}{3} + \frac{20}{9} + \dots$)

38. Evaluate : $1^2 + 3^2 + 5^2 + \dots + 29^2$

(Ans : 4495)

39. Evaluate : $8^3 + 9^3 + \dots + 17^3$

40. The sum of the squares of first 'n' natural numbers is 285, while the sum of their cubes is 2025. Find 'n'.

(Ans : 9)

OBJECTIVE TYPE QUESTIONS

1. When the denominator number of $\frac{257}{500}$ writes in the form of $2^m \times 5^n$, then $m + n$ is

- (a) 6 (b) 5
(c) 23 (d) none of these

Ans : (b)

2. The common difference of the A.P $\frac{1}{2q}, \frac{1-2q}{2q}, \frac{1-4q}{2q}, \dots$

- (a) 0 (b) 1 (c) -1 (d) q

Ans : (c)

3. The first 3 terms of A.P are $3y - 1, 3y + 5$ and $5y + 1$, then y is

- (a) 5 (b) 1 (c) -5 (d) 4

Ans : (a)

4. For an A.P, $S_n = n^2 - n + 1$, the 2nd term is
 (a) 2 (b) 3 (c) 4 (d) -2

Ans : (b)

5. The next term of an A.P :
 $-12, -9, -6, -3, \dots$ is
 (a) 3 (b) 6
 (c) 0 (d) none of these

Ans : (c)

6. The sum of 6 terms of the A.P 1, 3, 5, 7, is
 (a) 25 (b) 49 (c) 36 (d) 30

Ans : (c)

7. Which term of the series $-3, -1, 5, \dots$ is 53 ?
 (a) 12 (b) 13 (c) 14 (d) 15

Ans : (d)

8. The common ratio of the G.P $\frac{-5}{2}, \frac{25}{4}, \frac{-125}{8}, \dots$ is
 (a) 15 (b) $\frac{35}{4}$ (c) $\frac{-5}{2}$ (d) $\frac{5}{2}$

Ans : (c)

9. If the 3rd term of G.P is 4, then the product of its first 5 terms is
 (a) 4^3 (b) 4^5 (c) 4^4 (d) 4^2

Ans : (b)

10. If a, b, c are in A.P, a, b, d in G.P, then $a, a-b, d-c$ will be in
 (a) A.P (b) G.P
 (c) A.P and G.P (d) none of these

Ans : (b)

11. The 3rd term of a G.P is the square of first term. If the 2nd term is 8, then the 6th term is

(a) 120 (b) 124 (c) 128 (d) 132

Ans : (d)

12. If $a_1 = a_2 = 2$, $a_n = a_{n-1} - 1$, then a_5 is

(a) 1 (b) -1 (c) 0 (d) -2

Ans : (b)

13. The 9th term of the series $27 + 9 + 5\frac{2}{5} + \dots$ is

(a) $1\frac{10}{17}$ (b) $\frac{10}{17}$ (c) $\frac{16}{27}$ (d) $\frac{17}{27}$ **(Ans : (a))**

14. The n^{th} term of the series $3.8 + 6.11 + 9.14 + 12.17 + \dots$ will be

(c) $n(3n + 5)$ (d) $3n(3n + 5)$ **Ans : (d)**

15. If the n^{th} term of a G.P $5, \frac{-5}{2}, \frac{5}{4}, \dots$ is $\frac{5}{1024}$, then n is

(a) 11 (b) 10 (c) 9 (d) 4

Ans : (a)

16. If $1 + 2 + 3 + \dots + n = K$, then $1^3 + 2^3 + 3^3 + \dots + n^3$ is

(a) K^3 (b) K^2 (c) $\frac{K(K+1)}{2}$ (d) $(K+1)^3$ **Ans : (b)**

CHAPTER 3

ALGEBRA

I. SIMULTANEOUS LINEAR EQUATIONS IN THREE VARIABLES :

Key Points

- ✓ Any first degree equation containing two variables x and y is called a linear equation in two variables. The general form of linear equation in two variables x and y is $ax + by + c = 0$, where atleast one of a, b is non-zero and a, b, c are real numbers.
- ✓ The general form of a linear equation in three variables x, y and z is $ax + by + cz + d = 0$ where a, b, c, d are real numbers, and atleast one of a, b, c is non-zero.
- ✓ A linear equation in two variables of the form $ax + by + c = 0$, represents a straight line.
- ✓ A linear equation in three variables of the form $ax + by + cz + d = 0$, represents a plane.

The father's age is six times his son's age. Six years hence the age of father will be four times his son's age. Find the present ages (in years) of the son and father.

Solution :

Let the present age of father be x years and the present age of son be y years

Given, $x = 6y$

— (1)

$x + 6 = 4(y + 6)$

— (2)

Substituting (1) in (2), $6y + 6 = 4(y + 6)$

$6y + 6 = 4y + 24$ gives, $y = 9$

Therefore, son's age = 9 years and father's age = 54 years.

Solve $2x - 3y = 6$, $x + y = 1$

Solution :

$2x - 3y = 6$ — (1)

$x + y = 1$ — (2)


(1) $\times 1$ gives, $2x - 3y = 6$

(2) $\times 2$ gives, $2x + 2y = 2$

$-5y = 4$ gives, $y = \frac{-4}{5}$

Substituting $y = \frac{-4}{5}$ in (2), $x - \frac{4}{5} = 1$ we get,

$x = \frac{9}{5}$



Therefore $x = \frac{9}{5}$, $y = \frac{-4}{5}$

Example 3.3

Solve the following system of linear equations in three variables

$$3x - 2y + z = 2, 2x + 3y - z = 5, x + y + z = 6.$$

Solution :

$$3x - 2y + z = 2 \text{ --- (1)}$$

$$2x + 3y - z = 5 \text{ --- (2)}$$

$$x + y + z = 6 \text{ --- (3)}$$

Adding (1) and (2), $3x - 2y + z = 2$

$$\begin{array}{r} 2x + 3y - z = 5 \quad (+) \\ \hline 5x + y = 7 \quad \text{--- (4)} \end{array}$$

Adding (2) and (3), $2x + 3y - z = 5$

$$\begin{array}{r} x + y + z = 6 \quad (+) \\ \hline \end{array}$$

$$4 \times (4) - (5) \quad 20x + 4y = 28$$

$$\begin{array}{r} 3x + 4y = 11 \quad (-) \\ \hline 17x = 17 \quad \text{gives, } x = 1 \end{array}$$

Substituting $x = 1$ in (4), $5 + y = 7$ gives, $y = 2$

Substituting $x = 1$, $y = 2$ in (3), $1 + 2 + z = 6$ we get, $z = 3$

$$\text{Therefore, } x = 1, y = 2, z = 3$$

Example 3.4

In an interschool athletic meet, with 24 individual events, securing a total of 56 points, a first place secures 5 points, a second place secures 3 points, and a third place secures 1 point. Having as many third place finishers as first and second place finishers, find how many athletes finished in each place.

Solution :

Let the number of I, II and III place finishers be x , y and z respectively.

Total number of events = 24;

Total number of points = 56.

Hence, the linear equations in three variables are

$$x + y + z = 24 \text{ --- (1)}$$

$$5x + 3y + z = 56 \text{ --- (2)}$$

$$x + y = z \text{ --- (3)}$$

Substituting (3) in (1) we get, $z + z = 24$ gives, $z = 12$

Therefore, (3) equation will be, $x + y = 12$

$$(2) \text{ is } 5x + 3y = 44$$

$$3 \times (3) \text{ is } 3x + 3y = 36 \quad (-)$$

$$\begin{array}{r} 2x = 8 \quad \text{we get, } x = 4 \end{array}$$

Substituting $x = 4$, $z = 12$ in (3) we get, $y = 12 - 4 = 8$

Therefore, Number of first place finishers is 4

Number of second place finishers is 8

Number of third place finishers is 12.

Example 3.5

Solve $x + 2y - z = 5$; $x - y + z = -2$;

$$-5x - 4y + z = -11$$

Solution :

$$\text{Let, } x + 2y - z = 5 \text{ --- (1)}$$

$$x - y + z = -2 \text{ --- (2)}$$

$$-5x - 4y + z = -11 \text{ --- (3)}$$

Adding (1) and (2) we get,

$$x + 2y - z = 5$$

$$x - y + z = -2 \quad (+)$$

$$\begin{array}{r} 2x + y = 3 \end{array} \quad \text{--- (4)}$$

Subtracting (2) and (3),

$$\begin{array}{rcl}
 x - y + z & = & -2 \\
 -5x - 4y + z & = & -11 \quad (-) \\
 \hline
 6x + 3y & = & 9 \quad \text{--- (5)}
 \end{array}$$

Dividing by 3 $2x + y = 3$

Subtracting (4) and (5),

$$\begin{array}{rcl}
 2x + y & = & 3 \\
 2x + y & = & 3 \\
 \hline
 0 & = & 0
 \end{array}$$

Here we arrive at an identity $0 = 0$.

Hence the system has an infinite number of solutions.

Example 3.6

Solve $3x + y - 3z = 1$; $-2x - y + 2z = 1$;

Solution :

$$\text{Let } 3x + y - 3z = 1 \text{ --- (1)}$$

$$-2x - y + 2z = 1 \text{ --- (2)}$$

$$-x - y + z = 2 \text{ --- (3)}$$

Adding (1) and (2), $3x + y - 3z = 1$

$$\begin{array}{rcl}
 -2x - y + 2z & = & 1 \quad (+) \\
 \hline
 x - z & = & 2 \quad \text{--- (4)}
 \end{array}$$

Adding (1) and (3), $3x + y - 3z = 1$

$$\begin{array}{rcl}
 -x - y + z & = & 2 \quad (+) \\
 \hline
 2x - 2z & = & 3 \quad \text{--- (5)}
 \end{array}$$

Now, (5) - 2 × (4) we get,

$$\begin{array}{rcl}
 2x - 2z & = & 3 \\
 2x - 2z & = & 4 \quad (-) \\
 \hline
 0 & = & -1
 \end{array}$$

Here we arrive at a contradiction as $0 \neq -1$.

This means that the system is inconsistent and has no solution.

Example 3.7

Solve $\frac{x}{2} - 1 = \frac{y}{6} + 1 = \frac{z}{7} + 2$; $\frac{y}{3} + \frac{z}{2} = 13$

Solution :

Considering, $\frac{x}{2} - 1 = \frac{y}{6} + 1$

$$\frac{x}{2} - \frac{y}{6} = 1 + 1 \text{ gives,}$$

$$\frac{6x - 2y}{12} = 2 \text{ we get, } 3x - y = 12 \quad \text{--- (1)}$$

Considering, $\frac{x}{2} - 1 = \frac{z}{7} + 2$

$$\frac{x}{2} - \frac{z}{7} = 1 + 2 \text{ gives,}$$

$$\frac{7x - 2z}{14} = 3 \text{ we get, } 7x - 2z = 42 \quad \text{--- (2)}$$

Also, from $\frac{y}{3} + \frac{z}{2} = 13$ gives,

$$\frac{2y + 3z}{6} = 13 \text{ we get, } 2y + 3z = 78 \quad \text{--- (3)}$$

Eliminating z from (2) and (3)

$$(2) \times 3 \text{ gives, } 21x - 6z = 126$$

$$\begin{array}{rcl}
 (3) \times 2 \text{ gives, } & 4y + 6z & = 156 \quad (+) \\
 \hline
 & 21x + 4y & = 282
 \end{array}$$

$$\begin{array}{rcl}
 (1) \times 4 \text{ gives, } & 12x - 4y & = 48 \quad (+) \\
 \hline
 & 33x & = 330 \quad \text{so, } x = 10
 \end{array}$$

Substituting $x = 10$ in (1), $30 - y = 12$ we get, $y = 18$

Substituting $x = 10$ in (2), $70 - 2z = 42$ then, $z = 14$

Therefore, $x = 10$, $y = 18$, $z = 14$.

Example 3.8

Solve :

$$\frac{1}{2x} + \frac{1}{4y} - \frac{1}{3z} = \frac{1}{4}; \frac{1}{x} = \frac{1}{3y}; \frac{1}{x} - \frac{1}{5y} + \frac{4}{z} = 2\frac{2}{15}$$

Solution :

$$\text{Let } \frac{1}{x} = p, \frac{1}{y} = q, \frac{1}{z} = r$$

The given equations are written as

$$\frac{p}{2} + \frac{q}{4} - \frac{r}{3} = \frac{1}{4}$$

$$p = \frac{q}{3}$$

$$p - \frac{q}{5} + 4r = 2\frac{2}{15} = \frac{32}{15}$$

By simplifying we get,

$$6p + 3q - 4r = 3 \quad \text{--- (1)}$$

$$3p = q \quad \text{--- (2)}$$

$$15p - 3q + 60r = 32 \quad \text{--- (3)}$$

Substituting (2) in (1) and (3) we get,

$$15p - 4r = 3 \quad \text{--- (4)}$$

$$6p + 60r = 32 \text{ reduces to}$$

$$3p + 30r = 16 \quad \text{--- (5)}$$

Solving (4) and (5),

$$15p - 4r = 3$$

$$15p + 150r = 80 \quad (-)$$

$$\begin{array}{r} 15p + 150r = 80 \\ -154r = -77 \end{array} \quad \text{we get, } r = \frac{1}{2}$$

$$\text{Substituting } r = \frac{1}{2} \text{ in (4) we get, } 15p - 2 =$$

$$3 \text{ gives, } p = \frac{1}{3}$$

$$\text{From (2), } q = 3p \text{ we get } q = 1$$

$$\text{Therefore, } x = \frac{1}{p} = 3, y = \frac{1}{q} = 1, z = \frac{1}{r} = 2.$$

That is, $x = 3, y = 1, z = 2$.**Example 3.9**

The sum of thrice the first number, second number and twice the third number is 5. If thrice the second number is subtracted from the sum of first number and thrice the third we get 2. If the third number is subtracted from the sum of twice the first, thrice the second, we get 1. Find the numbers.

Solution :Let the three numbers be x, y, z

From the given data we get the following equations,

$$3x + y + 2z = 5 \quad \text{--- (1)}$$

$$x + 3z - 3y = 2 \quad \text{--- (2)}$$

$$2x + 3y - z = 1 \quad \text{--- (3)}$$

$$(2) \times 3 \text{ gives, } 3x - 9y + 9z = 6 \quad (-)$$

$$\begin{array}{r} 10y - 7z = -1 \end{array} \quad \text{--- (4)}$$

$$(1) \times \text{ gives, } 6x + 2y + 4z = 10$$

$$(3) \times 3 \text{ gives, } 6x + 9y - 3z = 3 \quad (-)$$

$$\begin{array}{r} -7y + 7z = 7 \end{array}$$

$$\text{Adding (4) and (5), } 10y - 7z = -1$$

$$\begin{array}{r} -7y + 7z = 7 \end{array}$$

$$\begin{array}{r} 3y = 6 \end{array} \text{ gives } y = 2$$

Substituting $y = 2$ in (5), $-14 + 7z = 7$ gives,

$$z = 3$$

Substituting $y = 2$ and $z = 3$ in (1),

$$3x + 2 + 6 = 5 \text{ we get } x = -1$$

Therefore, $x = -1, y = 2, z = 3$.

EXERCISE 3.1**1. Solve the following system of linear equations in three variables**

(i) $x + y + z = 5; 2x - y + z = 9; x - 2y + 3z = 16$

(ii) $\frac{1}{x} - \frac{2}{y} + 4 = 0; \frac{1}{y} - \frac{1}{z} + 1 = 0; \frac{2}{z} + \frac{3}{x} = 14$

(iii) $x + 20 = \frac{3y}{2} + 10 = 2z + 5 = 110 - (y + z)$

Solution:

i) Given $x + y + z = 5$ — (1)

$2x - y + z = 9$ — (2)

$x - 2y + 3z = 16$ — (3)

(1) - (3) $\Rightarrow 3y - 2z = -11$ — (4)

(2) $\Rightarrow 2x - y + z = 9$

Subtracting $\frac{-3y - z = -1}{-3y - z = -1}$ — (5)

Solving (4) & (5)

$3y - 2z = -11$

$-3y - z = -1$

Adding $\frac{-3z = -12}{-3z = -12}$

$z = 4$

Sub $z = 4$ in (5)

$-3y - 4 = -1$

$\Rightarrow -3y = 3$

$\Rightarrow y = -1$

sub $y = -1, z = 4$ in (1)

$\Rightarrow x - 1 + 4 = 5$

$\Rightarrow x = 2$

 \therefore Solution set :

$x = 2, y = -1, z = 4$

(ii) $\frac{1}{x} - \frac{2}{y} + 4 = 0$

$\frac{1}{y} - \frac{1}{z} + 1 = 0$

$\frac{2}{z} + \frac{3}{x} = 14$

Let $\frac{1}{x} = a, \frac{1}{y} = b, \frac{1}{z} = c$

$\therefore a - 2b = -4$ — (1)

$b - c = -1$ — (2)

$2c + 3a = 14$ — (3)

Solving (1) & (2)

(1) $\times 1 \Rightarrow a - 2b = -4$

(2) $\times 2 \Rightarrow 2b - 2c = -2$

Adding $\frac{a - 2c = -6}{a - 2c = -6}$ — (4)

Solving (3) & (4)

$3a + 2c = 14$

$a - 2c = -6$

Adding $\frac{4a = 8}{4a = 8}$

$a = 2$

$\therefore (1) \Rightarrow -2b = -6$

$b = 3$

(2) $\Rightarrow 3 - c = -1$

$\Rightarrow c = 4$

$\therefore a = 2, b = 3, c = 4$

$\Rightarrow x = \frac{1}{2}, y = \frac{1}{3}, z = \frac{1}{4}$

 \therefore Solution set :

$\left\{ \frac{1}{2}, \frac{1}{3}, \frac{1}{4} \right\}$

$$(iii) \quad x + 20 = \frac{3y}{2} + 10 = 2z + 5 = 110 - (y + z)$$

$$x + 20 = \frac{3y}{2} + 10$$

$$\Rightarrow 2x + 40 = 3y + 20$$

$$\Rightarrow \boxed{2x - 3y = -20} \quad \dots\dots\dots (1)$$

$$\frac{3y}{2} + 10 = 2z + 5$$

$$\Rightarrow 3y + 20 = 4z + 10$$

$$\Rightarrow \boxed{3y - 4z = -10} \quad \dots\dots\dots (2)$$

$$2z + 5 = 110 - y - z$$

$$\Rightarrow \boxed{3z + y = 105} \quad \dots\dots\dots (3)$$

Solving (1) & (2)

$$3y - 4z = -10$$

$$\text{Adding } 2x - 4z = -30 \quad \dots\dots\dots (4)$$

Solving (2) & (3)

$$(2) \Rightarrow 3y - 4z = -10$$

$$(3) \times 3 \Rightarrow \frac{9z + 3y = 315}{\text{Subtracting, } -13z = -325}$$

$$z = 25$$

Sub $z = 25$ in (4)

$$2x - 100 = -30$$

$$\Rightarrow 2x = 70$$

$$\Rightarrow x = 35$$

Sub $x = 35$ in (1)

$$70 - 3y = -20$$

$$3y = 90$$

$$\therefore y = 30$$

$$\therefore \text{Solution : } x = 35, y = 30, z = 35$$

2. Discuss the nature of solutions of the following system of equations

i) $x + 2y - z = 6$; $-3x - 2y + 5z = -12$;

$$x - 2z = 3$$

ii) $2y + z = 3(-x + 1)$; $-x + 3y - z = -4$

$$3x + 2y + z = -\frac{1}{2}$$

iii) $\frac{y+z}{4} = \frac{z+x}{3} = \frac{x+y}{2}$; $x + y + z = 27$

Solution :

Given $x + 2y - z = 6$ — (1)

$$-3x - 2y + 5z = -12 \quad \text{--- (2)}$$

$$x - 2z = 3 \quad \text{--- (3)}$$

Adding (1) & (2) $\Rightarrow -2x + 4z = -6$

$$\div \text{ by } (-2) \Rightarrow x - 2z = 3 \quad \dots\dots\dots (4)$$

Subtracting (3) & (4)

$$x - 2z = 3$$

$$\frac{x - 2z = 3}{0 = 0}$$

\therefore The system of equation has infinite number of solutions.

ii) Given

$$2y + z = 3(-x + 1) \Rightarrow$$

$$3x + 2y + z = 3 \quad \dots\dots\dots (1)$$

$$-x + 3y - z = -4 \quad \dots\dots\dots (2)$$

$$3x + 2y + z = -\frac{1}{2} \quad \dots\dots\dots (3)$$

Solving (1) & (2)

$$(1) + (2) \Rightarrow 2x + 5y = -1 \quad \dots\dots\dots (4)$$

Solving (2) + (3)

$$(2) + (3) \Rightarrow 2x + 5y = -\frac{9}{2} \quad \dots\dots\dots (5)$$

Solving (4) & (5)

$$\begin{array}{rcl}
 (4) - (5) & \Rightarrow & 2x + 5y = -1 \\
 & & 2x + 5y = -\frac{9}{2} \\
 \hline
 & & 0 = \frac{9}{2} - 1 \\
 & & 0 = \frac{7}{2}
 \end{array}$$

This is a contradiction

$$\text{Since } 0 \neq \frac{7}{2}$$

\therefore The system has no solution.

iii) Given $\frac{y+z}{4} = \frac{z+x}{3} = \frac{x+y}{2}$; $x + y + z = 27$

$$\Rightarrow \frac{y+z}{4} = \frac{z+x}{3} \quad \& \quad \frac{z+x}{3} = \frac{x+y}{2}$$

$$\Rightarrow 3y + 3z = 4z + 4x \quad \& \quad 2z + 2x = 3x + 3y$$

$$\Rightarrow 4x - 3y + z = 0 \quad \& \quad x + 3y - 2z = 0$$

$$\therefore 4x - 3y + z = 0 \quad \dots\dots\dots (1)$$

$$x + 3y - 2z = 0 \quad \dots\dots\dots (2)$$

$$x + y + z = 27 \quad \dots\dots\dots (3)$$

From (1) & (2)

$$\frac{a_1}{a_2} \neq \frac{b_1}{b_2} \neq \frac{c_1}{c_2} \text{ ie } \frac{4}{1} \neq \frac{-3}{3} \neq \frac{-1}{2}$$

From (2) & (3), $\frac{a_1}{a_2} \neq \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$ ie $1 \neq 3 \neq -2$

\therefore The system of equations has a unique solutions.

- 3. Vani, her father and her grand father have an average age of 53. One-half of her grand father's age plus one-third of her father's age plus one fourth of Vani's age is 65. Four years ago if Vani's grand-father was four times as old as Vani then how old are they all now ?**

Solution :

Let the present age of Vani, her father, grand father be x, y, z respectively.

By data given,

$$\frac{x+y+z}{3} = 53 \Rightarrow x + y + z = 159 \quad \dots\dots\dots (1)$$

$$\frac{1}{2}z + \frac{1}{3}y + \frac{1}{4}x = 65$$

$$\Rightarrow \frac{6z + 4y + 3x}{12} = 65 \Rightarrow$$

$$3x + 4y + 6z = 780 \quad \dots\dots\dots (2)$$

$$(z - 4) = 4(x - 4) \Rightarrow 4x - z = 12 \quad \dots\dots\dots (3)$$

Solving (1) & (2)

$$(1) \times (4) \Rightarrow 4x + 4y + 4z = 636$$

$$(2) \Rightarrow$$

$$\text{Subtracting} \quad \frac{4x + 4y + 4z = 636}{3x + 4y + 6z = 780} \quad \dots\dots\dots (4)$$

Solving (3) & (4)

$$(3) \times (2) \Rightarrow 8x - 2z = 24$$

$$(4) \Rightarrow 3x - 2z = -144$$

$$\text{Subtracting} \quad \frac{8x - 2z = 24}{3x - 2z = -144} \quad \dots\dots\dots$$

$$5x = 168$$

$$x = \frac{168}{7} = 24$$

Sub x = 24 in (3)

$$96 - z = 12$$

$$z = 84$$

$$\therefore (1) \Rightarrow 24 + y + 84 = 159$$

$$\Rightarrow y = 51$$

\therefore Vani's present age = 24 years

Father's present age = 51 years

Grand father's age = 84 years

4. The sum of the digits of a three-digit number is 11. If the digits are reversed, the new number is 46 more than five times the former number. If the hundreds digit plus twice the tens digit is equal to the units digit, then find the original three digit number ?

Solution :

Let x be the 100's digit

y be the 10's digit

z be the unit's digit of a 3 digit no.

\therefore The required number is $100x + 10y + z$
and the reversed number is $100z + 10y + x$

Given $\boxed{x + y + z = 11}$ (1)

\therefore By the data given,

$$100z + 10y + x = 5(100x + 10y + z) + 46$$

$$\Rightarrow \boxed{499x + 40y - 95z = -46}$$
 (2)

and

$$\text{also } x + 2y = z$$

$$\Rightarrow \boxed{x + 2y - z = 0}$$
 (3)

Adding (1) & (3), $2x + 3y = 11$ (4)

Solving (1) & (2)

$$(1) \times 95 \Rightarrow 95x + 95y + 95z = 1045$$

$$(2) \times 1 \Rightarrow 499x + 40y - 95z = -46$$

$$\text{Subtracting, } \underline{94x + 135y} = 999$$

$$\div \text{ by } 27 \Rightarrow 22x + 5y = 37$$
 (5)

Solving (4) & (5)

$$(4) \times 11 \Rightarrow 22x + 33y = 121$$

$$(5) \Rightarrow 22x + 5y = 37$$

$$\text{Subtracting, } \underline{28y = 84}$$

$$y = 3$$

Sub $y = 3$ in (4)

$$2x + 9 = 11$$

$$2x = 2$$

$$x = 1$$

Sub in (1)

$$x + y + z = 11$$

$$\Rightarrow 1 + 3 + z = 11$$

\therefore The 3 digit no is

$$100x + 10y + z$$

$$= 100(1) + 10(3) + 7$$

$$= 137 \text{ and reversed digit number } 731.$$

5. There are 12 pieces of five, ten and twenty rupee currencies whose total value is ₹ 105. When first 2 sorts are interchanged in their numbers its value will be increased by ₹ 20. Find the number of currencies in each sort.

Solution :

Let x be the number of 5 rupee currencies

Let y be the number of 10 rupee currencies

Let z be the number of 20 rupee currencies

By the data given,

$$x + y + z = 12$$
 (1)

$$5x + 10y + 20z = 105$$
 (2)

$$10x + 5y + 20z = 125$$
 (3)

Solving (1) & (2)

$$(1) \times 5 \Rightarrow 5x + 5y + 5z = 60$$

$$(2) \Rightarrow 5x + 10y + 20z = 105$$

$$\begin{array}{r} \text{Subtracting,} \\ \underline{-5y - 15z = -45} \\ \Rightarrow y + 3z = 9 \quad \dots\dots\dots (4) \end{array}$$

Solving (2) & (3)

$$(2) \times 2 \Rightarrow 10x + 20y + 40z = 210$$

$$(3) \Rightarrow 10x + 5y + 20z = 125$$

$$\begin{array}{r} \text{Subtracting,} \\ \underline{15y + 20z = 85} \\ \Rightarrow 3y + 4z = 17 \quad \dots\dots (5) \end{array}$$

$$(4) \times 3 \Rightarrow 3y + 9z = 27$$

$$\begin{array}{r} (5) \Rightarrow 3y + 4z = 17 \\ \underline{ + 5z = 10} \\ 5z = 10 \end{array}$$

Sub $z = 2$ in (4)

$$y + 6 = 9 \Rightarrow y = 3$$

$$\therefore (1) \Rightarrow x + 3 + 2 = 12 \Rightarrow x = 7$$

$$\therefore \text{No. of 5 rupee notes} = 7$$

$$\text{No. of 10 rupee notes} = 3$$

$$\text{No. of 20 rupee notes} = 2$$

II. GCD AND LCM OF POLYNOMIALS :

- ✓ **Step 1 :** First, divide $f(x)$ by $g(x)$ to obtain $f(x) = g(x)q(x) + r(x)$ where $q(x)$ is the quotient and $r(x)$ is the remainder. Then, $\deg[r(x)] < \deg[g(x)]$.
- ✓ **Step 2 :** If the remainder $r(x)$ is non-zero, divide $g(x)$ by $r(x)$ to obtain $g(x) = r(x)q_1(x) + r_1(x)$ where $r_1(x)$ is the new remainder. Then $\deg[r_1(x)] < \deg[r(x)]$. If the remainder $r_1(x)$ is zero, then $r(x)$ is the required GCD.
- ✓ **Step 3 :** If $r_1(x)$ is non-zero, then continue the process until we get zero as remainder. The divisor at this stage will be the required GCD.
- ✓ If the $f(x)$ and $g(x)$ are two polynomials of same degree then the polynomial carrying the highest coefficient will be the dividend.
- ✓ The Least Common Multiple of two or more algebraic expressions is the expression of lowest degree (or power) such that the expressions exactly divide it.

Example 3.10

Find the GCD of the polynomials $x^3 + x^2 - x + 2$ and $2x^3 - 5x^2 + 5x - 3$.

Solution :

Let $f(x) = 2x^3 - 5x^2 + 5x - 3$ and $g(x) = x^3 + x^2 - x + 2$

$$\begin{array}{r} 2 \\ \underline{2x^3 - 5x^2 + 5x - 3} \\ 2x^3 + 2x^2 - 2x + 4 \quad (-) \\ \underline{-7x^2 + 7x - 7} \\ -7(x^2 - x + 1) \end{array}$$

$-7(x^2 - x + 1) = 0$, note that -7 is not a divisor of $g(x)$

Now dividing $g(x) = x^3 + x^2 - x + 2$ by the new remainder $x^2 - x + 1$ (leaving the constant factor), we get

$$\begin{array}{r}
 x + 2 \\
 x^2 - x + 1 \overline{) x^3 + x^2 - x + 2} \\
 \underline{x^3 - x^2 + x} (-) \\
 2x^2 - 2x + 2 \\
 \underline{2x^2 - 2x + 2} (-) \\
 0
 \end{array}$$

Here, we get zero remainder

Therefore, $\text{GCD}(2x^3 - 5x^2 + 5x - 3, x^3 + x^2 - x + 2) = x^2 - x + 1$

Example 3.11

Find the GCD of $6x^3 - 30x^2 + 60x - 48$ and $3x^3 - 12x^2 + 21x - 18$

Solution :

Let $f(x) = 6x^3 - 30x^2 + 60x - 48 = 6$

$(x^3 - 5x^2 + 10x - 8)$ and

$g(x) = 3x^3 - 12x^2 + 21x - 18 = 3$

$(x^3 - 4x^2 + 7x - 6)$

Now, we shall find the GCD of $x^3 - 5x^2 + 10x - 8$ and $x^3 - 4x^2 + 7x - 6$

$$\begin{array}{r}
 1 \\
 x^3 - 5x^2 + 10x - 8 \overline{) x^3 - 4x^2 + 7x - 6} \\
 \underline{x^3 - 5x^2 + 10x - 8} (-) \\
 x^2 - 3x + 2
 \end{array}$$

$$\begin{array}{r}
 x - 2 \\
 x^2 - 3x + 2 \overline{) x^3 - 5x^2 + 10x - 8} \\
 \underline{x^3 - 3x^2 + 2x} (-) \\
 -2x^2 + 8x - 8 \\
 \underline{-2x^2 + 6x - 4} (-) \\
 2x - 4 \\
 = 2(x - 2)
 \end{array}$$

$$\begin{array}{r}
 x - 1 \\
 x - 2 \overline{) x^2 - 3x + 2} \\
 \underline{x^2 - 2x} (-) \\
 -x + 2 \\
 \underline{-x + 2} (-) \\
 0
 \end{array}$$

Here, we get zero as remainder.

GCD of leading coefficients 3 and 6 is 3.

Thus, $\text{GCD}[(6x^3 - 30x^2 + 60x - 48, 3x^3 - 12x^2 + 21x - 18)] = 3(x - 2)$

Example 3.12

Find the LCM of the following

$$\begin{array}{ll}
 \text{i) } 8x^4 y^2, 48x^2 y^4 & \text{ii) } 5x - 10, 5x^2 - 20 \\
 4 \quad 2 & 3 \quad 2 \quad 2
 \end{array}$$

Solution :

i) $8x^4 y^2, 48x^2 y^4$

First let us find the LCM of the numerical coefficients.

That is, LCM(8, 48)

$$= 2 \times 2 \times 2 \times 6 = 48$$

Then find the LCM of the terms involving variables.

That is, $\text{LCM}(x^4 y^2, x^2 y^4) = x^4 y^4$

Finally find the LCM of the given expression.

We conclude that the LCM of the given expression is the product of the LCM of the numerical coefficient and the LCM of the terms with variables.

$$\text{Therefore, } \text{LCM}(8x^4 y^2, 48x^2 y^4) = 48x^4 y^4$$

$$\begin{array}{r}
 2 \overline{) 8, 48} \\
 2 \overline{) 4, 24} \\
 2 \overline{) 2, 12} \\
 1, 6
 \end{array}$$

ii) $5x - 10, 5x^2 - 20$

$$5x - 10 = 5(x - 2)$$

$$5x^2 - 20 = 5(x^2 - 4) = 5(x + 2)(x - 2)$$

Therefore, LCM $[(5x - 10), (5x^2 - 20)]$

$$= 5(x + 2)(x - 2)$$

iii) $x^4 - 1, x^2 - 2x + 1$

$$x^4 - 1 = (x^2)^2 - 1 = (x^2 + 1)(x^2 - 1)$$

$$= (x^2 + 1)(x + 1)(x - 1)$$

$$x^2 - 2x + 1 = (x - 1)^2$$

Therefore, LCM $[(x^4 - 1), (x^2 - 2x + 1)]$

$$= (x^2 + 1)(x + 1)(x - 1)^2$$

iv) $x^3 - 27, (x - 3)^2, x^2 - 9$

$$x^3 - 27 = (x - 3)(x^2 + 3x + 9);$$

$$(x - 3) = (x - 3); (x - 9)$$

$$= (x + 3)(x - 3)$$

Therefore, LCM $[(x^3 - 27), (x - 3)^2, (x^2 - 9)] = (x - 3)^2(x + 3)(x^2 + 3x + 9)$

EXERCISE 3.2

1. Find the GCD of the given polynomials

i) $x^4 + 3x^3 - x - 3, x^3 + x^2 - 5x + 3$

ii) $x^4 - 1, x^3 - 11x^2 + x - 11$

iii) $3x^4 + 6x^3 - 12x^2 - 24x, 4x^4 + 14x^3 + 8x^2 - 8x$

iv) $3x^3 + 3x^2 + 3x + 3, 6x^3 + 12x^3 + 6x + 12$

Solution :

i) Let $f(x) = x^4 + 3x^3 - x - 3$

$$g(x) = x^3 + x^2 - 5x + 3$$

To find the GCD of $f(x), g(x)$

Divide $f(x)$ by $g(x)$

$$\begin{array}{r} x^3 + x^2 - 5x + 3 \overline{) x^4 + 3x^3 + 0x^2 - x - 3} \\ \underline{x^4 + x^3 - 5x^2 + 3x} \\ 2x^3 + 5x^2 - 10x + 6 \\ \underline{2x^3 + 2x^2 - 10x + 6} \\ 3x^2 + 6x - 9 \end{array}$$

$$= 3(x^2 + 2x - 3) \neq 0$$

Now, divide $g(x)$ by $x^2 + 2x - 3$ (excluding 3)

$$\begin{array}{r} x^2 + 2x - 3 \overline{) x^3 + x^2 - 5x + 3} \\ \underline{x^3 + 2x^2 - 3x} \\ -x^2 - 2x + 3 \\ \underline{-x^2 - 2x + 3} \\ 0 \end{array}$$

\therefore Remainder becomes 0.

\therefore The corresponding quotient is the HCF

$$\therefore \text{HCF} = x^2 + 2x - 3$$

ii) Let $f(x) = x^4 - 1$

$$g(x) = x^3 - 11x^2 + x - 11$$

$$\begin{array}{r} x^3 - 11x^2 + x - 11 \overline{) x^4 - 1} \\ \underline{x^4 - 11x^3 + x^2 - 11x} \\ 11x^3 - x^2 + 11x - 1 \\ \underline{11x^3 - 121x^2 + 11x - 121} \\ 120x^2 + 120 \\ = 120(x^2 + 1) \neq 0 \end{array}$$

Now, divide $g(x)$ by $x^2 + 1$

$$\begin{array}{r}
 x - 11 \\
 x^2 + 1 \overline{) x^3 - 11x^2 + x - 11} \\
 \underline{x^3 + 0x^2 + x} \\
 - 11x^2 - 11 \\
 \underline{- 11x^2 - 11} \\
 0
 \end{array}$$

$$\therefore \text{HCF} = x^2 + 1$$

iii) Let $f(x) = 4x^4 + 14x^3 + 8x^2 - 8x$
 $= 2x(2x^3 + 7x^2 + 4x - 4)$
 $g(x) = 3x^4 + 6x^3 - 12x^2 - 24x$
 $= 3x(1x^3 + 2x^2 - 4x - 8)$

$$\begin{array}{r}
 2 \\
 x^3 + 2x^2 - 4x - 8 \overline{) 2x^3 + 7x^2 + 4x - 4} \\
 \underline{2x^3 + 4x^2 - 8x - 16} \\
 3x^2 + 12x + 12 \\
 = 3(x^2 + 4x + 4) \neq 0
 \end{array}$$

$$\begin{array}{r}
 x - 2 \\
 x^2 + 4x + 4 \overline{) x^3 + 2x^2 - 4x - 8} \\
 \underline{x^3 + 4x^2 + 4x} \\
 - 2x^2 - 8x - 8 \\
 \underline{- 2x^2 - 8x - 8} \\
 0
 \end{array}$$

$$\therefore \text{G.C.D} = x(x^2 + 4x + 4)$$

iv) Let $f(x) = 6x^3 + 12x^2 + 6x + 12$
 $= 6(x^3 + 2x^2 + x + 2)$
 $g(x) = 3x^3 + 3x^2 + 3x + 3$
 $= 3(x^3 + x^2 + x + 1)$
 $\text{GCD of } 6, 3 = 3$

$$\begin{array}{r}
 1 \\
 x^3 + x^2 + x + 1 \overline{) x^3 + 2x^2 + x + 2} \\
 \underline{x^3 + x^2 + x + 1} \\
 x^2 + 1
 \end{array}$$

$$= x^2 + 1 \neq 0$$

$$\begin{array}{r}
 x + 1 \\
 x^2 + 1 \overline{) x^3 + x^2 + x + 1} \\
 \underline{x^3 + 0x^2 + x + 0} \\
 x^2 + 1 \\
 \underline{x^2 + 1} \\
 0
 \end{array}$$

$$\therefore \text{G.C.D} = 3(x^2 + 1)$$

2. Find the LCM of the given expressions.

$$x^2, x^3, x^2, x^3, x^2, x^2$$

iii) $16m, -12m^2n^2, 8n^2$

iv) $p^2 - 3p + 2, p^2 - 4$

v) $2x^2 - 5x - 3, 4x^2 - 36$

vi) $(2x^2 - 3xy)^2, (4x - 6y)^2, 8x^3 - 27y^3$

Solution :

i) $4x^2y = \underline{2} \times \underline{2} \times x^2 \times y$
 $8x^2y^2 = \underline{2} \times \underline{2} \times 2 \times x^2 \times y^2$
 $\therefore \text{LCM} = 2 \times 2 \times 2 \times x^2 \times y^2$
 $= 8x^2y^2$

ii) $-9a^3b^2 = -\underline{3} \times \underline{3} \times a^3 \times b^2$
 $12a^2b^2c = \underline{3} \times \underline{2} \times \underline{2} \times a^2 \times b^2 \times c$
 $\therefore \text{LCM} = -3 \times 3 \times 2 \times 2 \times a^3 \times b^2 \times c$
 $= -36a^3b^2c$

iii) $16m = \underline{2} \times \underline{2} \times (2) \times 2 \times m$
 $-12m^2n^2 = -\underline{3} \times \underline{2} \times \underline{2} \times m^2 \times n^2$
 $8n^2 = \underline{2} \times \underline{2} \times (2) \times n^2$
 $\therefore \text{LCM} = 2 \times 2 \times 2 \times 2 \times -3 \times m^2 \times n^2$
 $= -48m^2n^2$

$$\begin{aligned}\text{iv) } p^2 - 3p + 2 &= \underline{(p-2)} (p-1) \\ p^2 - 4 &= \underline{(p-2)} (p+2) \\ \therefore \text{LCM} &= (p-2) (p-1) (p+2)\end{aligned}$$

$$\begin{aligned}\text{v) } 2x^2 - 5x - 3 &= \underline{(x-3)} (2x+1) \\ 4x^2 - 36 &= (2x)^2 - (6)^2\end{aligned}$$

-5	-6
-6	+1
2	2

$$-3, \quad \frac{1}{2}$$

$$= (2x+6) (2x-6)$$

$$= 2 (x+3) 2 (x-3)$$

$$= 4 (x+3) (x-3)$$

$$\therefore \text{LCM} = 4(x-3) (x+3) (2x+1)$$

$$\begin{aligned}\text{vi) } (2x^2 - 3xy)^2 &= (x (2x - 3y))^2 \\ &= x^2 (2x - 3y)^2\end{aligned}$$

$$\begin{aligned}(4x - 6y)^3 &= (2 (2x - 3y))^3 \\ &= 8 (2x - 3y)^3\end{aligned}$$

$$\begin{aligned}8x^3 - 27y^3 &= (2x)^3 - (3y)^3 \\ &= (2x-3y) (4x^2 + 6xy + 9y^2)\end{aligned}$$

$$\begin{aligned}\therefore \text{LCM} &= (2x-3y)^3 8x^2 (4x^2 + 6xy + 9y^2) \\ &= 8x^2 (2x-3y)^3 (4x^2 + 6xy + 9y^2)\end{aligned}$$

Key Points

- ✓ The product of two polynomials is the product of their LCM and GCD. That is, $f(x) \times g(x) = \text{LCM}[f(x), g(x)] \times \text{GCD}[f(x), g(x)]$.

EXERCISE 3.3

1. Find the LCM and GCD for the following and verify that $f(x) \times g(x) = \text{LCM} \times \text{GCD}$
- i) $21x^2y, 35xy^2$ ii) $(x^3-1)(x+1), (x^3+1)$ iii) $(x^2y+xy^2), (x^2+xy)$

Solution :

$$\begin{aligned}\text{i) Given } f(x) &= 21x^2y = 7 \times 3 \times x^2 \times y \\ g(x) &= 35xy^2 = 7 \times 5 \times x \times y^2 \\ \therefore \text{GCD} &= 7 \times x \times y = 7xy \\ \therefore \text{LCM} &= 7 \times x^2 \times y^2 \times 15 = 105x^2y^2 \\ \therefore f(x) \times g(x) &= 21x^2y \times 35xy^2 \\ &= 735x^3y^3 \\ \therefore \text{LCM} \times \text{GCD} &= 105x^2y^2 \times 7xy \\ &= 735x^3y^3 \\ \therefore f(x) \times g(x) &= \text{LCM} \times \text{GCD}\end{aligned}$$

$$\begin{aligned}\text{ii) Given } f(x) &= (x^3-1)(x+1) \\ &= (x-1)(x^2+x+1) \underline{(x+1)} \\ g(x) &= x^3+1 \\ &= \underline{(x+1)} (x^2-x+1) \\ \therefore \text{GCD} &= x+1 \\ \therefore \text{LCM} &= (x+1)(x-1)(x^2+x+1)(x^2-x+1) \\ &= (x^3+1)(x^3-1) \\ &= x^6-1 \\ \therefore f(x) \times g(x) &= (x-1)(x^2+x+1)(x+1)(x+1) \\ &\quad (x^2-x+1) \\ &= (x^3-1)(x+1)(x^3+1) \\ &= (x+1)(x^6-1) \\ \therefore \text{LCM} \times \text{GCD} &= (x^6-1)(x+1) \\ &= 735x^3y^3 \\ \therefore f(x) \times g(x) &= \text{LCM} \times \text{GCD}\end{aligned}$$

$$\begin{aligned}
 \text{iii) } f(x) &= x^2y + xy^2 \\
 &= xy(x + y) \\
 g(x) &= x^2 + xy \\
 &= x(x + y) \\
 \therefore \text{GCD} &= x(x + y) \\
 \therefore \text{LCM} &= x(x + y)y \\
 &= (x^3 + 1)(x^3 - 1) \\
 &= x^6 - 1 \\
 \therefore f(x) \times g(x) &= xy(x + y) \times x(x + y) \\
 &= x^2y(x + y)^2 \\
 \therefore \text{LCM} \times \text{GCD} &= xy(x + y)x(x + y) \\
 &= x^2y(x + y)^2 \\
 \therefore f(x) \times g(x) &= \text{LCM} \times \text{GCD}
 \end{aligned}$$

2. Find the LCM of each pair of the following polynomials

$$a - 2$$

ii) $x^4 - 27a^3x$, $(x - 3a)^2$ whose GCD is $(x - 3a)$

Solution :

$$\begin{aligned}
 \text{i) Let } f(x) &= a^2 + 4a - 12 \\
 &= (a + 6)(a - 2) \\
 g(x) &= a^2 - 5a + 6 \\
 &= (a - 3)(a - 2) \\
 \text{GCD} &= a - 2 \\
 \therefore \text{LCM} &= \frac{f(x) \times g(x)}{\text{GCD}} \\
 &= \frac{(a + 6)(a - 2) \times (a - 3)(a - 2)}{a - 2} \\
 &= (a + 6)(a - 3)(a - 2)
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) Let } f(x) &= x^4 - 27a^3x \\
 &= x(x^3 - 27a^3) \\
 &= x(x^3 - (3a)^3) \\
 &= x[(x - 3a)(x^2 + 3ax + 9a^2)]
 \end{aligned}$$

$$\begin{aligned}
 g(x) &= (x - 3a)^2 \\
 &= (x - 3a)(x - 3a) \\
 \text{GCD} &= x - 3a \\
 \therefore \text{LCM} &= \frac{f(x) \times g(x)}{\text{GCD}} \\
 &= \frac{x(x - 3a)(x^2 + 3ax + 9a^2)(x - 3a)^2}{x - 3a} \\
 &= x(x^2 + 3ax + 9a^2)(x - 3a)^2
 \end{aligned}$$

3. Find the GCD of each pair of the following polynomials

i) $12(x^4 - x^3)$, $8(x^4 - 3x^3 + 2x^2)$ whose LCM is $24x^3(x - 1)(x - 2)$

ii) $(x^3 + y^3)$, $(x^4 + x^2y^2 + y^4)$ whose LCM is $(x^3 + y^3)(x^2 + xy + y^2)$

Solution :

$$\begin{aligned}
 \text{i) } f(x) &= 12(x^4 - x^3) \\
 &= 12x^3(x - 1) \\
 g(x) &= 8(x^4 - 3x^3 + 2x^2) \\
 &= 8 \times x^2(x^2 - 3x + 2) \\
 &= 8 \times x^2(x - 2)(x - 1) \\
 \text{LCM} &= 24x^3(x - 1)(x - 2)
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{GCD} &= \frac{f(x) \times g(x)}{\text{LCM}} \\
 &= \frac{12x^3(x - 1)8x^2(x - 2)(x - 1)}{24x^3(x - 1)(x - 2)} \\
 &= 4x^2(x - 1)
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) } f(x) &= (x^3 + y^3) = (x + y)(x^2 - xy + y^2) \\
 g(x) &= x^4 + x^2y^2 + y^4 = (x^2 - xy + y^2)(x^2 + xy + y^2) \\
 \text{LCM} &= (x^3 + y^3)(x^2 + xy + y^2) \\
 &= (x + y)(x^2 - xy + y^2)(x^2 + xy + y^2) \\
 \therefore \text{GCD} &= \frac{f(x) \times g(x)}{\text{LCM}} \\
 &= \frac{(x + y)(x^2 - xy + y^2) \times (x^2 - xy + y^2)(x^2 + xy + y^2)}{(x + y)(x^2 - xy + y^2)(x^2 + xy + y^2)} \\
 &= x^2 - xy + y^2
 \end{aligned}$$

4. Given the LCM and GCD of the two polynomials $p(x)$ and $q(x)$ find the unknown polynomial in the following table

S. No.	LCM	GCD
i)	$a^3 - 10a^2 + 11a + 70$	$a - 7$
ii)	$(x^2 + y^2)(x^4 + x^2y^2 + y^4)$	$(x^2 - y^2)$

$p(x)$	$q(x)$
$a^2 - 12a + 35$	
	$(x^4 - y^4)(x^2 + y^2 - xy)$

Solution :

i) Given $LCM = a^3 - 10a^2 + 11a + 70$,
 $GCD = a - 7$
 $p(x) = a^2 - 12a + 35$, $q(x) = ?$

$$q(x) = \frac{LCM \times GCD}{p(x)} = \frac{(a^3 - 10a^2 + 11a + 70)(a - 7)}{(a - 7)(a - 5)}$$

$$= \frac{(a - 5)(a^5 - 5a - 14)(a - 7)}{(a - 7)(a - 5)}$$

$$= a^2 - 5a - 14$$

$$q(x) = (a - 7)(a + 2)$$

ii) Given $LCM = (x^2 + y^2)(x^4 + x^2y^2 + y^4)$

$$GCD = (x^2 - y^2)$$

$$q(x) = (x^4 - y^4)(x^2 + y^2 - xy) \quad p(x) = ?$$

$$p(x) = \frac{LCM \times GCD}{q(x)}$$

$$= \frac{(x^2 + y^2)(x^4 + x^2y^2 + y^4)(x^2 - y^2)}{(x^2 + y^2)(x^2 - y^2)(x^2 + y^2 - xy)}$$

$$= \frac{(x^2 + y^2 - xy)(x^2 - xy + y^2)}{x^2 + y^2 - xy}$$

$$= x^2 + xy + y^2$$

Key Points

- ✓ An expression is called a rational expression if it can be written in the form $\frac{p(x)}{q(x)}$ where $p(x)$ and $q(x)$ are polynomials and $q(x) \neq 0$.
- ✓ A rational expression is the ratio of two polynomials.
- ✓ A rational expression $\frac{p(x)}{q(x)}$ is said to be in its lowest form if $GCD(p(x), q(x)) = 1$.
- ✓ A value that makes a rational expression (in its lowest form) undefined is called an excluded value.

Example 3.13

Reduce the rational expressions to its lowest form

(i) $\frac{x-3}{x^2-9}$ (ii) $\frac{x^2-16}{x^2+8x+16}$

Solution :

$$(i) \frac{x-3}{x^2-9} = \frac{x-3}{(x+3)(x-3)} = \frac{1}{x+3}$$

$$(ii) \frac{x^2-16}{x^2+8x+16} = \frac{(x+4)(x-4)}{(x+4)^2} = \frac{x-4}{x+4}$$

Example 3.14

Find the excluded values of the following expressions (if any).

i) $\frac{x+10}{8x}$ (ii) $\frac{7p+2}{8p^2+13p+5}$ (iii) $\frac{x}{x^2+1}$

Solution :

i) $\frac{x+10}{8x}$

The expression $\frac{x+10}{8x}$ is undefined when

$8x = 0$ or $x = 0$. Hence the excluded value is 0.

$$(ii) \frac{7p+2}{8p^2+13p+5}$$

The expression $\frac{7p+2}{8p^2+13p+5}$ is undefined when

$$8p^2 + 13p + 5 = 0 \text{ that is}$$

$$(8p + 5)(p + 1) = 0$$

$$p = \frac{-5}{8}, p = -1.$$

The excluded values are $\frac{-5}{8}$ and -1 .

$$(iii) \frac{x}{x^2+1}$$

Here $x^2 \geq 0$ for all x .

Therefore, $x^2 + 1 \geq 0 + 1 = 1$. Hence, $x^2 + 1 \neq 0$ for any x .

Therefore, there can be no real excluded values for the given rational expression $\frac{x}{x^2+1}$

EXERCISE 3.4

1. Reduce each of the following rational expressions to its lowest form.

$$i) \frac{x^2-1}{x^2+x} \quad ii) \frac{x^2-11x+18}{x^2-4x+4}$$

$$iii) \frac{9x^2+81x}{x^3+8x^2-9x} \quad iv) \frac{p^2-3p-40}{2p^3-24p^2+64p}$$

Solution :

$$\begin{aligned} i) \frac{x^2-1}{x^2+x} &= \frac{(x+1)(x-1)}{x(x+1)} \\ &= \frac{x-1}{x} \end{aligned}$$

$$\begin{aligned} ii) \frac{x^2-11x+18}{x^2-4x+4} &= \frac{(x-9)(x-2)}{(x-2)(x-2)} \\ &= \frac{x-9}{x-2} \end{aligned}$$

$$\begin{aligned} iii) \frac{9x^2+81x}{x^3+8x^2-9x} &= \frac{9x(x+9)}{x(x^2+8x-9)} \\ &= \frac{9(x+9)}{(x+9)(x-1)} \\ &= \frac{9}{x-1} \end{aligned}$$

$$\begin{aligned} iv) \frac{p^2-3p-40}{\frac{3}{2}} &= \frac{p^2-3p-40}{2p(p^2-12p+32)} \\ &= \frac{(p-8)(p+5)}{2p(p-8)(p+4)} \\ &= \frac{p+5}{2p(p-4)} \end{aligned}$$

2. Find the excluded values, if any of the following expressions.

$$i) \frac{y}{y^2-25} \quad ii) \frac{t}{t^2-5t+6}$$

$$iii) \frac{x^2+6x+8}{x^2+x-2} \quad iv) \frac{x^3-27}{x^3+x^2-6x}$$

Solution :

$$i) \frac{y}{y^2-25} = \frac{y}{(y+5)(y-5)}$$

The expression is undefined if $y = -5$, $y = 5$

\therefore The excluded values are $5, -5$

$$\begin{aligned} \text{ii) } \frac{t}{t^2 - 5t + 6} \\ = \frac{t}{(t-3)(t-2)} \end{aligned}$$

The expression is undefined if $t = 3, t = 2$

\therefore The excluded values are 3, 2

$$\begin{aligned} \text{iii) } \frac{x^2 + 6x + 8}{x^2 + x - 2} \\ = \frac{(x+4)(x+2)}{(x+2)(x-1)} \\ = \frac{x+4}{x-1} \end{aligned}$$

This expression is not defined if $x = 1$

$$\begin{aligned} \text{iv) } \frac{x^3 - 27}{x^3 + x^2 - 6x} \\ = \frac{x^3 - 3^3}{x(x^2 + x - 6)} \\ = \frac{(x-3)(x^2 + 3x + 9)}{x(x+3)(x-2)} \end{aligned}$$

This expression is not defined for

$$x = 0, x = -3, x = 2$$

\therefore The excluded values are 0, -3, 2

V. MULTIPLICATION AND DIVISION OF RATIONAL EXPRESSIONS:

Key Points

✓ If $\frac{p(x)}{q(x)}$ and $\frac{r(x)}{s(x)}$ are two rational expressions where $q(x) \neq 0, s(x) \neq 0$, their product is

$$\frac{p(x)}{q(x)} \times \frac{r(x)}{s(x)} = \frac{p(x) \times r(x)}{q(x) \times s(x)}$$

✓ If $\frac{p(x)}{q(x)}$ and $\frac{r(x)}{s(x)}$ are two rational expressions where $q(x), s(x) \neq 0$, then,

$$\frac{p(x)}{q(x)} \div \frac{r(x)}{s(x)} = \frac{p(x)}{q(x)} \times \frac{s(x)}{r(x)} = \frac{p(x) \times s(x)}{q(x) \times r(x)}$$

Example 3.15

i) Multiply $\frac{x^3}{9y^2}$ by $\frac{27y}{x^5}$ ii) Multiply $\frac{x^4b^2}{x-1}$ by $\frac{x^2-1}{a^4b^3}$

Solution :

$$\text{i) } \frac{x^3}{9y^2} \times \frac{27y}{x^5} = \frac{3}{x^2y}$$

$$\text{ii) } \frac{x^4b^2}{x-1} \times \frac{x^2-1}{a^4b^3} = \frac{x^4 \times b^2}{x-1} \times \frac{(x+1)(x-1)}{a^4 \times b^3} = \frac{x^4(x+1)}{a^4b}$$

Example 3.16

Find

$$i) \frac{14x^4}{y} \div \frac{7x}{3y^4} \quad ii) \frac{x^2 - 16}{x + 4} \div \frac{x - 4}{x + 4}$$

$$iii) \frac{16x^2 - 2x - 3}{3x^2 - 2x - 1} \div \frac{8x^2 + 11x + 3}{3x^2 - 11x - 4}$$

Solution :

$$i) \frac{14x^4}{y} \div \frac{7x}{3y^4} = \frac{14x^4}{y} \times \frac{3y^4}{7x} = 6x^3y^3$$

$$ii) \frac{x^2 - 16}{x + 4} \div \frac{x - 4}{x + 4} = \frac{(x + 4)(x - 4)}{(x + 4)} \times \left(\frac{x + 4}{x - 4} \right) = x + 4$$

$$iii) \frac{16x^2 - 2x - 3}{3x^2 - 2x - 1} \div \frac{8x^2 + 11x + 3}{3x^2 - 11x - 4}$$

$$= \frac{16x^2 - 2x - 3}{3x^2 - 2x - 1} \times \frac{3x^2 - 11x - 4}{8x^2 + 11x + 3}$$

$$= \frac{(8x + 3)(2x - 1)}{(x - 1)(x + 1)} \times \frac{(3x + 1)(x - 4)}{x^2 - 1}$$

$$= \frac{(2x - 1)(x - 4)}{(x - 1)(x + 1)} = \frac{2x^2 - 9x + 4}{x^2 - 1}$$

EXERCISE 3.5**1. Simplify**

$$i) \frac{4x^2y}{2z^2} \times \frac{6xz^3}{20y^4} \quad ii) \frac{p^2 - 10p + 21}{p - 7} \times \frac{p^2 + p - 12}{(p - 3)^2}$$

$$iii) \frac{5t^3}{4t - 8} \times \frac{6t - 12}{10t}$$

Solution :

$$i) \frac{4x^2y}{2z^2} \times \frac{6xz^3}{20y^4}$$

$$= \frac{\cancel{4}^2 x^2 \cancel{y}}{\cancel{2}^1 z^2} \times \frac{\cancel{6}^3 xz^{\cancel{3}^1}}{\cancel{20}^4 y^{\cancel{4}^1}}$$

$$= \frac{3}{5} \frac{x^3z}{y^3}$$

$$ii) \frac{p^2 - 10p + 21}{p - 7} \times \frac{p^2 + p - 12}{(p - 3)^2}$$

$$= \frac{(\cancel{p-7})(\cancel{p-3})}{\cancel{p-7}} \times \frac{(p+4)(\cancel{p-3})}{(\cancel{p-3})^2}$$

$$= p + 4$$

$$iii) \frac{5t^3}{4t - 8} \times \frac{6t - 12}{10t}$$

$$= \frac{5t^3}{4(t - 2)} \times \frac{6(t - 2)}{10t}$$

$$= \frac{3t^2}{4}$$

2. Simplify

$$i) \frac{x + 4}{3x + 4y} \times \frac{9x^2 - 16y^2}{2x^2 + 3x - 20}$$

$$ii) \frac{x^3 - y^3}{3x^2 + 9xy + 6y^2} \times \frac{x^2 + 2xy + y^2}{x^2 - y^2}$$

Solution :

$$i) \frac{x + 4}{3x + 4y} \times \frac{9x^2 - 16y^2}{2x^2 + 3x - 20}$$

$$= \frac{x + 4}{3x + 4y} \times \frac{(3x + 4y)(3x - 4y)}{(x + 4)(2x - 5)}$$

$$= \frac{3x - 4y}{2x - 5}$$

3	-40
8	-5
2	2
4,	-5/2

$$\begin{aligned}
 \text{ii) } & \frac{x^3 - y^3}{3x^2 + 9xy + 6y^2} \times \frac{x^2 + 2xy + y^2}{x^2 - y^2} \\
 &= \frac{\cancel{(x-y)} (x^2 + xy + y^2)}{3(x^2 + 3xy + 2y^2)} \times \frac{\cancel{(x+y)} \cancel{(x+y)}}{\cancel{(x+y)} \cancel{(x-y)}} \\
 &= \frac{(x^2 + xy + y^2) \cancel{(x+y)}}{3(x+2y) \cancel{(x+y)}} \\
 &= \frac{x^2 + xy + y^2}{3(x+2y)}
 \end{aligned}$$

3. Simplify

$$\text{i) } \frac{2a^2 + 5a + 3}{2a^2 + 7a + 6} \div \frac{a^2 + 6a + 5}{-5a^2 - 35a - 50}$$

$$\text{ii) } \frac{b^2 + 3b - 28}{b^2 + 4b + 4} \div \frac{b^2 - 49}{b^2 - 5b - 14}$$

$$\text{iii) } \frac{x+2}{4y} \div \frac{x^2 - x - 6}{12y^2}$$

$$\text{iv) } \frac{12t^2 - 22t + 8}{3t} \div \frac{3t^2 + 2t - 8}{2t^2 + 4t}$$

Solution :

$$\begin{aligned}
 \text{i) } & \frac{2a^2 + 5a + 3}{2a^2 + 7a + 6} \div \frac{a^2 + 6a + 5}{-5a^2 - 35a - 50} \\
 &= \frac{(2a+3)(a+1)}{(2a+3)(a+2)} \times \frac{-5(a^2 + 7a + 10)}{(a+5)(a+1)} \\
 &= \frac{\cancel{(2a+3)} \cancel{(a+1)}}{\cancel{(2a+3)} \cancel{(a+2)}} \times \frac{-5\cancel{(a+5)} \cancel{(a+2)}}{\cancel{(a+5)} \cancel{(a+1)}} \\
 &= -5
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) } & \frac{b^2 + 3b - 28}{b^2 + 4b + 4} \div \frac{b^2 - 49}{b^2 - 5b - 14} \\
 &= \frac{(b+7)(b-4)}{(b+2)(b+2)} \times \frac{(b-7)(b+2)}{(b+7)(b-7)} \\
 &= \frac{b-4}{b+2}
 \end{aligned}$$

$$\begin{aligned}
 \text{iii) } & \frac{x+2}{4y} \div \frac{x^2 - x - 6}{12y^2} \\
 &= \frac{x+2}{4y} \times \frac{12y^2}{(x-3)(x+2)} \\
 &= \frac{3y}{x-3}
 \end{aligned}$$

$$\begin{aligned}
 \text{iv) } & \frac{12t^2 - 22t + 8}{3t} \div \frac{3t^2 + 2t - 8}{2t^2 + 4t} \\
 &= \frac{2(6t^2 - 11t + 4)}{3t} \times \frac{2t(t+2)}{3t^2 + 2t - 8} \\
 &= \frac{2(3t-4)(2t-1)}{3t} \times \frac{2t(t+2)}{(3t-4)(t+3)} \\
 &= \frac{4(2t-1)}{3}
 \end{aligned}$$

4. If $x = \frac{a^2 + 3a - 4}{3a - 3}$ and $y = \frac{a^2 + 2a - 8}{2a - 2a - 4}$
find the value of $x^2 y^{-2}$.

Solution :

Given

$$\begin{aligned}
 x &= \frac{a^2 + 3a - 4}{3a^2 - 3}, \quad y = \frac{a^2 + 2a - 8}{2a^2 - 2a - 4} \\
 &= \frac{(a+4)(a-1)}{3(a+1)(a-1)} = \frac{(a+4)(a-2)}{2(a-2)(a+1)} \\
 &= \frac{a+4}{3(a+1)} = \frac{a+4}{2(a+1)}
 \end{aligned}$$

$$\begin{aligned}
 \therefore x^2 y^{-2} &= \frac{x^2}{y^2} \\
 &= \frac{(a+4)^2}{9(a+1)^2} \times \frac{4(a+1)^2}{(a+4)^2} \\
 &= \frac{4}{9}
 \end{aligned}$$

5. If a polynomial $p(x) = x^2 - 5x - 14$ is divided by another polynomial $q(x)$ we get

$$\frac{x-7}{x+2} \text{ find } q(x).$$

Solution :

Given

$$\frac{p(x)}{q(x)} = \frac{x-7}{x+2}$$

$$\Rightarrow \frac{x^2 - 5x - 14}{q(x)} = \frac{x-7}{x+2}$$

$$\Rightarrow \frac{(x-7)(x+2)}{q(x)} = \frac{x-7}{x+2}$$

$$\Rightarrow q(x) = (x+2)^2$$

$$\Rightarrow q(x) = x^2 + 4x + 4 \text{ is another polynomial}$$

VI. ADDITION AND SUBTRACTION OF RATIONAL EXPRESSION:

Example 3.17

Find $\frac{x^2 + 20x + 36}{x^2 - 3x - 28} - \frac{x^2 + 12x + 4}{x^2 - 3x - 28}$

Solution :

$$\begin{aligned} &= \frac{x^2 + 20x + 36}{x^2 - 3x - 28} - \frac{x^2 + 12x + 4}{x^2 - 3x - 28} \\ &= \frac{(x^2 + 20x + 36) - (x^2 + 12x + 4)}{x^2 - 3x - 28} \\ &= \frac{8x + 32}{x^2 - 3x - 28} = \frac{8(x+4)}{(x-7)(x+4)} = \frac{8}{x-7} \end{aligned}$$

Example 3.18

Simplify $\frac{1}{x^2 - 5x + 6} + \frac{1}{x^2 - 3x + 2} - \frac{1}{x^2 - 8x + 15}$

Solution :

$$\begin{aligned} &= \frac{1}{x^2 - 5x + 6} + \frac{1}{x^2 - 3x + 2} - \frac{1}{x^2 - 8x + 15} \\ &= \frac{1}{(x-2)(x-3)} + \frac{1}{(x-2)(x-1)} - \frac{1}{(x-5)(x-3)} \\ &= \frac{(x-1)(x-5) + (x-3)(x-5) - (x-1)(x-2)}{(x-1)(x-2)(x-3)(x-5)} \\ &= \frac{(x^2 - 6x + 5) + (x^2 - 8x + 15) - (x^2 - 3x + 2)}{(x-1)(x-2)(x-3)(x-5)} \\ &= \frac{x^2 - 11x + 8}{(x-1)(x-2)(x-3)(x-5)} \\ &= \frac{(x-9)(x-2)}{(x-1)(x-2)(x-3)(x-5)} \\ &= \frac{x-9}{(x-1)(x-3)(x-5)} \end{aligned}$$

EXERCISE 3.6

1. Simplify

i) $\frac{x(x+1)}{x-2} + \frac{x(1-x)}{x-2}$ ii) $\frac{x+2}{x+3} + \frac{x-1}{x-2}$

iii) $\frac{x^3}{x-y} + \frac{y^3}{y-x}$

$$\begin{aligned} \text{i) } &\frac{x(x+1)}{x-2} + \frac{x(1-x)}{x-2} \\ &= \frac{x^2 + x + x - x^2}{x-2} \\ &= \frac{2x}{x-2} \end{aligned}$$

$$\begin{aligned} \text{ii) } &\frac{x+2}{x+3} + \frac{x-1}{x-2} \\ &= \frac{(x-2)(x+2) + (x+3)(x-1)}{(x+3)(x-2)} \\ &= \frac{(x^2 - 4) + (x^2 + 2x - 3)}{(x+3)(x-2)} \\ &= \frac{2x^2 + 2x - 7}{(x+3)(x-2)} \end{aligned}$$

$$\begin{aligned}
 \text{iii) } & \frac{x^3}{x-y} + \frac{y^3}{y-x} \\
 &= \frac{x^3}{x-y} - \frac{y^3}{x-y} \\
 &= \frac{x^3 - y^3}{x-y} \\
 &= \frac{(x-y)(x^2 + xy + y^2)}{x-y} \\
 &= x^2 + xy + y^2
 \end{aligned}$$

2. Simplify

$$\text{i) } \frac{(2x+1)(x-2)}{x-4} - \frac{(2x^2-5x+2)}{x-4}$$

$$\text{ii) } \frac{4x}{x^2-1} - \frac{x+1}{x-1}$$

$$\begin{aligned}
 \text{i) } & \frac{(2x+1)(x-2)}{x-4} - \frac{(2x^2-5x+2)}{x-4} \\
 &= \frac{2x^2 - 3x + 2 - 2x^2 + 5x - 2}{x-4} \\
 &= \frac{2x-4}{x-4} = \frac{2(x-2)}{x-4}
 \end{aligned}$$

$$\begin{aligned}
 \text{ii) } & \frac{4x}{x^2-1} - \frac{x+1}{x-1} \\
 &= \frac{4x}{(x+1)(x-1)} - \frac{x+1}{x-1} \\
 &= \frac{4x - (x+1)^2}{(x+1)(x-1)} \\
 &= \frac{4x - (x^2 + 2x + 1)}{(x+1)(x-1)} \\
 &= \frac{-x^2 + 2x - 1}{(x+1)(x-1)}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{-(x^2 - 2x + 1)}{(x+1)(x-1)} \\
 &= \frac{-(x-1)(x-1)}{(x+1)(x-1)} \\
 &= \frac{1-x}{1+x}
 \end{aligned}$$

3. Subtract $\frac{1}{x^2+2}$ **from** $\frac{2x^3+x^2+3}{(x^2+2)^2}$

Solution :

$$\begin{aligned}
 &= \frac{2x^3+x^2+3}{(x^2+2)^2} - \frac{1}{x^2+2} \\
 &= \frac{(2x^3+x^2+3) - (x^2+2)}{(x^2+2)^2} \\
 &= \frac{2x^3+x^2+3-x^2-2}{(x^2+2)^2} \\
 &= \frac{2x^3+1}{(x^2+2)^2}
 \end{aligned}$$

4. Which rational expression should be subtracted from $\frac{x^2+6x+8}{x^3+8}$ **to get** $\frac{3}{x^2-2x+4}$.

Solution :

$$\begin{aligned}
 & \frac{x^2+6x+8}{x^3+8} - \frac{3}{x^2-2x+4} \\
 &= \frac{(x+4)\cancel{(x+2)}}{\cancel{(x+2)}(x^2-2x+4)} - \frac{3}{x^2-2x+4} \\
 &= \frac{x+4-3}{x^2-2x+4} = \frac{x+1}{x^2-2x+4}
 \end{aligned}$$

5. If $A = \frac{2x+1}{2x-1}$, $B = \frac{2x-1}{2x+1}$ find

$$\frac{1}{A-B} - \frac{2B}{A^2-B^2}$$

Solution :

$$\begin{aligned} &= \frac{1}{A-B} - \frac{2B}{A^2-B^2} \\ &= \frac{1}{A-B} - \frac{2B}{(A+B)(A-B)} \\ &= \frac{A+B-2B}{(A+B)(A-B)} \\ &= \frac{A-B}{(A+B)(A-B)} \\ &= \frac{1}{A+B} \\ &= \frac{1}{\frac{2x+1}{2x-1} + \frac{2x-1}{2x+1}} \\ &= \frac{1}{\frac{(2x+1)^2 + (2x-1)^2}{4x^2-1}} \\ &= \frac{4x^2-1}{2[(2x+1)^2 + (2x-1)^2]} \quad (\because (a+b)^2 + (a-b)^2 = 2(a^2+b^2)) \\ &= \frac{4x^2-1}{2[4x^2+1]} \end{aligned}$$

6. If $A = \frac{x}{x+1}$, $B = \frac{1}{x+1}$ prove that

$$\frac{(A+B)^2 + (A-B)^2}{A \div B} = \frac{2(x^2+1)}{x(x+1)^2}$$

Solution :

Given

$$\begin{aligned} A &= \frac{x}{x+1}, B = \frac{1}{x+1} \\ &= \frac{(A+B)^2 + (A-B)^2}{\frac{A}{B}} \\ &= 2(A^2 + B^2) \times \frac{B}{A} \end{aligned}$$

$$= 2 \left[\frac{x^2}{(x+1)^2} + \frac{1}{(x+1)^2} \right] \times \frac{\frac{1}{x+1}}{\frac{x}{x+1}}$$

$$= \frac{2(1+x^2)}{(x+1)^2} \times \frac{1}{x}$$

$$= \frac{2(x^2+1)}{x(x+1)^2}$$

$$= \text{RHS}$$

Hence Proved

7. Pari needs 4 hours to complete a work. His friend Yuvan needs 6 hours to complete the same work. How long will it take to complete if they work together?

Solution :

Time taken by Pari to Complete a work
= 4 hrs.

\therefore In 1 hour, he completes $\frac{1}{4}$ part of work

Time taken by Yuvan to complete the same work = 6 hrs.

\therefore In 1 hour, he completes $\frac{1}{6}$ part of work

\therefore Total work completion by both in 1 hr

$$= \frac{1}{4} + \frac{1}{6}$$

$$= \frac{3+2}{12}$$

$$= \frac{5}{12} \text{ part of work}$$

\therefore Time taken by both together to complete the work = $\frac{12}{5}$ hrs

$$= 2 \frac{2}{5} \text{ hrs}$$

$$= 2 \text{ hrs } 24 \text{ min.}$$

8. Iniya bought 50 kg of fruits consisting of apples and bananas. She paid twice as much per kg for the apple as she did for the banana. If Iniya bought ₹ 1800 worth of apples and ₹ 600 worth bananas, then how many kgs of each fruit did she buy?

Solution :

Let x be the weight and p be the price of an apple.

Let y be the weight and q be the price of a banana

$$\therefore \text{By data given, } x + y = 50 \quad \text{--- (1)}$$

$$px = 1800 \quad \text{--- (2)}$$

$$qy = 600 \quad \text{--- (3)}$$

Also, $p = 2q$.

$$\therefore (2) \Rightarrow 2qx = 1800$$

$$\Rightarrow q = \frac{900}{x}$$

$$(3) \Rightarrow \frac{900}{x} y = 600$$

$$\Rightarrow y = \frac{2x}{3}$$

$$(1) \Rightarrow x + \frac{2x}{3} = 50 \Rightarrow 5x = 150$$

$$\Rightarrow x = 30$$

$$y = 50 - 30 = 20.$$

\therefore She bought 30 kg of apple & 20 kg of banana.

VII. SQUARE ROOT OF POLYNOMIALS BY FACTORISATION :

- ✓ The square root of a given positive real number is another number which when multiplied with itself is the given number
- ✓ $|q(x)| = \sqrt{p(x)}$ where $|q(x)|$ is the absolute value of $q(x)$.

Example 3.19

Find the square root of the following expressions

$$(i) 256(x-a)^8(x-b)^4(x-c)(x-d)^{20} \quad (ii) \frac{144 a^8 b^{12} c^{16}}{81 f^{12} g^4 h^{14}}$$

Solution :

$$i) \sqrt{256(x-a)^8(x-b)^4(x-c)^{16}(x-d)^{20}} = 16 \left| (x-a)^4(x-b)^2(x-c)^8(x-d)^{10} \right|$$

$$ii) \sqrt{\frac{144 a^8 b^{12} c^{16}}{81 f^{12} g^4 h^{14}}} = \frac{4}{3} \left| \frac{a^4 b^6 c^8}{f^6 g^2 h^7} \right|$$

Example 3.20

Find the square root of the following expressions

$$(i) 16x^2 + 9y^2 - 24xy + 24x - 18y + 9 \quad (ii) (6x^2 + x - 1)(3x^2 + 2x - 1)(2x^2 + 3x + 1)$$

$$iii) \left[\sqrt{15}x^2 + (\sqrt{3} + \sqrt{10})x + \sqrt{2} \right] \left[\sqrt{5}x^2 + (2\sqrt{5} + 1)x + 2 \right] \left[\sqrt{3}x^2 + (\sqrt{2} + 2\sqrt{3})x + 2\sqrt{2} \right]$$

Solution :

$$\begin{aligned}
 i) \quad & \sqrt{16x^2 + 9y^2 - 24xy + 24x - 18y + 9} \\
 &= \sqrt{(4x)^2 + (-3y)^2 + (3)^2 + 2(4x)(-3y) + 2(-3y)(3) + 2(4x)(3)} \\
 &= \sqrt{(4x - 3y + 3)^2} = |4x - 3y + 3|
 \end{aligned}$$

$$\begin{aligned}
 ii) \quad & \sqrt{(6x^2 + x - 1)(3x^2 + 2x - 1)(2x^2 + 3x + 1)} \\
 &= \sqrt{(3x - 1)(2x + 1)(3x - 1)(x + 1)(2x + 1)(x + 1)} = |(3x - 1)(2x + 1)(x + 1)|
 \end{aligned}$$

iii) First let us factorize the polynomials

$$\begin{aligned}
 \sqrt{15x^2 + (\sqrt{3} + \sqrt{10})x + \sqrt{2}} &= \sqrt{15x^2 + \sqrt{3}x + \sqrt{10}x + \sqrt{2}} \\
 &= \sqrt{3}x(\sqrt{5}x + 1) + \sqrt{2}(\sqrt{5}x + 1) \\
 &= (\sqrt{5}x + 1) \times (\sqrt{3}x + \sqrt{2}) \\
 \sqrt{5x^2 + (2\sqrt{5} + 1)x + 2} &= \sqrt{5x^2 + 2\sqrt{5}x + x + 2} \\
 &= \sqrt{5}x(x + 2) + 1(x + 2) = (\sqrt{5}x + 1)(x + 2) \\
 \sqrt{x^2 + (\sqrt{3} + \sqrt{2})x + 2} &= x(\sqrt{3}x + \sqrt{2}) + 2(\sqrt{3}x + \sqrt{2}) = (x + 2)(\sqrt{3}x + \sqrt{2})
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 & \sqrt{[\sqrt{15x^2 + (\sqrt{3} + \sqrt{10})x + \sqrt{2}}][\sqrt{5x^2 + (2\sqrt{5} + 1)x + 2}][\sqrt{3x^2 + (\sqrt{2} + 2\sqrt{3})x + 2\sqrt{2}}]} \\
 &= \sqrt{(\sqrt{5}x + 1)(\sqrt{3}x + \sqrt{2})(\sqrt{5}x + 1)(x + 2)(\sqrt{3}x + \sqrt{2})(x + 2)} = |(\sqrt{5}x + 1)(\sqrt{3}x + \sqrt{2})(x + 2)|
 \end{aligned}$$

EXERCISE 3.7**1. Find the square root of the following rational expressions.**

$$i) \frac{400x^4y^{12}z^{16}}{100x^8y^4z^4} \quad ii) \frac{7x^2 + 2\sqrt{14}x + 2}{x^2 - \frac{1}{2}x + \frac{1}{16}}$$

$$iii) \frac{121(a+b)^2(x+y)^8(b-c)^8}{81(b-c)^4(a-b)^{12}(b-c)^4}$$

Solution :

$$\begin{aligned}
 i) \quad & \sqrt{\frac{400x^4y^{12}z^{16}}{100x^8y^4z^4}} \\
 &= \frac{20}{10} \left| \frac{x^2y^6z^8}{x^4y^2z^2} \right| \\
 &= 2 \left| \frac{y^4z^6}{x^2} \right|
 \end{aligned}$$

$$\begin{aligned}
 \text{ii)} \quad & \sqrt{\frac{7x^2 + 2\sqrt{14}x + 2}{x^2 - \frac{1}{2}x + \frac{1}{16}}} \\
 &= \sqrt{\frac{(\sqrt{7}x + \sqrt{2})^2}{\left(x - \frac{1}{4}\right)^2}} \\
 &= \left| \frac{\sqrt{7}x + \sqrt{2}}{x - \frac{1}{4}} \right| \\
 &= 4 \left| \frac{\sqrt{7}x + \sqrt{2}}{4x - 1} \right|
 \end{aligned}$$

$$\begin{aligned}
 \text{iii)} \quad & \sqrt{\frac{121(a+b)^8 (x+y)^8 (b-c)^8}{81(b-c)^4 (a-b)^{12} (b-c)^4}} \\
 &= \frac{11}{9} \left| \frac{(a+b)^4 (x+y)^4 (b-c)^4}{(b-c)^2 (a-b)^6 (b-c)^2} \right| \\
 &= \frac{11}{9} \left| \frac{(a+b)^4 (x+y)^4}{(a-b)^6} \right|
 \end{aligned}$$

2. Find the square root of the following

i) $4x^2 + 20x + 25$

ii) $9x^2 - 24xy + 30xz - 40yz + 25z^2 + 16y^2$

iii) $1 + \frac{1}{x^6} + \frac{2}{x^3}$

iv) $(4x^2 - 9x + 2)(7x^2 - 13x - 2)(28x^2 - 3x - 1)$

v) $\left(2x^2 + \frac{17}{6}x + 1\right)\left(\frac{3}{2}x^2 + 4x + 2\right)\left(\frac{4}{3}x^2 + \frac{11}{3}x + 2\right)$

Solution :

$$\begin{aligned}
 \text{i)} \quad & \sqrt{4x^2 + 20x + 25} \\
 &= \sqrt{(2x + 5)^2} \\
 &= |2x + 5|
 \end{aligned}$$

$$\begin{aligned}
 \text{ii)} \quad & \sqrt{9x^2 - 24xy + 30xz - 40yz + 25z^2 + 16y^2} \\
 &= \sqrt{(3x - 4y + 5z)^2} \\
 &= |3x - 4y + 5z|
 \end{aligned}$$

$$\begin{aligned}
 \text{iii)} \quad & \sqrt{1 + \frac{1}{x^6} + \frac{2}{x^3}} \\
 &= \sqrt{\left(1 + \frac{1}{x^3}\right)^2} \\
 &= \left|1 + \frac{1}{x^3}\right|
 \end{aligned}$$

$$\begin{aligned}
 \text{iv)} \quad & \sqrt{(4x^2 - 9x + 2)(7x^2 - 13x - 2)(28x^2 - 3x - 1)} \\
 & \quad \quad \quad \text{(by factorisation)} \\
 &= \sqrt{(7x + 1)^2 (4x - 1)^2 (x - 2)^2} \\
 &= |(7x + 1)(4x - 1)(x - 2)|
 \end{aligned}$$

$$\begin{aligned}
 \text{v)} \quad & \sqrt{\left(2x^2 + \frac{17}{6}x + 1\right)\left(\frac{3}{2}x^2 + 4x + 2\right)\left(\frac{4}{3}x^2 + \frac{11}{3}x + 2\right)} \\
 &= \sqrt{\frac{(12x^2 + 17x + 6)}{6} \cdot \frac{(3x^2 + 8x + 4)}{2} \cdot \frac{(4x^2 + 11x + 6)}{3}} \\
 &= \sqrt{\frac{(4x + 3)(3x + 2) \cdot (3x + 2)(x + 2) \cdot (4x + 3)(x + 2)}{36}} \\
 &= \frac{1}{6} \sqrt{(4x + 3)^2 \cdot (3x + 2)^2 \cdot (x + 2)^2} \\
 &= \frac{1}{6} |(4x + 3)(3x + 2)(x + 2)|
 \end{aligned}$$

VIII. SQUARE ROOT OF POLYNOMIALS BY DIVISION METHOD:

Key Points

- ✓ The long division method in finding the square root of a polynomial is useful when the degree of the polynomial is higher.
- ✓ Before proceeding to find the square root of a polynomial, one has to ensure that the degrees of the variables are in descending or ascending order.

Example 3.21

Find the square root of $64x^4 - 16x^3 + 17x^2 - 2x + 1$

Solution :

$$\begin{array}{r}
 8x^2 - x + 1 \\
 8x^2 \overline{) 64x^4 - 16x^3 + 17x^2 - 2x + 1} \\
 \underline{64x^4} \qquad \qquad \qquad (-) \\
 16x^2 - x \qquad \qquad \qquad \underline{16x^2 - x} \qquad \qquad \qquad (-) \\
 16x^2 - 2x + 1 \qquad \qquad \underline{16x^2 - 2x + 1} \qquad \qquad \qquad (-) \\
 \qquad \qquad \qquad \qquad \qquad \underline{16x^2 - 2x + 1} \qquad \qquad \qquad (-) \\
 \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad 0
 \end{array}$$

Therefore

$$\sqrt{64x^4 - 16x^3 + 17x^2 - 2x + 1} = |8x^2 - x + 1|$$

Example 3.22

Find the square root of the expression

$$\frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2}$$

Solution :

$$\begin{array}{r}
 \frac{2x}{y} + 5 - \frac{3y}{x} \\
 \frac{2x}{y} \overline{) \frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2}} \\
 \underline{\frac{4x^2}{y^2}} \qquad \qquad \qquad (-) \\
 \frac{4x}{y} + 5 \overline{) \frac{20x}{y} + 13} \qquad \qquad \qquad (-) \\
 \underline{\frac{20x}{y} + 25} \\
 \frac{4x}{y} + 10 - \frac{3y}{x} \overline{) -12 - \frac{30y}{x} + \frac{9y^2}{x^2}} \qquad \qquad \qquad (-) \\
 \underline{-12 - \frac{30y}{x} + \frac{9y^2}{x^2}} \\
 \qquad \qquad \qquad \qquad \qquad \underline{0}
 \end{array}$$

Hence,

$$\sqrt{\frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2}} = \left| \frac{2x}{y} + 5 - \frac{3y}{x} \right|$$

Example 3.23

If $9x^4 + 12x^3 + 28x^2 + ax + b$ is a perfect square, find the values of a and b .

Solution :

$$\begin{array}{r}
 3x^2 + 2x + 4 \\
 3x^2 \overline{) 9x^4 + 12x^3 + 28x^2 + ax + b} \\
 \underline{9x^4} \quad (-) \\
 6x^2 + 2x \\
 \underline{6x^2 + 4x + 4} \quad (-) \\
 6x^2 + 4x + 4 \\
 \underline{6x^2 + 4x + 4} \quad (-) \\
 0
 \end{array}$$

Because the given polynomial is a perfect square $a - 16 = 0$, $b - 16 = 0$

Therefore $a = 16$, $b = 16$.

1. Find the square root of the following polynomials by division method

(i) $x^4 - 12x^3 + 42x^2 - 36x + 9$

(ii) $37x^2 - 28x^3 + 4x^4 + 42x + 9$

(iii) $16x^4 + 8x^2 + 1$

(iv) $121x^4 - 198x^3 - 183x^2 + 216x + 144$

Solution :

(i) $x^4 - 12x^3 + 42x^2 - 36x + 9$

$$\begin{array}{r}
 x^2 - 6x + 3 \\
 x^2 \overline{) x^4 - 12x^3 + 42x^2 - 36x + 9} \\
 \underline{x^4} \\
 2x^2 - 6x \\
 \underline{2x^2 - 6x} \\
 2x^2 - 12x + 3 \\
 \underline{2x^2 - 12x + 3} \\
 0
 \end{array}$$

$$\therefore \sqrt{x^4 - 12x^3 + 42x^2 - 36x + 9} = |x^2 - 6x + 3|$$

(ii) $37x^2 - 28x^3 + 4x^4 + 42x + 9$

$$\begin{array}{r}
 2x^2 - 7x - 3 \\
 2x^2 \overline{) 4x^4 - 28x^3 + 37x^2 + 42x + 9} \\
 \underline{4x^4} \\
 4x^2 - 7x \\
 \underline{4x^2 - 14x - 3} \\
 -12x^2 + 42x + 9 \\
 \underline{-12x^2 + 42x + 9} \\
 0
 \end{array}$$

$$\therefore \sqrt{4x^4 - 28x^3 + 37x^2 + 42x + 9} = |2x^2 - 7x - 3|$$

(iii) $16x^4 + 8x^2 + 1$

$$\begin{array}{r}
 4x^2 + 1 \\
 4x^2 \overline{) 16x^4 + 0x^3 + 8x^2 + 0x + 1} \\
 \underline{16x^4} \\
 8x^2 + 1 \\
 \underline{8x^2 + 1} \\
 0
 \end{array}$$

$$\therefore \sqrt{16x^4 + 8x^2 + 1} = |4x^2 + 1|$$

(iv) $121x^4 - 198x^3 - 183x^2 + 216x + 144$

$$\begin{array}{r}
 11x^2 - 9x - 12 \\
 11x^2 \overline{) 121x^4 - 198x^3 - 183x^2 + 216x + 144} \\
 \underline{121x^4} \\
 22x^2 - 9x \\
 \underline{22x^2 - 18x - 12} \\
 -198x^3 - 183x^2 \\
 \underline{-198x^3 + 81x^2} \\
 -264x^2 + 216x + 144 \\
 \underline{-264x^2 + 216x + 144} \\
 0
 \end{array}$$

$$\therefore \sqrt{121x^4 - 198x^3 - 183x^2 + 216x + 144} = |11x^2 - 9x - 12|$$

2. Find the square root of the expression

$$\frac{x^2}{y^2} - \frac{10x}{y} + 27 - \frac{10y}{x} + \frac{y^2}{x^2}$$

Solution :

$$\begin{array}{r} \frac{x}{y} - 5 + \frac{y}{x} \\ \hline \frac{x}{y} \left[\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2} \right] \\ \frac{x^2}{y^2} \\ \hline \frac{2x}{y} - 5 \quad \begin{array}{r} -10\frac{x}{y} + 27 \\ -10\frac{x}{y} + 25 \end{array} \\ \hline \frac{2x}{y} - 10 + \frac{y}{x} \quad \begin{array}{r} 2 - 10\frac{y}{x} + \frac{y^2}{x^2} \\ 2 - 10\frac{y}{x} + \frac{y^2}{x^2} \end{array} \\ \hline 0 \end{array}$$

$$\therefore \sqrt{\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}} = \left| \frac{x}{y} - 5 + \frac{y}{x} \right|$$

3. Find the values of a and b if the following polynomials are perfect squares

(i) $4x^4 - 12x^3 + 37x^2 + bx + a$ (ii) $ax^4 + bx^3 + 361x^2 + 220x + 100$

Solution :

$$\begin{array}{r} 2x^2 - 3x + 7 \\ \hline 2x^2 \quad \begin{array}{r} 4x^4 - 12x^3 + 37x^2 + bx + a \\ 4x^4 \end{array} \\ \hline 4x^2 - 3x \quad \begin{array}{r} -12x^3 + 37x^2 \\ -12x^3 + 9x^2 \end{array} \\ \hline \end{array}$$

$$\begin{array}{r} 4x^2 - 6x + 7 \quad \begin{array}{r} 28x^2 + bx + a \\ 28x^2 - 42x + 49 \\ (+) \quad (-) \\ \hline 0 \end{array} \end{array}$$

(The given polynomial is a perfect square)

$$\begin{aligned} \therefore a - 49 &= 0, & b + 42 &= 0 \\ a &= 49 & b &= -42 \end{aligned}$$

(ii) $ax^4 + bx^3 + 361x^2 + 220x + 100$

Solution :

$$\begin{array}{r} 10 + 11x + 12x^2 \\ \hline 10 \quad \begin{array}{r} 100 + 220x + 361x^2 + 6x^3 + ax^4 \\ 100 \end{array} \\ \hline 20 + 11x \quad \begin{array}{r} 220x + 361x^2 \\ 220x + 121x^2 \end{array} \\ \hline 20 + 22x + 12x^2 \quad \begin{array}{r} 240x^2 + \quad bx^3 + \quad ax^4 \\ \quad \quad \quad 2 \quad \quad \quad 3 \quad \quad \quad 4 \\ (-) \quad (-) \\ \hline 0 \end{array} \end{array}$$

(The given polynomial is a perfect square)

$$\begin{aligned} \therefore a - 144 &= 0, & b - 264 &= 0 \\ a &= 144 & b &= 264 \end{aligned}$$

4. Find the values of m and n if the following expressions are perfect squares

i) $\frac{1}{x^4} - \frac{6}{x^3} + \frac{13}{x^2} + \frac{m}{x} + n$

ii) $x^4 - 8x^3 + mx^2 + nx + 16$

Solution

$$\begin{array}{r} \frac{1}{x^2} - \frac{3}{x} + 2 \\ \hline \frac{1}{x^2} \quad \begin{array}{r} \frac{1}{x^4} - \frac{6}{x^3} + \frac{13}{x^2} + \frac{m}{x} + n \\ \frac{1}{x^4} \end{array} \end{array}$$

$$\frac{2}{x^2} - \frac{3}{x}$$

$$\frac{-6}{x^3} + \frac{13}{x^2}$$

$$\frac{-6}{x^3} + \frac{a}{x^2}$$

$$\frac{2}{x^2} - \frac{6}{x} + 2$$

$$\frac{4}{x^2} + \frac{m}{x} + n$$

$$\frac{4}{x^2} - \frac{12}{x} + 4$$

$$(+)\quad (-)$$

$$0$$

(The given polynomial is a perfect square)

$$\therefore m + 12 = 0, \quad n - 4 = 0$$

$$m = -12 \quad n = 4$$

$$\text{ii) } x^4 - 8x^3 + mx^2 + nx + 16$$

$$x^2 - 4x + 4$$

$$x^2 \quad x^4 - 8x^3 + mx^2 + nx + 16$$

$$x^4$$

$$2x^2 - 4x$$

$$-8x^3 + mx^2$$

$$-8x^3 + 16x^2$$

$$2x^2 - 8x + 4$$

$$(m - 16)x^2 + nx + 16$$

$$8x^2 - 32x + 16$$

$$(-)\quad (+)\quad (-)$$

$$0$$

(The given polynomial is a perfect square)

$$\therefore m - 16 - 8 = 0, \quad n + 32 = 0$$

$$m = 24 \quad n = -32$$

IX. QUADRATIC EQUATIONS, ZEROS AND ROOTS :

- ✓ A quadratic expression is an expression of degree n in variable x is $a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n$ where $a_0 \neq 0$ and a_1, a_2, \dots, a_n are real numbers, $a_0, a_1, a_2, \dots, a_n$ are called coefficients of the expression.
- ✓ An expression of degree 2 is called a Quadratic Expression which is expressed as $p(x) = ax^2 + bx + c$, $a \neq 0$ and a, b, c are real numbers.
- ✓ Let $p(x)$ be a polynomial. $x = a$ is called zero of $p(x)$ if $p(a) = 0$.
- ✓ Let $ax^2 + bx + c = 0$, ($a \neq 0$) be a quadratic equation. The values of x such that the expression $ax^2 + bx + c$ becomes zero are called roots of the quadratic equation $ax^2 + bx + c = 0$.

- ✓ If α and β are roots of a quadratic equation $ax^2 + bx + c = 0$ then

$$\alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad \beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

$$\checkmark \quad \alpha + \beta = \frac{-b + \sqrt{b^2 - 4ac} - b - \sqrt{b^2 - 4ac}}{2a} = \frac{-b}{a}$$

$$\checkmark \quad \alpha\beta = \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \times \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) = \frac{c}{a}$$

- ✓ $x^2 - (\text{sum of roots})x + \text{product of roots} = 0$ is the general form of the quadratic equation when the roots are given.

Example 3.24

Find the zeroes of the quadratic expression $x^2 + 8x + 12$.

Solution :

$$\text{Let } p(x) = x^2 + 8x + 12 = (x + 2)(x + 6)$$

$$p(-2) = 4 - 16 + 12 = 0$$

$$p(-6) = 36 - 48 + 12 = 0$$

Therefore -2 and -6 are zeros of $p(x) = x^2 + 8x + 12$

Example 3.25

Write down the quadratic equation in general form for which sum and product of the roots are given below.

$$i) 9, 14 \quad ii) -\frac{7}{2}, \frac{5}{2} \quad iii) -\frac{3}{5}, -\frac{1}{2}$$

Solution :

(i) General form of the quadratic equation when the roots are given is

$$x^2 - (\text{sum of the roots})x + \text{product of the roots} = 0$$

$$x^2 - 9x + 14 = 0$$

$$ii) x^2 - \left(-\frac{7}{2}\right)x + \frac{5}{2} = 0 \text{ gives } 2x^2 + 7x + 5 = 0$$

$$iii) x^2 - \left(-\frac{3}{5}\right)x + \left(-\frac{1}{2}\right) = 0 \Rightarrow \frac{10x^2 + 6x - 5}{10} = 0$$

$$\text{Therefore } 10x^2 + 6x - 5 = 0$$

Example 3.26

Find the sum and product of the roots for each of the following quadratic equations :

$$(i) x^2 + 8x - 65 = 0 \quad (ii) 2x^2 + 5x + 7 = 0$$

$$(iii) kx^2 - k^2x - 2k^3 = 0$$

Solution :

Let α and β be the roots of the given quadratic equation

$$(i) x^2 + 8x - 65 = 0$$

$$a = 1, b = 8, c = -65$$

$$\alpha + \beta = -\frac{b}{a} = -8 \text{ and } \alpha\beta = \frac{c}{a} = -65$$

$$\alpha + \beta = -8 ; \alpha\beta = -65$$

$$(ii) 2x^2 + 5x + 7 = 0$$

$$a = 2, b = 5, c = 7$$

$$\alpha + \beta = -\frac{b}{a} = -\frac{5}{2} \text{ and } \alpha\beta = \frac{c}{a} = \frac{7}{2}$$

$$\alpha + \beta = -\frac{5}{2} ; \alpha\beta = \frac{7}{2}$$

$$(iii) kx^2 - k^2x - 2k^3 = 0$$

$$a = k, b = -k^2, c = -2k^3$$

$$\alpha + \beta = -\frac{b}{a} = \frac{-(-k^2)}{k}$$

$$\text{and } \alpha\beta = \frac{c}{a} = \frac{-2k^3}{k} = -2k^2$$

EXERCISE 3.9

1. Determine the quadratic equations, whose sum and product of roots are

$$i) -9, 20 \quad ii) \frac{5}{3}, 4$$

$$iii) \frac{-3}{2}, -1 \quad iv) -(2-a)^2, (a+5)^2$$

Solution :

i) Given

$$\text{Sum of the roots, } \text{SOR} = -9$$

$$\text{Product of the roots, } \text{POR} = 20$$

The required quadratic equation is

$$x^2 - (\text{SOR})x + (\text{POR}) = 0$$

$$\Rightarrow x^2 - (-9)x + 20 = 0$$

$$\Rightarrow x^2 + 9x + 20 = 0$$

- ii) Given $\text{SOR} = \frac{5}{3}$, $\text{POR} = 4$
 \therefore The required quadratic equation is

$$x^2 - \frac{5}{3}x + 4 = 0$$

$$\Rightarrow 3x^2 - 5x + 12 = 0$$

- iii) Given $\text{SOR} = \frac{-3}{2}$, $\text{POR} = -1$
 \therefore The required equation is

$$x^2 - \left(\frac{-3}{2}\right)x + (-1) = 0$$

$$\Rightarrow 2x^2 + 3x - 2 = 0$$

- iv) Given $\text{SOR} = -(2-a)^2$, $\text{POR} = (a+5)^2$
 \therefore The required equation is

$$x^2 - (-(2-a)^2)x + (a+5)^2 = 0$$

$$\Rightarrow x^2 + (2-a)^2x + (a+5)^2 = 0$$

2. Find the sum and product of the roots for each of the following quadratic equations

(i) $x^2 + 3x - 28 = 0$ (ii) $x^2 + 3x = 0$

iii) $3 + \frac{1}{a} = \frac{10}{a^2}$ (iv) $3y^2 - y - 4 = 0$

Solution :

i) $x^2 + 3x - 28 = 0$

Given equation is $x^2 + 3x - 28 = 0$

$a = 1$, $b = 3$, $c = -28$

\therefore Sum of the roots $= \alpha + \beta = \frac{-b}{a} = \frac{-3}{1} = -3$

Product of the roots $= \alpha \beta = \frac{c}{a} = \frac{-28}{1} = -28$

ii) $x^2 + 3x = 0$

$a = 1$, $b = 3$, $c = 0$

$\therefore \alpha + \beta = \frac{-b}{a} = \frac{-3}{1} = -3$

$\alpha \beta = \frac{c}{a} = \frac{0}{1} = 0$

iii) Given $3 + \frac{1}{a} = \frac{10}{a^2}$

$\Rightarrow \frac{3a+1}{a} = \frac{10}{a^2}$

$\Rightarrow 3a+1 = \frac{10}{a}$

$\Rightarrow 3a^2 + a - 10 = 0$

$A = 3$, $B = 1$, $C = -10$

$\frac{-B}{A} = \frac{-1}{3}$

$\alpha \beta = \frac{C}{A} = \frac{-10}{3}$

(iv) $3y^2 - y - 4 = 0$

$a = 3$, $b = -1$, $c = -4$

$\therefore \alpha + \beta = \frac{-b}{a} = \frac{1}{3}$

$\alpha \beta = \frac{c}{a} = \frac{-4}{3}$

X. SOLVING QUADRATIC EQUATIONS BY FACTORISATION :**Example 3.27**

Solve $2x^2 - 2\sqrt{6}x + 3 = 0$

Solution :

$$2x^2 - 2\sqrt{6}x + 3 = 2x^2 - \sqrt{6}x - \sqrt{6}x + 3$$

(by splitting the middle term)

$$\begin{aligned}
 &= \sqrt{2}x(\sqrt{2}x - \sqrt{3}) - \sqrt{3}(\sqrt{2}x - \sqrt{3}) \\
 &= (\sqrt{2}x - \sqrt{3})(\sqrt{2}x - \sqrt{3})
 \end{aligned}$$

Now, equating the factors to zero we get,

$$(\sqrt{2}x - \sqrt{3})(\sqrt{2}x - \sqrt{3}) = 0$$

$$\sqrt{2}x - \sqrt{3} = 0 \quad \text{or} \quad \sqrt{2}x - \sqrt{3} = 0$$

$$\sqrt{2}x = \sqrt{3} \quad \text{or} \quad \sqrt{2}x = \sqrt{3}$$

$$\text{Therefore the solution is } x = \frac{\sqrt{3}}{\sqrt{2}}$$

Example 3.28

Solve $2m^2 + 19m + 30 = 0$

Solution :

$$2m^2 + 19m + 30 = 2m^2 + 4m + 15m + 30$$

$$= 2m(m + 2) + 15(m + 2)$$

$$= (m + 2)(2m + 15)$$

Now, equating the factors to zero we get,

$$(m + 2)(2m + 15) = 0$$

$$m + 2 = 0 \text{ gives, } m = -2 \text{ or } 2m + 15 = 0$$

$$\text{we get, } m = \frac{-15}{2}$$

$$\text{Therefore the roots are } -2, \frac{-15}{2}$$

Some equations which are not quadratic can be solved by reducing them to quadratic equations by suitable substitutions. Such examples are illustrated below.

Example 3.29

Solve $x^4 - 13x^2 + 42 = 0$

Solution :

$$\text{Let } x^2 = a. \text{ Then, } (x^2)^2 - 13x^2 + 42 = a^2 - 13a + 42 = (a - 7)(a - 6)$$

$$\text{Given, } (a - 7)(a - 6) = 0 \text{ we get, } a = 7 \text{ or } 6.$$

$$\text{Since } a = x^2, x^2 = 7 \text{ then, } x = \pm\sqrt{7} \text{ or } x^2 = 6 \text{ we get, } x = \pm\sqrt{6}$$

$$\text{Therefore the roots are } x = \pm\sqrt{7}, \pm\sqrt{6}$$

Example 3.30

Solve $\frac{x}{x-1} + \frac{x-1}{x} = 2\frac{1}{2}$

$$\text{Let } y = \frac{x}{x-1} \text{ then } \frac{1}{y} = \frac{x-1}{x}$$

$$\text{Therefore, } \frac{x}{x-1} + \frac{x-1}{x} = 2\frac{1}{2} \text{ becomes } y + \frac{1}{y} = \frac{5}{2}$$

$$2y^2 - 5y + 2 = 0 \text{ then, } y = \frac{1}{2}, 2$$

$$\frac{x}{x-1} = \frac{1}{2} \text{ we get, } 2x = x - 1 \text{ implies } x = -1$$

$$\frac{x}{x-1} = 2 \text{ we get, } x = 2x - 2 \text{ implies } x = 2$$

$$\text{Therefore the roots are } x = -1, 2$$

EXERCISE 3.10

1. Solve the following quadratic equations by factorization method

i) $4x^2 - 7x - 2 = 0$

ii) $3(p^2 - 6) = p(p + 5)$

iii) $\sqrt{a(a-7)} = 3\sqrt{2}$

iv) $\sqrt{2x^2 + 7x + 5\sqrt{2}} = 0$

v) $2x^2 - x + \frac{1}{8} = 0$

Solution :

i) Given $4x^2 - 7x - 2 = 0$

-7	-8
-8	+1

$$\Rightarrow 4x^2 - 8x + x - 2 = 0$$

$$\Rightarrow 4x(x-2) + 1(x-2) = 0$$

$$\Rightarrow (4x+1)(x-2) = 0$$

$$\Rightarrow 4x = -1 \text{ (or) } x - 2 = 0$$

$$x = -\frac{1}{4} \text{ (or) } x = 2$$

$$\text{Roots are } \left\{ -\frac{1}{4}, 2 \right\}$$

ii) Given $3(p^2 - 6) = p(p + 5)$

-5	-36
-9	+4
2	2

$$\Rightarrow 3p^2 - 18 = p^2 + 5p$$

$$\Rightarrow 2p^2 - 5p - 18 = 0$$

$$\Rightarrow p = \frac{9}{2}, -2$$

iii) Given $\sqrt{a(a-7)} = 3\sqrt{2}$

Squaring on both sides

$$a^2 - 7a = 18$$

$$a^2 - 7a - 18 = 0$$

$$(a-9)(a+2) = 0$$

$$a = 9 \text{ (or) } -2$$

$$\text{Roots are } 9, -2$$

iv) $\sqrt{2x^2 + 7x + 5\sqrt{2}} = 0$

7	10
5	2

$$\Rightarrow \sqrt{2x^2 + 2x + 5x + 5\sqrt{2}} = 0$$

$$\Rightarrow \sqrt{2x}(x + \sqrt{2}) + 5(x + \sqrt{2}) = 0$$

$$\Rightarrow (\sqrt{2x} + 5)(x + \sqrt{2}) = 0$$

$$\therefore \sqrt{2x} + 5 = 0 \text{ (or) } x + \sqrt{2} = 0$$

$$\Rightarrow x = -\frac{5}{\sqrt{2}} \text{ (or) } x = -\sqrt{2}$$

$$\therefore \text{Roots are } -\frac{5}{\sqrt{2}}, -\sqrt{2}$$

v) $2x^2 - x + \frac{1}{8} = 0$

$$\Rightarrow 16x^2 - 8x + 1 = 0$$

$$\Rightarrow 16x^2 - 4x - 4x + 1 = 0$$

$$\Rightarrow 4x(4x-1) - 1(4x-1) = 0$$

$$\Rightarrow (4x-1)(4x-1) = 0$$

$$x = \frac{1}{4}, \frac{1}{4}$$

$$\therefore \text{Roots are } \frac{1}{4}, \frac{1}{4}$$

2. The number of volleyball games that must be scheduled in a league with n teams is given by $G(n) = \frac{n^2 - n}{2}$ where each team plays with every other team exactly once.

A league schedules 15 games. How many teams are in the league?

Solution :

By data given, $G(n) = \frac{n^2 - n}{2} = 15$

$$\Rightarrow n^2 - n = 30$$

$$\Rightarrow n^2 - n - 30 = 0$$

$$\Rightarrow (n-6)(n+5) = 0$$

$$\Rightarrow n = 6, -5$$

$$\therefore \text{Number of terms in the league} = 6$$

XI. SOLVING QUADRATIC EQUATIONS BY COMPLETING THE SQUARES AND BY FORMULA:**Key Points**

- ✓ **Step 1** Write the quadratic equation in general form $ax^2 + bx + c = 0$.
- ✓ **Step 2** Divide both sides of the equation by the coefficient of x^2 if it is not 1.
- ✓ **Step 3** Shift the constant term to the right hand side.
- ✓ **Step 4** Add the square of one-half of the coefficient of x to both sides.
- ✓ **Step 5** Write the left hand side as a square and simplify the right hand side.
- ✓ **Step 6** Take the square root on both sides and solve for x .
- ✓ The formula for finding roots of a quadratic equation $ax^2 + bx + c = 0$ is $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Example 3.31Solve $x^2 - 3x - 2 = 0$

$$x^2 - 3x - 2 = 0$$

$$x^2 - 3x = 2 \quad (\text{Shifting the Constant to RHS})$$

$$x^2 - 3x + \left(\frac{3}{2}\right)^2 = 2 + \left(\frac{3}{2}\right)^2$$

$$\left(\text{Add } \left| \frac{1}{2} (\text{co-efficient of } x) \right|^2 \text{ to both sides} \right)$$

$$\left(x - \frac{3}{2} \right)^2 = \frac{17}{4}$$

(writing the LHS as complete square)

$$x - \frac{3}{2} = \pm \frac{\sqrt{17}}{2}$$

(Taking the square root on both sides)

$$x = \frac{3}{2} + \frac{\sqrt{17}}{2} \text{ or } x = \frac{3}{2} - \frac{\sqrt{17}}{2}$$

Therefore, $x = \frac{3 + \sqrt{17}}{2}, \frac{3 - \sqrt{17}}{2}$

Example 3.32Solve $2x^2 - x - 1 = 0$

$$2x^2 - x - 1 = 0$$

$$x^2 - \frac{x}{2} - \frac{1}{2} = 0$$

($\div 2$ make co-efficient of x^2 as 1)

$$x^2 - \frac{x}{2} = \frac{1}{2}$$

$$x^2 - \frac{x}{2} + \left(\frac{1}{4}\right)^2 = \frac{1}{2} + \left(\frac{1}{4}\right)^2$$

$$\left(x - \frac{1}{4} \right)^2 = \frac{9}{16} = \left(\frac{3}{4} \right)^2$$

$$x - \frac{1}{4} = \pm \frac{3}{4} \Rightarrow x = 1, -\frac{1}{2}$$

Example 3.33Solve $x^2 + 2x - 2 = 0$ by formula method**Solution :**

Compare $x^2 + 2x - 2 = 0$ with the standard form $ax^2 + bx + c = 0$

$$a = 1, b = 2, c = -2$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

substituting the values of a, b and c in the formula we get,

$$x = \frac{-2 \pm \sqrt{(2)^2 - 4(1)(-2)}}{2(1)} = \frac{-2 \pm \sqrt{12}}{2} = -1 \pm \sqrt{3}$$

$$\text{Therefore, } x = -1 + \sqrt{3}, -1 - \sqrt{3}$$

Example 3.34

Solve $2x^2 - 3x - 3 = 0$ by formula method

Solution :

Compare $2x^2 - 3x - 3 = 0$ with the standard form $ax^2 + bx + c = 0$

$$a = 2, b = -3, c = -3$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

substituting the values of a, b and c in the formula we get,

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(2)(-3)}}{2(2)} = \frac{3 \pm \sqrt{33}}{4}$$

$$\text{Therefore, } x = \frac{3 + \sqrt{33}}{4}, x = \frac{3 - \sqrt{33}}{4}$$

Example 3.35

Solve $3p^2 + 2\sqrt{5}p - 5 = 0$ by formula method

Solution :

Compare $3p^2 + 2\sqrt{5}p - 5 = 0$ with the standard form $ax^2 + bx + c = 0$

$$a = 3, b = 2\sqrt{5}, c = -5$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

substituting the values of a, b and c in the formula we get,

$$p = \frac{-2\sqrt{5} \pm \sqrt{(2\sqrt{5})^2 - 4(3)(-5)}}{2(3)}$$

$$= \frac{-2\sqrt{5} \pm \sqrt{80}}{6}$$

$$= \frac{-\sqrt{5} \pm 2\sqrt{5}}{3}$$

$$\text{Therefore, } x = \frac{\sqrt{5}}{3}, -\sqrt{5}$$

Example 3.36

Solve $pqx^2 = (p + q)^2 x + (p + q)^2 = 0$ by formula method

Solution :

Compare the coefficients of the given equation

$$a = pq, b = -(p + q)^2, c = (p + q)^2$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

substituting the values of a, b and c in the formula we get,

$$x = \frac{[-(p + q)^2] \pm \sqrt{[-(p + q)^2]^2 - 4(pq)(p + q)^2}}{2pq}$$

$$= \frac{(p + q)^2 \pm \sqrt{(p + q)^4 - 4(pq)(p + q)^2}}{2pq}$$

$$= \frac{(p + q)^2 \pm \sqrt{(p + q)^2 [(p + q)^2 - 4pq]}}{2pq}$$

$$= \frac{(p + q)^2 \pm \sqrt{(p + q)^2 (p^2 + q^2 + 2pq - 4pq)}}{2pq}$$

$$= \frac{(p + q)^2 \pm \sqrt{(p + q)^2 (p - q)^2}}{2pq}$$

$$= \frac{(p + q)^2 \pm (p + q)(p - q)}{2pq}$$

$$\text{Therefore, } x = \frac{p+q}{2pq} \times pq, \frac{p+q}{2pq} \times 2q$$

$$\text{we get, } x = \frac{p+q}{q}, \frac{p+q}{p}$$

EXERCISE 3.11

1. Solve the following quadratic equations by completing the square method

i) $9x^2 - 12x + 4 = 0$ ii) $\frac{5x+7}{x-1} = 3x+2$

Solution :

- i) Given equation is

$$9x^2 - 12x + 4 = 0$$

$$\Rightarrow 9x^2 - 12x = -4$$

$$\Rightarrow x^2 - \frac{12}{9}x = \frac{-4}{9} \quad (\div \text{ by } 9)$$

$$\Rightarrow x^2 - \frac{4}{3}x = \frac{-4}{9}$$

$$\Rightarrow x^2 - \frac{4}{3}x + \left(\frac{2}{3}\right)^2 = \frac{-4}{9} + \left(\frac{2}{3}\right)^2$$

$$\Rightarrow \left(x - \frac{2}{3}\right)^2 = \frac{-4}{9} + \frac{4}{9}$$

$$\Rightarrow \left(x - \frac{2}{3}\right)^2 = 0$$

$$\Rightarrow \left(x - \frac{2}{3}\right)\left(x - \frac{2}{3}\right) = 0$$

$$\Rightarrow x = \frac{2}{3}, \frac{2}{3}$$

$$\therefore \text{Solution set} = \left\{\frac{2}{3}, \frac{2}{3}\right\}$$

- ii) Given equation is

$$\frac{5x+7}{x-1} = 3x+2$$

$$\Rightarrow 5x+7 = (3x+2)(x-1)$$

$$\Rightarrow 5x+7 = 3x^2 - x - 2$$

$$\Rightarrow 3x^2 - 6x - 9 = 0$$

$$\Rightarrow x^2 - 2x - 3 = 0$$

$$\Rightarrow x^2 - 2x = 3$$

$$\Rightarrow x^2 - 2x + 1 = 3 + 1$$

$$\Rightarrow (x-1)^2 = 4$$

$$\Rightarrow (x-1) = \pm 2$$

$$\Rightarrow x-1 = 2, x-1 = -2$$

$$\Rightarrow x = 3, x = -1$$

$$\text{Solution set} = \{3, -1\}$$

2. Solve the following quadratic equations by formula method

i) $2x^2 - 5x + 2 = 0$

ii) $\sqrt{2}f^2 - 6f + 3\sqrt{2} = 0$

iii) $3y^2 - 20y - 23 = 0$

iv) $36y^2 - 12ay + (a^2 - b^2) = 0$

- i) Given equation is

$$2x^2 - 5x + 2 = 0$$

$$a = 2, b = -5, c = 2$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{5 \pm \sqrt{25 - 16}}{4}$$

$$= \frac{5 \pm \sqrt{9}}{4}$$

$$= \frac{5 \pm 3}{4}$$

$$= \frac{5+3}{4}, \frac{5-3}{4}$$

$$= 2, \frac{1}{2}$$

ii) Given equation is

$$\sqrt{2}f^2 - 6f + 3\sqrt{2} = 0$$

$$a = \sqrt{2}, b = -6, c = 3\sqrt{2}$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{6 \pm \sqrt{36 - 4(\sqrt{2})(3\sqrt{2})}}{2\sqrt{2}} \\ &= \frac{6 \pm \sqrt{36 - 24}}{2\sqrt{2}} \\ &= \frac{6 \pm 2\sqrt{3}}{2\sqrt{2}} \\ &= \frac{6 + 2\sqrt{3}}{2\sqrt{2}}, \frac{6 - 2\sqrt{3}}{2\sqrt{2}} \\ &= \frac{3 + \sqrt{3}}{\sqrt{2}}, \frac{3 - \sqrt{3}}{\sqrt{2}} \end{aligned}$$

iii) Given equation is

$$3y^2 - 20y - 23 = 0$$

$$a = 3, b = -20, c = -23$$

$$\begin{aligned} y &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{20 \pm \sqrt{400 - 4(3)(-23)}}{6} \\ &= \frac{20 \pm \sqrt{400 + 276}}{6} \\ &= \frac{20 \pm \sqrt{676}}{6} \\ &= \frac{20 \pm 26}{6} \\ &= \frac{20 + 26}{6}, \frac{20 - 26}{6} \\ &= \frac{46}{6}, \frac{-6}{6} \\ &= \frac{23}{3}, -1 \end{aligned}$$

iv) Given equation is

$$36y^2 - 12ay + (a^2 - b^2) = 0$$

$$A = 36, B = -12a, C = a^2 - b^2$$

$$\begin{aligned} \therefore y &= \frac{-B \pm \sqrt{B^2 - 4AC}}{2A} \\ &= \frac{12a \pm \sqrt{144a^2 - 4(36)(a^2 - b^2)}}{2(36)} \\ &= \frac{12a \pm \sqrt{144a^2 - 144(a^2 - b^2)}}{72} \\ &= \frac{12a \pm \sqrt{144b^2}}{72} \\ &= \frac{12a \pm 12b}{72} \\ &= \frac{a \pm b}{6} \\ &= \frac{a+b}{6}, \frac{a-b}{6} \end{aligned}$$

3. A ball rolls down a slope and travels a distance $dt = t^2 - 0.75t$ feet in t seconds. Find the time when the distance travelled by the ball is 11.25 feet.

Solution :

By data given,

$$d = t^2 - 0.75t \text{ where } d = 11.25 \text{ ft}$$

$$\Rightarrow t^2 - 0.75t = 11.25$$

$$\Rightarrow t^2 - 0.75t - 11.25 = 0$$

$$\Rightarrow (t - 3.75)(t + 3) = 0$$

$$\Rightarrow t - 3.75 = 0, \quad t + 3 = 0$$

$$\therefore t = 3.75 \quad t = -3$$

$$\text{But } t \neq -3$$

$$\therefore t = 3.75 \text{ sec}$$

XII. SOLVING PROBLEMS INVOLVING QUADRATIC EQUATIONS :

Example 3.37

The product of Kumaran's age (in years) two years ago and his age four years from now is one more than twice his present age. What is his present age?

Solution :

Let the present age of Kumaran be x years.

Two years ago, his age = $(x - 2)$ years.

Four years from now, his age = $(x + 4)$ years.

Given, $(x - 2)(x + 4) = 1 + 2x$

$x^2 + 2x - 8 = 1 + 2x$ gives $(x - 3)(x + 3) = 0$
then, $x = \pm 3$

Therefore, $x = 3$ (Rejecting -3 as age cannot be negative)

Kumaran's present age is 3 years.


Example 3.38

A ladder 17 feet long is leaning against a wall. If the ladder, vertical wall and the floor from the bottom of the wall to the ladder form a right triangle, find the height of the wall where the top of the ladder meets if the distance between bottom of the wall to bottom of the ladder is 7 feet less than the height of the wall?

Solution :

Let the height of the wall $AB = x$ feet

As per the given data $BC = (x - 7)$ feet



In the right triangle ABC, $AC = 17$ ft,

$$BC = (x - 7) \text{ feet}$$

By Pythagoras theorem, $AC^2 = AB^2 + BC^2$

$$(17)^2 = x^2 + (x - 7)^2 ; 289 = x^2 + x^2 - 14x + 49$$

$$x^2 - 7x - 120 = 0 \text{ hence, } (x - 15)(x + 8) = 0 \text{ then, } x = 15 \text{ (or) } -8$$

Therefore, height of the wall $AB = 15$ ft
(Rejecting -8 as height cannot be negative).

Example 3.39

A flock of swans contained x^2 members. As the clouds gathered, $10x$ went to a lake and one-eighth of the members flew away to a garden. The remaining three pairs played about in the water. How many swans were there in total?

Solution :

As given there are x^2 swans

As per the given data $x^2 - 10x - \frac{1}{8}x^2 = 6$
we get, $7x^2 - 80x - 48 = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{80 \pm \sqrt{6400 - 4(7)(-48)}}{14}$$

$$= \frac{80 \pm 88}{14}$$

$$\text{Therefore, } x = 12, -\frac{4}{7}$$

Here $x = -\frac{4}{7}$ is not possible as the number of swans cannot be negative.

Hence, $x = 12$. Therefore total number of swans is $x^2 = 144$.

Example 3.40

A passenger train takes 1 hr more than an express train to travel a distance of 240 km from Chennai to Virudhachalam. The speed of passenger train is less than that of an express train by 20 km per hour. Find the average speed of both the trains.

Solution :

Let the average speed of passenger train be x km/hr.

Then the average speed of express train will be $(x + 20)$ km/hr

Time taken by the passenger train to cover distance of 240 km = $\frac{240}{x}$ hr

Time taken by express train to cover distance = $\frac{240}{x + 20}$

$$\text{Given, } \frac{240}{x} = \frac{240}{x + 20} + 1$$

$$240 \left| \frac{1}{x} - \frac{1}{x + 20} \right| = 1 \text{ gives, } 240 \left| \frac{x + 20 - x}{x(x + 20)} \right| = 1$$

$$\text{we get, } 4800 = (x^2 + 20x)$$

$x^2 + 2x - 4800 = 0$ gives, $(x + 80)(x - 60) = 0$ we get, $x = -80$ or 60 .

Therefore $x = 60$ (Rejecting -80 as speed cannot be negative)

Average speed of the passenger train is 60 km/hr

Average speed of the express train is 80 km/hr.

EXERCISE 3.12

1. If the difference between a number and its reciprocal is $\frac{24}{5}$, find the number.

Solution :

Let x be the required number

$\frac{1}{x}$ be its reciprocal

$$\text{Given } x - \frac{1}{x} = \frac{24}{5}$$

$$\Rightarrow \frac{x^2 - 1}{x} = \frac{24}{5}$$

$$\Rightarrow 5x^2 - 5 = 24x$$


$$\Rightarrow 5x^2 - 24x - 5 = 0$$

$$\Rightarrow x = 5, -\frac{1}{5}$$

-24	-25
$\frac{-25}{5}$	$\frac{+1}{5}$
$-5,$	$\frac{1}{5}$

\therefore The required numbers are $5, -\frac{1}{5}$

2. A garden measuring 12m by 16m is to have a pedestrian pathway that is ' w ' meters wide installed all the way around so that it increases the total area to 285 m². What is the width of the pathway?

Solution :

Given the dimensions of the garden

$$= 16\text{m} \times 12\text{m}$$

Let 'w' be the equal width of the pedestrian pathway.

\therefore By the data given,

56	- 372
$\frac{62}{4}$	$\frac{-6}{4}$
$\frac{31}{2}$	$\frac{-3}{2}$

$$\Rightarrow (16 + 2w)(12 + 2w) = 285$$

$$\Rightarrow 4w^2 + 56w + 192 - 285 = 0$$

$$\Rightarrow 4w^2 + 56w - 93 = 0$$

$$\Rightarrow -31\frac{1}{2} \quad 3\frac{1}{2}$$

But w can't be -ve

$$\therefore w = 3\frac{1}{2}$$

$$= 1.5 \text{ m}$$

\therefore Width of the path away = 1.5 m

- 3. A bus covers a distance of 90 km at a uniform speed. Had the speed been 15 km/hour more it would have taken 30 minutes less for the journey. Find the original speed of the bus.**

Solution :

Let 'x' Km/hr be the original speed of the bus.

Distance covered at a uniform speed = 90 Km

$$\text{Time taken} = \frac{90}{x}$$

Had the speed been 15 Km/hr more,

$$\text{Time taken} = \frac{90}{x+15}$$

\therefore By Data given

$$\frac{90}{x+15} - \frac{90}{x} = \frac{1}{2} \quad \left(\because 30 \text{ min} = \frac{1}{2} \text{ hr} \right)$$

$$\Rightarrow 90 \left(\frac{1}{x+15} - \frac{1}{x} \right) = \frac{1}{2}$$

$$\Rightarrow \frac{x - x - 15}{x(x+15)} = \frac{1}{180}$$

$$\Rightarrow 180 \times (-15) = x^2 + 15x$$

$$\Rightarrow x^2 + 15x - 2700 = 0$$

$$\Rightarrow (x+60)(x-45) = 0$$

$$\therefore x = -60, 45$$

\therefore Original speed = 45 Km/hr

- 4. A girl is twice as old as her sister. Five years hence, the product of their ages (in years) will be 375. Find their present ages.**

Solution :

Let the present ages of the girl and her sister be x, y

By data given,

i) $x = 2y$

ii) $(x+5)(y+5) = 375$

$$\Rightarrow (2y+5)(y+5) = 375$$

$$\Rightarrow 2y^2 + 15y - 350 = 0$$

$$\Rightarrow y = -35\frac{1}{2}, 10$$

y can't be -ve


$$\therefore y = 10$$

$$\therefore x = 2y \quad \Rightarrow \quad x = 20$$

\therefore Their present ages are 20, 10 years old.

5. A pole has to be erected at a point on the boundary of a circular ground of diameter 20 m in such a way that the difference of its distances from two diametrically opposite fixed gates P and Q on the boundary is 4 m. Is it possible to do so? If answer is yes at what distance from the two gates should the pole be erected?

Solution :



In the fig, $PQ = 20$ m = diameter of the circle with centre O.

Let A be the point of pole s.t

$$AP - AQ = 4\text{m} \quad AP = x, \quad AQ = x - 4$$

$$\text{Also, } PA^2 + QA^2 = PQ^2$$

(Angle in a semicircle is 90°)

$$\Rightarrow x^2 + (x - 4)^2 = 20^2$$

$$\Rightarrow 2x^2 - 8x + 16 - 400 = 0$$

$$\Rightarrow 2x^2 - 8x - 384 = 0$$

$$\Rightarrow x^2 - 4x - 192 = 0$$

$$\Rightarrow (x - 16)(x + 12) = 0$$

$$\therefore x = 16 \text{ m}$$

$$x = 16 \Rightarrow x - 4 = 12\text{m}$$

\therefore Pole should be erected at a distance of 16 m, 12m from the two gates.

6. From a group of $2x^2$ black bees, square root of half of the group went to a tree. Again eight-ninth of the bees went to the same tree. The remaining two got caught up in a fragrant lotus. How many bees were there in total ?

Solution :

$$\text{Given number of black bees} = 2x^2$$

By the data given,

$$2x^2 - x - \frac{8}{9}(2x^2) = 2$$

$$\Rightarrow 2x^2 \left(1 - \frac{8}{9}\right) - x = 2$$

$$\Rightarrow 2x^2 \left(\frac{1}{9}\right) - x = 2$$

$$\Rightarrow 2x^2 - 9x = 18$$

$$\Rightarrow 2x^2 - 9x - 18 = 0$$

$$\Rightarrow x = 6, \quad \frac{-}{2}$$

$$\text{But } x \neq \frac{-3}{2}$$

$$\therefore x = 6$$

- 9	- 36
$\frac{-12}{2}$	$\frac{+3}{2}$
- 6	$\frac{3}{2}$


$$\therefore \text{Total number of bees} = 2x^2$$

$$= 2(36)$$

$$= 72$$

7. Music is been played in two opposite galleries with certain group of people. In the first gallery a group of 4 singers were singing and in the second gallery 9 singers were singing. The two galleries are separated by the distance of 70 m. Where should a person stand for hearing the same intensity of the singers voice? (Hint: The ratio of the sound intensity is equal to the square of the ratio of their corresponding distances).

Solution :



galleries be P, who stand at a distance of 'd' m.

$$\therefore G_1 G_2 = 70\text{m}, G_1 P = d\text{m}, G_2 P = (70 - d)\text{m}$$

Since the ratio of sound intensity is equal to the square of the ratio of their corresponding sides,

$$\frac{4}{9} = \frac{d^2}{(70 - d)^2}$$

$$\Rightarrow 4(70 - d)^2 = 9d^2$$

$$\Rightarrow 4(4900 - 140d + d^2) = 9d^2$$

$$\Rightarrow 19600 - 560d + 4d^2 = 9d^2$$

$$\Rightarrow 5d^2 + 560d - 19600 = 0$$

$$\Rightarrow d^2 + 112d - 3920 = 0$$

$$\Rightarrow (d + 140)(d - 28) = 0$$

$$\Rightarrow d = -140 \text{ (or) } 28$$

$$\therefore d = 28\text{m}$$


The person should stand 28m from gallery 1

(or)

42m from gallery-2 to hear the same intensity of the singers voice.

8. There is a square field whose side is 10 m. A square flower bed is prepared in its centre leaving a gravel path all round the flower bed. The total cost of laying the flower bed and gravelling the path at ₹ 3 and ₹ 4 per square metre respectively is ₹ 364. Find the width of the gravel path.

Solution :



Given, length of the square field = 10m

Let the side of the flower bed = x m

$$\therefore \text{Area of square field} = 100 \text{ m}^2 \text{ \& area of the flower bed} = x^2 \text{ m}^2$$

$$\therefore \text{Area of the gravel path} = (100 - x^2) \text{ m}^2$$

Given cost of laying flower bed = ₹3 /m².

Cost of laying gravel path = ₹4 /m².

\therefore By the problem,

$$3x^2 + 4(100 - x^2) = 364$$

$$\Rightarrow 3x^2 + 400 - 4x^2 = 364$$

$$\Rightarrow x^2 = 36$$

$$\Rightarrow x = 6$$

\therefore Length of flower bed = 6m

$$\therefore \text{Width of the path} = \frac{10 - 6}{2}$$

$$= \frac{4}{2}$$

$$= 2\text{m}$$

9. Two women together took 100 eggs to a market, one had more than the other. Both sold them for the same sum of money. The first then said to the second: "If I had your eggs, I would have earned ₹ 15", to which the second replied: "If I had your eggs, I would have earned ₹ $6\frac{2}{3}$ ". How many eggs did each had in the beginning?

Solution :

Let the number of eggs of woman 1 and 2 respectively be x, y , and their selling price be

$$p, q, \quad \Rightarrow x + y = 100 \quad \text{--- (1)}$$

* If both of them sold the eggs for the equal sum of money, $px = qy$.

* By the data given in the problem,

$$py = 15, \quad qx = 6\frac{2}{3}$$

$$qx = \frac{20}{3}$$

$$p = \frac{15}{y}, \quad q = \frac{20}{3x}$$

Also,

$$px = qy \Rightarrow \frac{15}{y}x = \frac{20}{3x}y$$

$$\Rightarrow \frac{3x}{y} = \frac{4y}{3x}$$

$$\Rightarrow 9x^2 = 4y^2$$

$$\Rightarrow 9x^2 = 4(100 - x)^2 \quad (\text{from (1)})$$

$$\Rightarrow 9x^2 = 4(x^2 - 200x + 10000)$$

$$\Rightarrow 5x^2 + 800x - 40000 = 0$$

$$\Rightarrow x^2 + 160x - 8000 = 0$$

$$\Rightarrow (x + 200)(x - 40) = 0$$

$$\Rightarrow x = 40$$

$$\therefore y = 60$$

\therefore Woman 1 had 40 eggs and
Woman 2 had 60 eggs.

10. The hypotenuse of a right angled triangle is 25 cm and its perimeter 56 cm. Find the length of the smallest side.

Solution :

$$\text{Given } b = 25 \text{ cm, } a + b + c = 56 \text{ cm}$$

$$\Rightarrow a + c = 56 - 25$$

$$\Rightarrow a + c = 31$$

$$\text{Let } a = x, c = 31 - x$$

$$\therefore \text{ In } \triangle ABC, \quad a^2 + c^2 = 25^2$$

$$x^2 + (31 - x)^2 = 625$$

$$2x^2 - 62x + 336 = 0$$

$$x^2 - 31x + 168 = 0$$

$$(x - 24)(x - 7) = 0$$

$$x = 24, 7$$

\therefore Length of smallest side = 7 cm

XIII. NATURE OF ROOTS OF A QUADRATIC EQUATION :

Key Points

✓ The roots of the quadratic equation $ax^2 + bx + c = 0$, $a \neq 0$ are found using the formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Values of Discriminant $\Delta = b^2 - 4ac$

$$\Delta > 0$$

$$\Delta = 0$$

$$\Delta < 0$$

Nature of Roots

Real and Unequal roots

Real and Equal roots

No Real root

Example 3.41

Determine the nature of roots for the following quadratic equations

i) $x^2 - x - 20 = 0$

ii) $9x^2 - 24x + 16 = 0$

iii) $2x^2 - 2x + 9 = 0$

Solution :

i) $x^2 - x - 20 = 0$

Here, $a = 1$, $b = -1$, $c = -20$

Now, $\Delta = b^2 - 4ac$

$$\Delta = (-1)^2 - 4(1)(-20) = 81$$

Here, $\Delta = 81 > 0$. So, the equation will have real and unequal roots

ii) $9x^2 - 24x + 16 = 0$

Here, $a = 9$, $b = -24$, $c = 16$

Now, $\Delta = b^2 - 4ac$

$$\Delta = (-24)^2 - 4(9)(16) = 0$$

Here, $\Delta = 0$. So, the equation will have real and unequal roots

iii) $2x^2 - 2x + 9 = 0$

Here, $a = 2$, $b = -2$, $c = 9$

Now, $\Delta = b^2 - 4ac$

$$\Delta = (-2)^2 - 4(2)(9) = -68$$

Here, $\Delta = -68 < 0$. So, the equation will have no real roots

Example 3.42

(i) Find the values of 'k', for which the quadratic equation $kx^2 - (8k + 4)x + 81 = 0$ has real and equal roots ?

equation $(k + 9)x^2 + (k + 1)x + 1 = 0$ has no real roots ?

Solution :

i) $kx^2 - (8k + 4)x + 81 = 0$

Since the equation has real and equal roots, $\Delta = 0$.

That is, $b^2 - 4ac = 0$

Here, $a = k$, $b = -(8k + 4)$, $c = 81$

That is, $[(8k + 4)]^2 - 4(k)(81) = 0$

$$64k^2 + 64k + 16 - 324k = 0$$

$$64k^2 - 260k + 16 = 0$$

dividing by 4 we get $16k^2 - 65k + 4 = 0$

$$(16k - 1)(k - 4) = 0 \text{ then, } k = \frac{1}{16} \text{ or } k = 4$$

ii) $(k + 9)x^2 + (k + 1)x + 1 = 0$

Since the equation has no real roots, $\Delta < 0$

That is, $b^2 - 4ac < 0$

Here, $a = k + 9$, $b = k + 1$, $c = 1$

That is, $(k + 1)^2 - 4(k + 9)(1) < 0$

$$k^2 + 2k + 1 - 4k - 36 < 0$$

$$k^2 - 2k - 35 < 0$$

$$(k + 5)(k - 7) < 0$$

Therefore $-5 < k < 7$. {If $\alpha < \beta$ and if $(x - \alpha)(x - \beta) < 0$ then, $\alpha < x < \beta$).

Example 3.43

Prove that the equation $x^2(p^2 + q^2) + 2x(pr + qs) + r^2 + s^2 = 0$ has no real roots. If $ps = qr$, then show that the roots are real and equal.

Solution :

The given quadratic equation is,

$$x^2(p^2 + q^2) + 2x(pr + qs) + r^2 + s^2 = 0$$

Here, $a = p^2 + q^2$, $b = 2(pr + qs)$, $c = r^2 + s^2$

Now $\Delta = b^2 - 4ac$

$$= [2(pr + qs)]^2 - 4(p^2 + q^2)(r^2 + s^2)$$

$$= 4[p^2r^2 + 2pqrs + q^2s^2 - p^2r^2 - p^2s^2 - q^2r^2 - q^2s^2]$$

$$= 4[-p^2s^2 + 2pqrs - q^2r^2]$$

$$= -4[(ps - qr)^2] < 0 \quad \dots\dots (1)$$

Since, $\Delta = b^2 - 4ac < 0$, the roots are not equal.

If $ps = qr$ then $= -4[ps - qr]^2$

$$= -4[qr - qr]^2 = 0 \quad (\text{using (1)})$$

Thus $\Delta = 0$ if $ps = qr$ and so the roots will be real and equal.

EXERCISE 3.13

1. Determine the nature of the roots for the following quadratic equations

i) $15x^2 + 11x + 2 = 0$

ii) $x^2 - x + 1 = 0$

iii) $\sqrt{2}t^2 - 3t + 3\sqrt{2} = 0$

iv) $9y^2 - 6\sqrt{2}y + 2 = 0$

v) $9a^2b^2x^2 - 24abcdx + 16c^2d^2 = 0$, $a \neq 0$, $b \neq 0$

Solution :

i) Given equation is $15x^2 + 11x + 2 = 0$

$$a = 15, b = 11, c = 2$$

$$\therefore \Delta = b^2 - 4ac$$

$$= 121 - 4 \times 15 \times 2$$

$$= 121 - 120$$

\therefore The equation will have real and unequal roots.

ii) Given equation is $x^2 - x - 1 = 0$

$$a = 1, b = -1, c = -1$$

$$\therefore \Delta = b^2 - 4ac$$

$$= 1 - 4(1)(-1)$$

$$= 1 + 4$$

$$= 5 > 0$$

\therefore The equation will have real and unequal roots.

iii) Given $\sqrt{2}t^2 - 3t + 3\sqrt{2} = 0$

$$a = \sqrt{2}, b = -3, c = 3\sqrt{2}$$

$$\therefore \Delta = b^2 - 4ac$$

$$= 9 - 4(\sqrt{2})(3\sqrt{2})$$

$$= 9 - 24$$

$$= -15 < 0$$

\therefore The roots are unreal.

iv) Given $9y^2 - 6\sqrt{2}y + 2 = 0$

$$a = 9, b = -6\sqrt{2}, c = 2$$

$$\therefore \Delta = b^2 - 4ac$$

$$= (-6\sqrt{2})^2 - 4(9)(2)$$

$$= 72 - 72$$

$$= 0$$

\therefore The roots are real & equal.

v) Given $9a^2b^2x^2 - 24abcdx + 16c^2d^2$

$$= 0, a, b \neq 0$$

$$a = 9a^2b^2, b = -24abcd, c = 16c^2d^2$$

$$\therefore \Delta = B^2 - 4AC$$

$$= 576 a^2b^2c^2d^2 - 4(9a^2b^2)(16c^2d^2)$$

$$= 576 a^2b^2c^2d^2 - 576 a^2b^2c^2d^2$$

$$= 0$$

\therefore The roots are real & equal.

2. Find the value(s) of 'k' for which the roots of the following equations are real and equal.

i) $(5k - 6)x^2 + 2kx + 1 = 0$

ii) $kx^2 + (6k + 2)x + 16 = 0$

Solution :

i) Given equation is $(5k - 6)x^2 + 2kx + 1 = 0$ are real & equal

$$a = 5k - 6, b = 2k, c = 1$$

$$\therefore \Delta = b^2 - 4ac = 0$$

$$\Rightarrow 4k^2 - 4(5k - 6)(1) = 0$$

$$\Rightarrow 4k^2 - 20k + 24 = 0$$

$$\Rightarrow k^2 - 5k + 6 = 0$$

$$\Rightarrow (k - 3)(k - 2) = 0$$

$$\therefore k = 3, 2$$

ii) Given the roots of $kx^2 + (6k + 2)x + 16 = 0$ are real & equal

$$a = k, b = 6k + 2, c = 16$$

$$\therefore \Delta = b^2 - 4ac = 0$$

- 10	9
$\frac{-9}{9}$	$\frac{-1}{9}$
- 1	$\frac{-1}{9}$

$$\Rightarrow (6k + 2)^2 - 4(k)(16) = 0$$

$$\Rightarrow 36k^2 + 24k + 4 - 64k = 0$$

$$\Rightarrow 36k^2 - 40k + 4 = 0$$

$$\Rightarrow 9k^2 - 10k + 1 = 0$$

$$\therefore k = 1, \frac{1}{9}$$

3. If the roots of $(a - b)x^2 + (b - c)x + (c - a) = 0$ are real and equal, then prove that

Solution :

i) Given equation is $(a - b)x^2 + (b - c)x + (c - a) = 0$ are real & equal

To prove : b, a, c are in A.P.

Here $A = a - b, B = b - c, C = c - a$

$$\therefore \Delta = B^2 - 4AC = 0$$

$$\Rightarrow (b - c)^2 - 4(a - b)(c - a) = 0$$

$$\Rightarrow (b^2 + c^2 - 2bc) - 4(ac - bc - a^2 + ab) = 0$$

$$\Rightarrow b^2 + c^2 - 2bc - 4ac + 4bc + 4a^2 - 4ab = 0$$

$$\Rightarrow 4a^2 + b^2 + c^2 - 4ab + 2bc - 4ac = 0$$

$$\Rightarrow (2a - b - c)^2 = 0$$

$$\Rightarrow 2a - b - c = 0$$

$$\Rightarrow b + c = 2a$$

$$\Rightarrow a = \frac{b + c}{2}$$

\therefore b, a, c are in A.P.

4. If a, b are real then show that the roots of the equation $(a-b)x^2 - 6(a+b)x - 9(a-b) = 0$ are real and unequal.

Solution :

To prove the roots of $(a-b)x^2 - 6(a+b)x - 9(a-b) = 0$ are real & unequal

Here $A = a - b$, $B = -6(a+b)$, $C = -9(a-b)$

$$\therefore \Delta = B^2 - 4AC = 0$$

$$= 36(a+b)^2 - 4(a-b)(-9(a-b)) = 0$$

$$= 36(a+b)^2 + 36(a-b)^2$$

$$= 36[(a+b)^2 + (a-b)^2]$$

$$= 36[2(a^2 + b^2)]$$

$$= 72(a^2 + b^2) > 0, \quad a \text{ \& \& } b \text{ are real}$$

\therefore The roots of the equation are real & unequal.

5. If the roots of the equation $(c^2 - ab)x^2 - 2(a^2 - bc)x + b^2 - ac = 0$ are real and equal prove that either $a = 0$ (or) $a^3 + b^3 + c^3 = 3abc$.

Solution :

Given roots of the equation $(c^2 - ab)x^2 - 2(a^2 - bc)x + b^2 - ac = 0$ are real & equal

To prove either $a = 0$ (or)

$$a^3 + b^3 + c^3 = 3abc$$

Here $A = c^2 - ab$, $B = -2(a^2 - bc)$, $C = b^2 - ac$

$$\text{Given } \Delta = B^2 - 4AC = 0$$

$$\Rightarrow 4(a^2 - bc)^2 - 4(c^2 - ab)(b^2 - ac) = 0$$

$$\Rightarrow (a^2 - bc)^2 - (c^2 - ab)(b^2 - ac) = 0$$

$$\Rightarrow (a^4 + b^2c^2 - 2a^2bc) - (b^2c^2 - ab^3 - ac^3 + a^2bc) = 0$$

$$\Rightarrow a^4 - 3a^2bc + ab^3 + ac^3 = 0$$

$$\Rightarrow a = 0 \text{ (or) } a^3 + b^3 + c^3 - 3abc = 0$$

$$\Rightarrow a^3 + b^3 + c^3 = 3abc$$

Hence proved.

XIV. RELATION BETWEEN ROOTS AND COEFFICIENTS OF A QUADRATIC EQUATION

Key Points

✓ Let α and β are the roots of the equation $ax^2 + bx + c = 0$ then.

$$\alpha + \beta = \frac{-b}{a} = \frac{-\text{Co-efficient of } x}{\text{Co-efficient of } x^2}$$

$$\alpha\beta = \frac{c}{a} = \frac{\text{Constant term}}{\text{Co-efficient of } x^2}$$

Example 3.44

If the difference between the roots of the equation $x^2 - 13x + k = 0$ is 17 find k .

Solution :

$$x^2 - 13x + k = 0 \text{ here, } a = 1, b = -13, c = k$$

Let α, β be the roots of the equation. Then

$$\alpha + \beta = \frac{-b}{a} = \frac{-(-13)}{1} = 13 \quad \dots\dots\dots (1)$$

$$\text{Also } \alpha - \beta = 17 \quad \dots\dots\dots (2)$$

$$(1) + (2) \text{ we get, } 2\alpha = 30 \text{ gives } \alpha = 15$$

$$\text{Therefore, } 15 + \beta = 13 \text{ (from (1)) gives } \beta = -2$$

$$\text{But } \alpha\beta = \frac{c}{a} = \frac{k}{1} \text{ gives } 15 \times (-2) = k$$

$$\text{we get, } k = -30$$

Example 3.45

If α and β are the roots of $x^2 + 7x + 10 = 0$ find the values of

$$i) (\alpha - \beta) \quad ii) \alpha^2 + \beta^2 \quad iii) \alpha^3 - \beta^3 \quad iv) \alpha^4 + \beta^4$$

$$v) \frac{\alpha}{\beta} + \frac{\beta}{\alpha} \quad vi) \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha}$$

Solution :

$$x^2 + 7x + 10 = 0 \text{ here, } a = 1, b = 7, c = 10$$

If α and β be the roots of the equation then,

$$\alpha + \beta = \frac{-b}{a} = \frac{-7}{1} = -7; \alpha\beta = \frac{c}{a} = \frac{10}{1} = 10$$

$$\begin{aligned} i) \alpha - \beta &= \sqrt{(\alpha + \beta)^2 - 4\alpha\beta} \\ &= \sqrt{(-7)^2 - 4 \times 10} = \sqrt{9} = 3 \end{aligned}$$

$$\begin{aligned} ii) \alpha^2 + \beta^2 &= (\alpha + \beta)^2 - 2\alpha\beta \\ &= (-7)^2 - 2 \times 10 = 29 \end{aligned}$$

$$\begin{aligned} iii) \alpha^3 + \beta^3 &= (\alpha + \beta)^3 + 3\alpha\beta(\alpha + \beta) \\ &= (3)^3 + 3(10)(3) = 117 \end{aligned}$$

$$\begin{aligned} iv) \alpha^4 + \beta^4 &= (\alpha^2 + \beta^2)^2 - 2\alpha^2\beta^2 \\ &= 29^2 - 2 \times (10)^2 \\ &= 641 \text{ (since from (iii), } \alpha^2 + \beta^2 = 29) \end{aligned}$$

$$\begin{aligned} v) \frac{\alpha}{\beta} + \frac{\beta}{\alpha} &= \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} \\ &= \frac{49 - 20}{10} = \frac{29}{10} \end{aligned}$$

$$\begin{aligned} vi) \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} &= \frac{\alpha^3 + \beta^3}{\alpha\beta} = \frac{(\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)}{\alpha\beta} \\ &= \frac{(-343) - 3(10 \times (-7))}{10} \\ &= \frac{-343 + 210}{10} = \frac{-133}{10} \end{aligned}$$

Example 3.46

If α, β are the roots of the equation $3x^2 + 7x - 2 = 0$, find the values of

$$i) \frac{\alpha}{\beta} + \frac{\beta}{\alpha} \quad ii) \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha}$$

Solution :

$$3x^2 + 7x - 2 = 0 \text{ here, } a = 3, b = 7, c = -2$$

since α, β are the roots of the equation

$$\begin{aligned} i) \alpha + \beta &= \frac{-b}{a} = \frac{-7}{3}; \alpha\beta = \frac{c}{a} = \frac{-2}{3} \\ \frac{\alpha}{\beta} + \frac{\beta}{\alpha} &= \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} \\ &= \frac{\left(\frac{-7}{3}\right)^2 - 2\left(\frac{-2}{3}\right)}{\frac{-2}{3}} \\ &= \frac{-61}{6} \end{aligned}$$

$$\begin{aligned}
 \text{ii) } \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} &= \frac{\alpha^3 + \beta^3}{\alpha\beta} = \frac{(\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)}{\alpha\beta} \\
 &= \frac{\left(-\frac{7}{3}\right)^3 - 3\left(-\frac{2}{3}\right)\left(-\frac{7}{3}\right)}{-\frac{7}{3}} \\
 &= \frac{67}{9}
 \end{aligned}$$

Example 3.47

If α, β are the roots of the equation $2x^2 - x - 1 = 0$, then form the equation whose roots are

i) $\frac{1}{\alpha}, \frac{1}{\beta}$ ii) $\alpha^2\beta, \beta^2\alpha$ iii) $2\alpha + \beta, 2\beta + \alpha$

Solution :

$$2x^2 - x - 1 = 0 \text{ here, } a = 2, b = -1, c = -1$$

$$\alpha + \beta = \frac{-b}{a} = \frac{-(-1)}{2} = \frac{1}{2}; \alpha\beta = \frac{c}{a} = -\frac{1}{2}$$

i) Given roots are $\frac{1}{\alpha}, \frac{1}{\beta}$

$$\text{Sum of the roots} = \frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta} = \frac{\frac{1}{2}}{-\frac{1}{2}} = -1$$

$$\text{Product of the roots} = \frac{1}{\alpha} \times \frac{1}{\beta} = \frac{1}{\alpha\beta} = \frac{1}{-\frac{1}{2}} = -2$$

The required equation is $x^2 - (\text{Sum of the roots})x + (\text{Product of the roots}) = 0$

$$x^2 - (-1)x - 2 = 0 \text{ gives } x^2 + x - 2 = 0$$

ii) Given roots are $\alpha^2\beta, \beta^2\alpha$

$$\begin{aligned}
 \text{Sum of the roots } \alpha^2\beta, \beta^2\alpha \\
 = \alpha\beta(\alpha + \beta) = -\frac{1}{2}\left(\frac{1}{2}\right) = -\frac{1}{4}
 \end{aligned}$$

Product of the roots $(\alpha^2\beta) \times (\beta^2\alpha)$

$$= \alpha^3\beta^3 = (\alpha\beta)^3 = \left(-\frac{1}{2}\right)^3 = -\frac{1}{8}$$

The required equation is $x^2 - (\text{Sum of the roots})x + (\text{Product of the roots}) = 0$

$$x^2 - \left(-\frac{1}{4}\right)x - \frac{1}{8} = 0 \text{ gives } 8x^2 + 2x - 1 = 0$$

iii) $2\alpha + \beta, 2\beta + \alpha$

Sum of the roots $2\alpha + \beta + 2\beta + \alpha$

$$= 3(\alpha + \beta) = 3\left(\frac{1}{2}\right) = \frac{3}{2}$$

Product of the roots $= (2\alpha + \beta)(2\beta + \alpha)$

$$= 4\alpha\beta + 2\alpha^2 + 2\beta^2 = \alpha\beta$$

$$= 5\alpha\beta + 2[(\alpha + \beta)^2 - 2\alpha\beta]$$

$$= 5\left(-\frac{1}{2}\right) + 2\left[\frac{1}{4} - 2 \times -\frac{1}{2}\right] = 0$$

The required equation is $x^2 - (\text{Sum of the roots})x + (\text{Product of the roots}) = 0$

$$x^2 - \frac{3}{2}x + 0 = 0 \text{ gives } 2x^2 - 3x = 0$$

EXERCISE 3.14

1. Write each of the following expression in terms of $\alpha + \beta$ and $\alpha\beta$.

$$i) \frac{\alpha}{3\beta} + \frac{\beta}{3\alpha}$$

$$ii) \frac{1}{\alpha^2\beta} + \frac{1}{\beta^2\alpha}$$

$$iii) (3\alpha - 1)(3\beta - 1) \quad iv) \frac{\alpha + 3}{\beta} + \frac{\beta + 3}{\alpha}$$

Solution :

$$\begin{aligned} i) \quad & \frac{\alpha}{3\beta} + \frac{\beta}{3\alpha} \\ &= \frac{\alpha^2 + \beta^2}{3\alpha\beta} \\ &= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{3\alpha\beta} \end{aligned}$$

$$\begin{aligned} ii) \quad & \frac{1}{\alpha^2\beta} + \frac{1}{\beta^2\alpha} \\ &= \frac{\beta + \alpha}{\alpha^2\beta^2} \\ &= \frac{\alpha + \beta}{(\alpha\beta)^2} \end{aligned}$$

$$\begin{aligned} iii) \quad & (3\alpha - 1)(3\beta - 1) \\ &= 9\alpha\beta - 3\alpha - 3\beta + 1 \\ &= 9\alpha\beta - 3(\alpha + \beta) + 1 \end{aligned}$$

$$\begin{aligned} iv) \quad & \frac{\alpha + 3}{\beta} + \frac{\beta + 3}{\alpha} \\ &= \frac{\alpha^2 + 3\alpha + \beta^2 + 3\beta}{\alpha\beta} \\ &= \frac{\alpha^2 + \beta^2 + 3(\alpha + \beta)}{\alpha\beta} \\ &= \frac{(\alpha + \beta)^2 - 2\alpha\beta + 3(\alpha + \beta)}{\alpha\beta} \end{aligned}$$

2. The roots of the equation $2x^2 - 7x + 5 = 0$ are α and β . Without solving for the roots, find

$$i) \frac{1}{\alpha} + \frac{1}{\beta} \quad ii) \frac{\alpha}{\beta} + \frac{\beta}{\alpha} \quad iii) \frac{\alpha + 2}{\beta + 2} + \frac{\beta + 2}{\alpha + 2}$$

Solution :

Given α, β are the roots of $2x^2 - 7x + 5 = 0$

$$a = 2, b = -7, c = 5$$

$$\alpha + \beta = \frac{-b}{a} = \frac{7}{2}$$

$$\alpha\beta = \frac{c}{a} = \frac{5}{2}$$

$$\begin{aligned} i) \quad & \frac{1}{\alpha} + \frac{1}{\beta} \\ &= \frac{\alpha + \beta}{\alpha\beta} \end{aligned}$$

$$\begin{aligned} &= \frac{\frac{7}{2}}{\frac{5}{2}} \\ &= \frac{7}{5} \end{aligned}$$

$$\begin{aligned} ii) \quad & \frac{\alpha}{\beta} + \frac{\beta}{\alpha} \\ &= \frac{\alpha^2 + \beta^2}{\alpha\beta} \\ &= \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} \end{aligned}$$

$$\begin{aligned} &= \frac{\frac{49}{4} - 5}{\frac{5}{2}} = \frac{29}{4} \times \frac{2}{5} = \frac{29}{10} \end{aligned}$$

$$\begin{aligned}
 \text{iii) } & \frac{\alpha+2}{\beta+2} + \frac{\beta+2}{\alpha+2} \\
 &= \frac{\alpha^2 + 4\alpha + 4 + \beta^2 + 4\beta + 4}{(\alpha+2)(\beta+2)} \\
 &= \frac{(\alpha^2 + \beta^2) + 4(\alpha + \beta) + 8}{\alpha\beta + 2\alpha + 2\beta + 4} \\
 &= \frac{(\alpha + \beta)^2 - 2\alpha\beta + 4(\alpha + \beta) + 8}{\alpha\beta + 2(\alpha + \beta) + 4} \\
 &= \frac{\frac{49}{4} - 5 + 2\left(\frac{7}{2}\right) + 8}{\frac{5}{2} + 2\left(\frac{7}{2}\right) + 4} \\
 &= \frac{\frac{49}{4} + 3 + 14}{\frac{5}{2} + 4} \\
 &= \frac{\frac{49 + 68}{4}}{\frac{13}{2}} \\
 &= \frac{117}{4} \times \frac{2}{13} \\
 &= \frac{13}{6}
 \end{aligned}$$

3. The roots of the equation $x^2 + 6x - 4 = 0$ are α, β . Find the quadratic equation whose roots are

i) α^2 and β^2 ii) $\frac{2}{\alpha}$ and $\frac{2}{\beta}$ iii) $\alpha^2\beta$ and $\beta^2\alpha$

Solution :

Given α, β are the roots of $x^2 + 6x - 4 = 0$

$$a = 1, b = 6, c = -4$$

$$\alpha + \beta = \frac{-b}{a} = -6$$

$$\alpha\beta = \frac{c}{a} = -4$$

i) To find the equation whose roots are α^2, β^2

$$\begin{aligned}
 \text{Sum} &= \alpha^2 + \beta^2 \\
 &= (\alpha + \beta)^2 - 2\alpha\beta \\
 &= (-6)^2 - 2(-4) \\
 &= 36 + 8 \\
 &= 44
 \end{aligned}$$

$$\begin{aligned}
 \text{Product} &= \alpha^2 \beta^2 \\
 &= (\alpha\beta)^2 \\
 &= (-4)^2 \\
 &= 16
 \end{aligned}$$

\therefore The required equation is

$$x^2 - (\text{Sum of the roots})x + \text{Product of the roots} = 0$$

$$\Rightarrow x^2 - 44x + 16 = 0$$

ii) To find the equation whose roots are

$$\begin{aligned}
 &\frac{2}{\alpha}, \frac{2}{\beta} \\
 \text{Sum} &= \frac{2}{\alpha} + \frac{2}{\beta} \\
 &= 2\left(\frac{1}{\alpha} + \frac{1}{\beta}\right) \\
 &= 2\left(\frac{\alpha + \beta}{\alpha\beta}\right) = 2\left(\frac{-6}{-4}\right) = 3
 \end{aligned}$$

$$\begin{aligned}
 \text{Product} &= \frac{2}{\alpha} \cdot \frac{2}{\beta} \\
 &= \frac{4}{\alpha\beta} \\
 &= \frac{4}{-4} = -1
 \end{aligned}$$

\therefore The required equation is $x^2 - 3x - 1 = 0$

iii) To find the equation whose roots are $\alpha^2\beta$, $\beta^2\alpha$

$$\begin{aligned}\text{Sum} &= \alpha^2\beta + \beta^2\alpha \\ &= \alpha\beta(\alpha + \beta) \\ &= -4(-6) \\ &= 24\end{aligned}$$

$$\begin{aligned}\text{Product} &= \alpha^2\beta \cdot \alpha\beta^2 \\ &= (\alpha\beta)^3 \\ &= (-4)^3 \\ &= -64\end{aligned}$$

$$\begin{aligned}\therefore \text{The required equation is} \\ \Rightarrow x^2 - 24x - 64 = 0\end{aligned}$$

4. If α, β are the roots of $7x^2 + ax + 2 = 0$ and if $\beta - \alpha = \frac{-13}{7}$ Find the values of a .

Solution :

Given α, β are the roots of $7x^2 + ax + 2 = 0$

$$\alpha + \beta = -\frac{a}{7}$$

$$\alpha\beta = \frac{2}{7}$$

$$\text{Also, } \beta - \alpha = \frac{-13}{7}$$

$$\Rightarrow \alpha - \beta = \frac{13}{7}$$

$$\Rightarrow (\alpha - \beta)^2 = \frac{169}{49}$$

$$\Rightarrow (\alpha + \beta)^2 - 4\alpha\beta = \frac{169}{49}$$

$$\Rightarrow \left(-\frac{a}{7}\right)^2 - 4\left(\frac{2}{7}\right) = \frac{169}{49}$$

$$\Rightarrow \frac{a^2}{49} - \frac{8}{7} = \frac{169}{49}$$

$$\Rightarrow \frac{a^2}{49} - \frac{56}{49} = \frac{169}{49}$$

$$\Rightarrow a^2 - 56 = 169$$

$$\Rightarrow a^2 = 225$$

$$\therefore a = 15, -15$$

5. If one root of the equation $2y^2 - ay + 64 = 0$ is twice the other then find the values of a .

Solution :

Let α, β be the roots of $2y^2 - ay + 64 = 0$

$$\alpha + \beta = \frac{a}{2}$$

$$\alpha\beta = 32$$

Given $\alpha = 2\beta$

$$\begin{array}{l|l}\therefore \alpha + \beta = \frac{a}{2} & \alpha\beta = 32 \\ \Rightarrow 3\beta = \frac{a}{2} & 2\beta \cdot \beta = 32 \\ \Rightarrow \beta = \frac{a}{6} & \Rightarrow \beta^2 = 16 \\ & \Rightarrow \beta = \pm 4 \\ & \therefore \frac{a}{6} = \pm 4\end{array}$$

$$a = \pm 24$$

6. If one root of the equation $3x^2 + kx + 81 = 0$ (having real roots) is the square of the other then find k .

Solution :

Let α, β be the roots of $3x^2 + kx + 81 = 0$

$$\alpha + \beta = -\frac{k}{3}$$

$$\alpha\beta = 27$$

Given $\alpha = \beta^2$

$$\begin{array}{l|l}\therefore \alpha + \beta = -\frac{k}{3} \quad \dots\dots\dots (1) & \alpha\beta = 27 \\ & \Rightarrow \beta^2 \cdot \beta = 27 \\ & \Rightarrow \beta^3 = 27 \\ & \therefore \beta = 3 \\ & \therefore \alpha = 9\end{array}$$

$$\therefore (1) \Rightarrow 9 + 3 = -\frac{k}{3}$$

$$\Rightarrow -\frac{k}{3} = 12$$

$$\Rightarrow k = -36$$

XV. QUADRATIC GRAPHS

Key Points

- ✓ A parabola represents a Quadratic function.
- ✓ A quadratic function has the form $f(x) = ax^2 + bx + c$, where a, b, c are constants, and $a \neq 0$.
- ✓ The coefficient a in the general equation is responsible for parabolas to open upward or downward and vary in “width” (“wider” or “skinnier”), but they all have the same basic “ \cup ” shape.
- ✓ The greater the quadratic coefficient, the narrower is the parabola.
- ✓ The lesser the quadratic coefficient, the wider is the parabola.
- ✓ A parabola is symmetric with respect to a line called the axis of symmetry. The point of intersection of the parabola and the axis of symmetry is called the vertex of the parabola. The graph of any second degree polynomial gives a curve called “parabola”.
- ✓ If the graph of the given quadratic equation intersect the X axis at two distinct points, then the given equation has two real and unequal roots.
- ✓ If the graph of the given quadratic equation touch the X axis at only one point, then the given
- ✓ If the graph of the given equation does not intersect the X axis at any point then the given equation has no real root.
- ✓ If the straight line intersects the parabola at two distinct points, then the x coordinates of those points will be the roots of the given quadratic equation.
- ✓ If the straight line just touch the parabola at only one point, then the x coordinate of the common point will be the single root of the quadratic equation.
- ✓ If the straight line doesn't intersect or touch the parabola then the quadratic equation will have no real roots.

Example 3.48

Discuss the nature of solutions of the following quadratic equations.

i) $x^2 + x - 12 = 0$

ii) $x^2 - 8x + 16 = 0$

iii) $x^2 + 2x + 5 = 0$


Solution :

i) $x^2 + x - 12 = 0$

Step 1 Prepare the table of values for the equation $y = x^2 + x - 12$

x	-5	-4	-3	-2	-1	0	1	2	3	4
y	8	0	-6	-10	-12	-12	-10	-6	0	8

Step 2 Plot the points for the above ordered pairs (x, y) on the graph using suitable scale.



Step 3 Draw the parabola and mark the co-ordinates of the parabola which intersect the X axis.

Step 4 The roots of the equation are the x coordinates of the intersecting points $(-4, 0)$ and $(3, 0)$ of the parabola with the X axis which are -4 and 3 respectively.


Since there are two points of intersection with the X axis, the quadratic equation $x^2 + x - 12$ has real and unequal roots.

ii) $x^2 - 8x + 16 = 0$

Step 1 Prepare the table of values for the equation $y = x^2 - 8x + 16$

x	-1	0	1	2	3	4	5	6	7	8
y	25	16	9	4	1	0	1	4	9	16

Step 2 Plot the points for the above ordered pairs (x, y) on the graph using suitable scale.



Step 3 Draw the parabola and mark the co-ordinates of the parabola which intersect with the X axis.

Step 4 The roots of the equation are the x coordinates of the intersecting points of the parabola with the X axis (4, 0) which is 4.

Since there is only one point of intersection with the X axis, the quadratic equation $x^2 - 8x + 16 = 0$ has real and equal roots.


iii) $x^2 + 2x + 5 = 0$

Let $y = x^2 + 2x + 5$

Step 1 Prepare the table of values for the equation $y = x^2 + 2x + 5$

x	-3	-2	-1	0	1	2	3
y	8	5	4	5	8	13	20

Step 2 Plot the above ordered pairs (x, y) on the graph using suitable scale.



Step 3 Join the points by a free-hand smooth curve this smooth curve is the graph of $y = x^2 + 2x + 5$

Step 4 The solutions of the given quadratic equation are the x coordinates of the intersecting points of the parabola the X axis.

Here the parabola doesn't intersect or touch the X axis.

So, we conclude that there is no real root for the given quadratic equation.

Example 3.49

Draw the graph of $y = 2x^2$ and hence solve $2x^2 - x - 6 = 0$

Solution :

Step 1 Draw the graph of $y = 2x^2$ by preparing the table of values as below

x	-2	-1	0	1	2
y	8	2	0	2	8


Step 2 To solve $2x^2 - x - 6 = 0$, subtract $2x^2 - x - 6 = 0$ from $y = 2x^2$

$$\text{that is } y = 2x^2 \quad (-)$$

$$0 = 2x^2 - x - 6$$

$$\underline{y = x + 6}$$

The equation $y = x + 6$ represents a straight line. Draw the graph of $y = x + 6$ by forming table of values as below



x	-2	-1	0	1	2
y	4	5	6	7	8

Step 3 Mark the points of intersection of the curve $y = 2x^2$ and the line $y = x + 6$. That is, $(-1.5, 4.5)$ and $(2, 8)$

Step 4 The x coordinates of the respective points forms the solution set $\{-1.5, 2\}$ for $2x^2 - x - 6 = 0$


Example 3.50

Draw the graph of $y = x^2 + 4x + 3$ and hence find the roots of $x^2 + x + 1 = 0$

Solution :

Step 1 Draw the graph of $y = x^2 + 4x + 3$ by preparing the table of values as below

x	-4	-3	-2	-1	0	1	2
y	3	0	-1	0	3	8	15



Step 2 To solve $x^2 + x + 1 = 0$, subtract $x^2 + x + 1 = 0$ from $y = x^2 + 4x + 3$

$$\begin{array}{rcl}
 \text{that is} & y = x^2 + 4x + 3 & (-) \\
 & 0 = x^2 + x + 1 & \\
 \hline
 & y = & 3x + 2
 \end{array}$$

The equation represent a straight line. Draw the graph of $y = 3x + 2$ by forming the table of values as below.

x	-2	-1	0	1	2
y	-4	-1	2	5	8

Step 3 Observe that the graph of $y = 3x + 2$ does not intersect or touch the graph of the parabola $y = x^2 + 4x + 3$

Thus $x^2 + x + 1 = 0$ has no real roots.


Example 3.51

Draw the graph of $y = x^2 + x - 2$ and hence solve $x^2 + x - 2 = 0$

Solution :

Step 1 Draw the graph of $y = x^2 + x - 2$ by preparing the table of values as below

x	-3	-2	-1	0	1	2
y	4	0	-2	-2	0	4



Step 2 To solve $x^2 + x - 2 = 0$, subtract $x^2 + x - 2 = 0$ from $y = x^2 + x - 2$

$$\begin{array}{rcl}
 \text{that is} & y = x^2 + x - 2 & (-) \\
 & 0 = x^2 + x + 2 & \\
 \hline
 & y = & 0
 \end{array}$$

The equation $y = 0$ represents the X axis.

Step 3 Mark the point of intersection of the curve $x^2 + x - 2$ with the X axis. That is $(-2, 0)$ and $(1, 0)$

Step 4 The x coordinates of the respective points form the solution set $\{-2, 1\}$ for $x^2 + x - 2 = 0$.


Example 3.52

Draw the graph of $y = x^2 - 4x + 3$ and use it to solve $x^2 - 6x + 9 = 0$

Solution :

Step 1 Draw the graph of $y = x^2 - 4x + 3$ by preparing the table of values as below

x	-2	-1	0	1	2	3	4
y	15	8	3	0	-1	0	3



Step 2 To solve $x^2 - 6x + 9 = 0$, subtract $x^2 - 6x + 9 = 0$ from $y = x^2 - 4x + 3$

$$\begin{array}{rcl} \text{that is } & y = x^2 - 4x + 3 & (-) \\ & 0 = x^2 - 6x + 9 & \\ \hline & y = & 2x - 6 \end{array}$$

The equation $y = 2x - 6$ represents a straight line. Draw the graph of $y = 2x - 6$ forming the table of values as below.

x	0	1	2	3	4	5
y	-6	-4	-2	0	2	4

The line $y = 2x - 6$ intersect $y = x^2 - 4x + 3$ only at one point.

Step 3 Mark the point of intersection of the curve $y = x^2 - 4x + 3$ and $y = 2x - 6$ that is (3, 0)

Therefore, the x coordinate 3 is the only solution for the equation $x^2 - 6x + 9 = 0$.

EXERCISE 3.15

1. Graph the following quadratic equations and state their nature of solutions.

i) $x^2 - 9x + 20 = 0$

ii) $x^2 - 4x + 4 = 0$

iii) $x^2 + x + 7 = 0$

iv) $x^2 - 9 = 0$

v) $x^2 - 6x + 9 = 0$


vi) $(2x - 3)(x + 2) = 0$

Solution :

i) $x^2 - 9x + 20 = 0$

Let $y = x^2 - 9x + 20$

X :	-2	-1	0	1	2	3	4	5	6	7
x^2 :	4	1	0	1	4	9	16	25	36	49
$-9x$:	18	9	0	-9	-18	-27	-36	-45	-54	-63
20 :	20	20	20	20	20	20	20	20	20	20
$y = x^2 - 9x + 20$:	42	30	20	12	6	2	0	0	2	6




- Plot the points $(-1, 30)$, $(0, 20)$, $(1, 12)$, $(2, 6)$, $(3, 2)$, $(4, 0)$, $(5, 0)$, $(6, 2)$, $(7, 6)$ on the graph.
- Join all the points by a free-hand smooth curve. This curve is the graph of $y = x^2 - 9x + 20$.
- Here, the curve meets x -axis at $(4, 0)$, $(5, 0)$.
 \therefore The equation has real roots and x - coordinates of the points are $x = 4, x = 5$.
 \therefore Solution = $\{4, 5\}$

ii) $x^2 - 4x + 4 = 0$

Let $y = x^2 - 4x + 4$


$x :$	-2	-1	0	1	2	3	4
$x^2 :$	4	1	0	1	4	9	16
$-4x :$	8	4	0	-4	-8	-12	-16
$4 :$	4	4	4	4	4	4	4
$y = x^2 - 4x + 4 :$	16	9	4	1	0	1	4



- Plot the points $(-2, 16)$, $(-1, 9)$, $(0, 4)$, $(1, 1)$, $(2, 0)$, $(3, 1)$, $(4, 4)$ on the graph.
- Join all the points by a free-hand smooth curve. This curve is the graph of $y = x^2 - 4x + 4$.
- Here, the curve meets x -axis at $(2, 0)$.
 \therefore The equation has 2 equal roots.
 \therefore The x - coordinates of the points is $x = 2$.
 \therefore Solution = $\{2, 2\}$

Let $y = x^2 + x + 7$

$x :$	-3	-2	-1	0	1	2	3
$x^2 :$	9	4	1	0	1	4	9
$x :$	-3	-2	-1	0	1	2	3
$7 :$	7	7	7	7	7	7	7
$y = x^2 + x + 7 :$	13	9	7	7	9	13	19




- Plot the points $(-3, 13)$, $(-2, 9)$, $(-1, 7)$, $(0, 7)$, $(1, 9)$, $(2, 13)$, $(3, 19)$ on the graph.
- Join all the points by a free-hand smooth curve. This curve is the graph of $y = x^2 + x + 7$.
- Here the curve does not meet the x -axis and the curve has no real roots.

iv) $x^2 - 9 = 0$

Let $y = x^2 - 9$

$x :$	-4	-3	-2	-1	0	1	2	3	4
$x^2 :$	16	9	4	1	0	1	4	9	16
$-9 :$	-9	-9	-9	-9	-9	-9	-9	-9	-9
$y = x^2 - 9 :$	7	0	-5	-8	-9	-8	-5	0	7




- Plot the points $(-4, 7)$, $(-3, 0)$, $(-2, -5)$, $(-1, -8)$, $(0, -9)$, $(1, -8)$, $(2, -5)$, $(3, 0)$, $(4, 7)$ on the graph.
- Join all the points by a free-hand smooth curve. This curve is the graph of $y = x^2 - 9$.
- Here, the curve meets x-axis at 2 points $(-3, 0)$, $(3, 0)$
 - \therefore The equation has real and unequal roots.
 - \therefore The x - coordinates are 3, -3 will be the solution.
 - \therefore Solution = $\{-3, 3\}$

v) $x^2 - 6x + 9 = 0$

Solution:

Let $y = x^2 - 6x + 9$

$x :$	-2	-1	0	1	2	3	4	5
$x^2 :$	4	1	0	1	4	9	16	25
$-6x :$	12	6	0	-6	-12	-18	-24	-30
$9 :$	9	9	9	9	9	9	9	9
$y = x^2 - 6x + 9 :$	25	16	9	4	1	0	1	4




- Plot the points $(-1, 16)$, $(0, 9)$, $(1, 4)$, $(2, 1)$, $(3, 0)$, $(4, 1)$, $(5, 4)$ on the graph.
- Join all the points by a free-hand smooth curve. This curve is the graph of $y = x^2 - 6x + 9$.
- Here, the curve meets x -axis at only one point $(3, 0)$ and the equation has real and equal roots.
 \therefore The x - coordinate 3 will be the solution.
 \therefore Solution = $\{3, 3\}$

vi) $(2x - 3)(x + 2) = 0$
 $\Rightarrow 2x^2 + x - 6 = 0$

Solution:

Let $y = 2x^2 + x - 6$

$x :$	-3	-2	-1	0	1	2
$2x^2 :$	18	8	2	0	2	8
$x :$	-3	-2	-1	0	1	2
$-6 :$	-6	-6	-6	-6	-6	-6
$y = 2x^2 + x - 6 :$	9	0	-5	-6	-3	4




- Plot the points $(-3, 9)$, $(-2, 0)$, $(-1, -5)$, $(0, -6)$, $(1, -3)$, $(2, 4)$ on the graph.
- Join all the points by a free-hand smooth curve. This curve is the graph of $y = 2x^2 + x - 6$.
- Here, the curve meets x -axis at two points $(-2, 0)$, $(1.5, 0)$ and
 \therefore The equation has real and unequal roots.
 \therefore The x - coordinates are $x = -2, -1.5$ will be the solution.
 \therefore Solution = $\{-2, 3/2\}$

2. Draw the graph of $y = x^2 - 4$ and hence solve $x^2 - x - 12 = 0$

Solution :

First, we draw the graph of $y = x^2 - 9$

$x :$	-4	-3	-2	-1	0	1	2	3	4
$x^2 :$	16	9	4	1	0	1	4	9	16
$-4 :$	-4	-4	-4	-4	-4	-4	-4	-4	-4
$y = x^2 - 4 :$	12	5	0	-3	-4	-3	0	5	12



- Plot the points $(-4, 12)$, $(-3, 5)$, $(-2, 0)$, $(-1, -3)$, $(0, -4)$, $(1, -3)$, $(2, 0)$, $(3, 5)$, $(4, 12)$ on the graph.
- To solve $x^2 - x - 12 = 0$, subtract $x^2 - x - 12 = 0$ from $y = x^2 - 4$.

$$\begin{array}{rcl}
 \text{from} & y = x^2 - 4 & \\
 & y = x^2 + 0x - 4 & \\
 & 0 = x^2 - x - 12 & \\
 \hline
 & y = & x + 8
 \end{array}$$

- We draw the graph of $y = x + 8$.

x	-4	-3	-2	-1	0	1	2	3	4
y	4	5	6	7	8	9	10	11	12

- The line meets the curve at $(-3, 5)$, $(4, 12)$.

\therefore The x - coordinates $x = -3$, $x = 4$ will be the solution of $x^2 - x - 12 = 0$.


\therefore Solution = $\{-3, 4\}$

3. Draw the graph of $y = x^2 + x$ and hence solve $x^2 + 1 = 0$

Solution :

First, we draw the graph of $y = x^2 + x$.

$x :$	-3	-2	-1	0	1	2
$x^2 :$	9	4	1	0	1	4
$x :$	-3	-2	-1	0	1	2
$y = x^2 + x :$	6	2	0	0	2	6



- Plot the points $(-3, 6)$, $(-2, 2)$, $(-1, 0)$, $(0, 0)$, $(1, 2)$, $(2, 6)$ on the graph.
- To solve $x^2 + 1 = 0$, subtract $x^2 + 1 = 0$ from $y = x^2 + x$.

$$\begin{array}{r} y = x^2 + x \\ 0 = x^2 - 0x + 1 \\ \hline y = \quad \quad x - 1 \end{array}$$

- Draw the graph of $y = x - 1$.

x	-4	-3	-2	-1	0	1	2	3	4	5
y	-5	-4	-3	-2	-1	0	1	2	3	4

- The line $y = x - 1$ does not meet the curve $y = x^2 + x$ and the equation has no real roots.

4. Draw the graph of $y = x^2 + 3x + 2$ and use it to solve $x^2 + 2x + 1 = 0$

Solution :

First, we draw the graph of $y = x^2 + 3x + 2$.

$x :$	-4	-3	-2	-1	0	1	2	3
$x^2 :$	16	9	4	1	0	1	4	9
$2 :$	2	2	2	2	2	2	2	2
$y = x^2 + 3x + 2 :$	6	2	0	0	2	6	12	20

- Plot the points $(-4, 6)$, $(-3, 2)$, $(-2, 0)$, $(-1, 0)$, $(0, 2)$, $(1, 6)$, $(2, 12)$, $(3, 20)$ on the graph.
- Join all the points to draw a free-hand smooth curve.
- To solve $x^2 + 2x + 1 = 0$, subtract $x^2 + 2x + 1 = 0$ from $y = x^2 + 3x + 2$.


$$\begin{array}{r} y = x^2 + 3x + 2 \\ 0 = x^2 + 2x + 1 \\ \hline y = \quad \quad x + 1 \end{array}$$

- Draw the graph of $y = x + 1$.

x	-4	-3	-2	-1	0	1	2	3	4
y	-3	-2	-1	0	1	2	3	4	5

- The line $y = x + 1$ meets the curve $y = x^2 + 3x + 2$ at $(-1, 0)$ only and the equation $x^2 + 2x + 1 = 0$ has 2 equal roots.

$$\therefore \text{Solution} = \{-1, -1\}$$



5. Draw the graph of $y = x^2 + 3x - 4$ and hence use it to solve $x^2 + 3x - 4 = 0$

Solution :

First, we draw the graph of $y = x^2 + 3x - 4$.

$x :$	-5	-4	-3	-2	-1	0	1	2	3
$x^2 :$	25	16	9	4	1	0	1	4	9
$3x :$	-15	-12	-9	-6	-3	0	3	6	9
$-4 :$	-4	-4	-4	-4	-4	-4	-4	-4	-4
$y = x^2 + 3x - 4 :$	6	0	-4	-6	-6	-4	0	6	14

- Plot the points $(-4, 0)$, $(-3, -4)$, $(-2, -6)$, $(-1, -6)$, $(0, -4)$, $(1, 0)$, $(2, 6)$, $(3, 14)$, $(4, 24)$ on the graph.
- Join all the points to draw a free-hand smooth curve.
- To solve $x^2 + 3x - 4 = 0$, subtract $x^2 + 3x - 4 = 0$ from $y = x^2 + 3x - 4$.


$$y = x^2 + 3x - 4$$

$$\frac{0 = x^2 + 3x - 4}{y = 0}$$

which is the equation of x - axis.

- The curve meets x -axis at $(-4, 0)$, $(1, 0)$ and the x co-ordinates of the points $x = -4$, $x = 1$ will be the solution of $x^2 + 3x - 4 = 0$.

\therefore Solution = $\{-4, 1\}$




6. Draw the graph of $y = x^2 - 5x - 6$ and hence solve $x^2 - 5x - 14 = 0$

Solution :

First, we draw the graph of $y = x^2 - 5x + 6$.

$x :$	-3	-2	-1	0	1	2	3	4	5	6	7
$x^2 :$	9	4	1	0	1	4	9	16	25	36	49
$-5x :$	15	10	5	0	-5	-10	-15	-20	-25	-30	-35
$-6 :$	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6
$y = x^2 - 5x + 6 :$	18	8	0	-6	-10	-12	-12	-10	-6	0	8



- Plot the points and join them by a hand-free smooth curve.
- To solve $x^2 - 5x - 14 = 0$, subtract $x^2 - 5x - 14 = 0$ from $y = x^2 - 5x - 6$.

$$\begin{array}{r}
 y = x^2 - 5x - 6 \\
 0 = x^2 - 5x - 14 \\
 \hline
 y = 8
 \end{array}$$

a line parallel to x -axis.


- The line $y = 8$ meets the curve $y = x^2 - 5x + 6$ at $(-2, 8)$, $(7, 8)$.
 The x co-ordinates of the points $x = -2$, $x = 7$ will be the solution of $x^2 - 5x - 14 = 0$.
 \therefore Solution = $\{-2, 7\}$

7. Draw the graph of $y = 2x^2 - 3x - 5$ and hence solve $2x^2 - 4x - 6 = 0$

Solution :

First, we draw the graph of $y = 2x^2 - 3x - 5$.

$x :$	-2	-1	0	1	2	3
$2x^2 :$	8	2	0	2	8	18
$-3x :$	6	3	0	-3	-6	-9
$-5 :$	-5	-5	-5	-5	-5	-5
$y = 2x^2 - 3x - 5 :$	9	0	-5	-6	-3	4



- Plot the points on the graph and join all of them by a hand-free smooth curve.
- To solve $2x^2 - 4x - 6 = 0$, subtract it from $y = 2x^2 - 3x - 5$.

$$\begin{array}{r}
 y = 2x^2 - 3x - 5 \\
 0 = 2x^2 - 4x - 6 \\
 \hline
 y = \quad \quad x + 1
 \end{array}$$

- Draw the graph of $y = x + 1$.

x	-4	-3	-2	-1	0	1	2	3	4
y	-3	-2	-1	0	1	2	3	4	5


- The line $y = x + 1$ meets the curve $y = 2x^2 - 3x - 5$ at $(-1, 0)$, $(3, 4)$.
The x co-ordinates of the points $x = -1$, $x = 3$ will be the solution set.
 \therefore Solution = $\{-1, 3\}$

8. Draw the graph of $y = (x - 1)(x + 3)$ and hence solve $x^2 - x - 6 = 0$

Solution :

First, we draw the graph of $y = x^2 + 2x - 3$.

$x :$	-4	-3	-2	-1	0	1	2	3
$x^2 :$	16	9	4	1	0	1	4	9
$2x :$	-8	-6	-4	-2	0	2	4	6
$-3 :$	-3	-3	-3	-3	-3	-3	-3	-3
$y = x^2 + 2x - 3 :$	5	0	-3	-4	-3	0	5	12



- Plot the points on the graph and join all of them by a hand-free smooth curve.
- To solve $x^2 - x - 6 = 0$, subtract it from $y = x^2 + 2x - 3$.

$$\begin{array}{r} y = x^2 + 2x - 3 \\ 0 = x^2 - x - 6 \\ \hline y = 3x + 3 \end{array}$$

- Draw the graph of $y = 3x + 3$.

x	-4	-3	-2	-1	0	1	2	3
y	-9	-6	-3	0	3	6	9	12

- The line meets the curve at $(-2, -3)$, $(3, 12)$.
The x co-ordinates of the points $x = -2$, $x = 3$ which are the solution.
 \therefore Solution = $\{-2, 3\}$

XVI. MATRICES

- ✓ A matrix is a rectangular array of elements. The horizontal arrangements are called rows and vertical arrangements are called columns.
- ✓ If a matrix A has m number of rows and n number of columns, then the order of the matrix A is (Number of rows) \times (Number of columns) that is, $m \times n$. We read $m \times n$ as m cross n or m by n .
- ✓ General form of a matrix A with m rows and n columns (order $m \times n$) can be written in the form

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{13} & \cdots & a_{2n} \\ a_{21} & a_{22} & \cdots & a_{23} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mj} & \cdots & a_{mn} \end{bmatrix}$$

where, a_{11}, a_{12}, \dots denote entries of the matrix.

- ✓ a_{ij} is the element in the i^{th} row and j^{th} column and is referred as $(i, j)^{\text{th}}$ element.
- ✓ The total number of entries in the matrix $A = (a_{ij})_{m \times n}$ is mn .
- ✓ A matrix is said to be a **row matrix** if it has only one row and any number of columns. A row matrix is also called as a row vector.

- ✓ A matrix is said to be a **column matrix** if it has only one column and any number of rows. It is also called as a column vector.
- ✓ A matrix in which the number of rows is equal to the number of columns is called a **square matrix**. Thus a matrix $A = (a_{ij})_{m \times n}$ will be a square matrix if $m = n$.
- ✓ In a square matrix, the elements of the form $a_{11}, a_{22}, a_{33}, \dots$ (i.e) a_{ii} are called leading diagonal elements.
- ✓ A square matrix, all of whose elements, except those in the leading diagonal are zero is called a **diagonal matrix**.
- ✓ A diagonal matrix in which all the leading diagonal elements are equal is called a **scalar matrix**.
- ✓ A square matrix in which elements in the leading diagonal are all “1” and rest are all zero is called an **identity matrix** (or) **unit matrix**.
- ✓ A matrix is said to be a **zero matrix** or null matrix if all its elements are zero.
- ✓ The matrix which is obtained by interchanging the elements in rows and columns of the given matrix A is called transpose of A and is denoted by A^T (read as A transpose).
- ✓ A square matrix in which all the entries above the leading diagonal are zero is called a lower
- ✓ If all the entries below the leading diagonal are zero, then it is called an **upper triangular matrix**.
- ✓ Two matrices A and B are said to be equal if and only if they have the same order and each element of matrix A is equal to the corresponding element of matrix B. That is, $a_{ij} = b_{ij}$ for all i, j .
- ✓ The negative of a matrix $A_{m \times n}$ denoted by $-A_{m \times n}$ is the matrix formed by replacing each element in the matrix $A_{m \times n}$ with its additive inverse.

Example 3.53

Consider the following information regarding the number of men and women workers in three factories I, II and III.

Factory	Men	Women
I	23	18
II	47	36
III	15	16

Represent the above information in the form of a matrix. What does the entry in the second row and first column represent?

Solution :

The information is represented in the form of a 3×2 matrix as follows

$$A = \begin{pmatrix} 23 & 18 \\ 47 & 36 \\ 15 & 16 \end{pmatrix}$$

The entry in the second row and first column represent that there are 47 men workers in factory II.

Example 3.54

If a matrix has 16 elements, what are the possible orders it can have?

Solution :

We know that a matrix of order $m \times n$, has mn elements. Thus to find all possible orders of a matrix with 16 elements, we will find all ordered pairs of natural numbers whose product is 16.

Such ordered pairs are (1,16), (16,1), (4,4), (8,2), (2,8)

Hence possible orders are 1×16 , 16×1 , 4×4 , 2×8 , 8×2 .

Example 3.55

Construct a 3×3 matrix whose elements are $a_{ij} = i^2 j^2$

The general 3×3 matrix is given by

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad a_{ij} = i^2 j^2$$

$$a_{11} = 1^2 \times 1^2 = 1 \times 1 = 1 \quad ;$$

$$a_{12} = 1^2 \times 2^2 = 1 \times 4 = 4 \quad ;$$

$$a_{13} = 1^2 \times 3^2 = 1 \times 9 = 9 \quad ;$$

$$a_{21} = 2^2 \times 1^2 = 4 \times 1 = 4 \quad ;$$

$$a_{22} = 2^2 \times 2^2 = 4 \times 4 = 16 \quad ;$$

$$a_{23} = 2^2 \times 3^2 = 4 \times 9 = 36 \quad ;$$

$$a_{31} = 3^2 \times 1^2 = 9 \times 1 = 9 \quad ;$$

$$a_{32} = 3^2 \times 2^2 = 9 \times 4 = 36 \quad ;$$

$$a_{33} = 3^2 \times 3^2 = 9 \times 9 = 81 \quad ;$$

Hence the required matrix is $A = \begin{pmatrix} 1 & 4 & 9 \\ 4 & 16 & 36 \\ 9 & 36 & 81 \end{pmatrix}$

Example 3.56

Find the value of a, b, c, d from the equation

$$\begin{pmatrix} a-b & 2a+c \\ 2a-b & 3c+d \end{pmatrix} = \begin{pmatrix} 1 & 5 \\ 0 & 2 \end{pmatrix}$$

Solution :

The given matrices are equal. Thus all corresponding elements are equal.

$$\text{Therefore, } a - b = 1 \quad \dots\dots\dots (1)$$

$$2a + c = 5 \quad \dots\dots\dots (2)$$

$$2a - b = 0 \quad \dots\dots\dots (3)$$

$$3c + d = 2 \quad \dots\dots\dots (4)$$

$$(3) \text{ gives } 2a - b = 0$$

$$2a = b \quad \dots\dots\dots (5)$$

Put $2a = b$ in equation (1),

$$a - 2a = 1 \text{ gives } a = -1$$

Put $a = -1$ in equation (5),

$$2(-1) = b \text{ gives } b = -2$$

Put $a = -1$ in equation (2),

$$2(-1) + c = 5 \quad \text{gives } c = 7$$

Put $c = 7$ in equation (4),

$$3(7) + d = 2 \quad \text{gives } d = -19$$

$$\text{Therefore, } a = -1, b = -2, c = 7, d = -19$$

EXERCISE 3.16

1. In the matrix $A = \begin{pmatrix} 8 & 9 & 4 & 3 \\ -1 & \sqrt{7} & \frac{\sqrt{3}}{2} & 5 \\ 1 & 4 & 3 & 0 \\ 6 & 8 & -11 & 1 \end{pmatrix}$, write

(i) The number of elements

(ii) The order of the matrix

(iii) Write the elements a_{22} , a_{23} , a_{24} , a_{34} , a_{43} , a_{44} .

Solution :

i) A has 4 rows and 4 columns

$$\begin{aligned} \text{Number of elements} &= 4 \times 4 \\ &= 16 \end{aligned}$$

ii) Order of the matrix $= 4 \times 4$

$$\begin{aligned} \text{iii) } a_{22} &= \sqrt{7}, a_{23} = \frac{\sqrt{3}}{2} \\ a_{24} &= 5, a_{34} = 0, a_{43} = -11, a_{44} = 1 \end{aligned}$$

2. If a matrix has 18 elements, what are the possible orders it can have? What if it has 6 elements?

Solution :

Given, a matrix has 18 elements

The possible orders of the matrix are

$$18 \times 1, 1 \times 18, 9 \times 2, 2 \times 9, 6 \times 3, 3 \times 6$$

If the matrix has 6 elements

$$\text{The order are } 1 \times 6, 6 \times 1, 3 \times 2, 2 \times 3$$

3. Construct a 3×3 matrix whose elements are given by

$$\text{i) } a_{ij} = |i - 2j| \quad \text{ii) } a_{ij} = \frac{(i+j)^3}{3}$$

Solution :

i) Given $a_{ij} = |i - 2j|$, 3×3

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

$$a_{11} = |1 - 2| = |-1| = 1$$

$$a_{12} = |1 - 4| = |-3| = 3$$

$$a_{13} = |1 - 6| = |-5| = 5$$

$$a_{21} = |2 - 2| = 0$$

$$a_{22} = |2 - 4| = |-2| = 2$$

$$a_{23} = |2 - 6| = |-4| = 4$$

$$a_{31} = |3 - 2| = |1| = 1$$

$$a_{32} = |3 - 4| = |-1| = 1$$

$$a_{33} = |3 - 6| = |-3| = 3$$

$$\therefore A = \begin{pmatrix} 1 & 3 & 5 \\ 0 & 2 & 4 \\ 1 & 1 & 3 \end{pmatrix}$$

$$\text{ii) } a_{ij} = \frac{(i+j)^3}{3}$$

$$a_{11} = \frac{8}{3}, \quad a_{12} = \frac{27}{3} = 9, \quad a_{13} = \frac{64}{3}$$

$$a_{21} = \frac{27}{3} = 9, \quad a_{22} = \frac{64}{3}, \quad a_{23} = \frac{125}{3}$$

$$a_{31} = \frac{64}{3}, \quad a_{32} = \frac{125}{3}, \quad a_{33} = \frac{216}{3} = 72$$

$$\therefore A = \begin{pmatrix} \frac{8}{3} & 9 & \frac{64}{3} \\ 9 & \frac{64}{3} & \frac{125}{3} \\ \frac{64}{3} & \frac{125}{3} & 72 \end{pmatrix}$$

4. If $A = \begin{pmatrix} 5 & 4 & 3 \\ 1 & -7 & 9 \\ 3 & 8 & 2 \end{pmatrix}$ then find the transpose of A.

Solution :

Given

$$A = \begin{pmatrix} 5 & 4 & 3 \\ 1 & -7 & 9 \\ 3 & 8 & 2 \end{pmatrix}$$

$$\therefore A^T = \begin{pmatrix} 5 & 1 & 3 \\ 4 & -7 & 8 \\ 3 & 9 & 2 \end{pmatrix}$$

5. If $A = \begin{pmatrix} \sqrt{7} & -3 \\ -\sqrt{5} & 2 \\ \sqrt{3} & -5 \end{pmatrix}$ then find the transpose of A.

Solution :

Given

$$A = \begin{pmatrix} \sqrt{7} & -3 \\ -\sqrt{5} & 2 \\ \sqrt{3} & -5 \end{pmatrix}$$

$$-A = \begin{pmatrix} -\sqrt{7} & 3 \\ \sqrt{5} & -2 \\ -\sqrt{3} & 5 \end{pmatrix}$$

$$\therefore \text{Transpose of } -A = \begin{pmatrix} -\sqrt{7} & \sqrt{5} & -\sqrt{3} \\ 3 & -2 & 5 \end{pmatrix}$$

6. If $A = \begin{pmatrix} 5 & 2 & 2 \\ -\sqrt{17} & 0.7 & \frac{5}{2} \\ 8 & 3 & 1 \end{pmatrix}$ then verify $(A^T)^T = A$

Solution :

Given

$$A = \begin{pmatrix} 5 & 2 & 2 \\ -\sqrt{17} & 0.7 & \frac{5}{2} \\ 8 & 3 & 1 \end{pmatrix}$$

$$A^T = \begin{pmatrix} 5 & -\sqrt{17} & 8 \\ 2 & 0.7 & 3 \\ 2 & \frac{5}{2} & 1 \end{pmatrix}$$

$$(A^T)^T = \begin{pmatrix} 5 & 2 & 2 \\ -\sqrt{17} & 0.7 & \frac{5}{2} \\ 8 & 3 & 1 \end{pmatrix} = A$$

7. Find the values of x, y and z from the following equations

$$i) \begin{pmatrix} 12 & 3 \\ x & \frac{3}{2} \end{pmatrix} = \begin{pmatrix} y & z \\ 3 & 5 \end{pmatrix} \quad ii) \begin{pmatrix} x+y & 2 \\ 5+z & xy \end{pmatrix} = \begin{pmatrix} 6 & 2 \\ 5 & 8 \end{pmatrix}$$

$$iii) \begin{pmatrix} x+y+z \\ x+z \\ y+z \end{pmatrix} = \begin{pmatrix} 9 \\ 5 \\ 7 \end{pmatrix}$$

Solution :

i) Given

$$\begin{pmatrix} 12 & 3 \\ x & \frac{3}{2} \end{pmatrix} = \begin{pmatrix} y & z \\ 3 & 5 \end{pmatrix}$$

$$\Rightarrow x = 3, y = 12, z = 3$$

ii) Given

$$\begin{pmatrix} x+y & 2 \\ 5+z & xy \end{pmatrix} = \begin{pmatrix} 6 & 2 \\ 5 & 8 \end{pmatrix}$$

$$\Rightarrow x+y=6, \quad xy=8, \quad 5+z=5$$

$$x=2 \text{ (or) } 4, \quad \Rightarrow z=0$$

$$y=4 \text{ (or) } 2$$

iii) Given

$$\begin{pmatrix} x+y+z \\ x+z \\ y+z \end{pmatrix} = \begin{pmatrix} 9 \\ 5 \\ 7 \end{pmatrix}$$

$$\begin{array}{l} \Rightarrow x+y+z=9 \\ \Rightarrow 5+y=9 \\ \Rightarrow y=4 \end{array} \quad \begin{array}{l} x+z=5 \\ \Rightarrow x+3=5 \\ \Rightarrow x=2 \end{array} \quad \begin{array}{l} y+z=7 \\ 4+z=7 \\ \Rightarrow z=3 \end{array}$$

XVII. OPERATIONS ON MATRICES

Key Points

- ✓ Two matrices can be added or subtracted if they have the same order. To add or subtract two matrices, simply add or subtract the corresponding elements.
- ✓ We can multiply the elements of the given matrix A by a non-zero number k to obtain a new matrix kA whose elements are multiplied by k . The matrix kA is called scalar multiplication of A .
- ✓ If A be any given matrix then $-A$ is the additive inverse of A .

Example 3.57

$$\text{If } A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}, B = \begin{pmatrix} 1 & 7 & 0 \\ 1 & 3 & 1 \\ 2 & 4 & 0 \end{pmatrix}, \text{ find } A+B$$

Solution :

$$A+B = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} + \begin{pmatrix} 1 & 7 & 0 \\ 1 & 3 & 1 \\ 2 & 4 & 0 \end{pmatrix} = \begin{pmatrix} 1+1 & 2+7 & 3+0 \\ 4+1 & 5+3 & 6+1 \\ 7+2 & 8+4 & 9+0 \end{pmatrix} = \begin{pmatrix} 2 & 9 & 3 \\ 5 & 8 & 7 \\ 9 & 12 & 9 \end{pmatrix}$$

Example 3.58

Two examinations were conducted for three groups of students namely group 1, group 2, group 3 and their data on average of marks for the subjects Tamil, English, Science and Mathematics are given below in the form of matrices A and B . Find the total marks of both the examinations for all the three groups.

$$A = \begin{matrix} & \begin{pmatrix} \text{Tamil} & \text{English} & \text{Science} & \text{Mathematics} \end{pmatrix} \\ \begin{pmatrix} \text{Group 1} \\ \text{Group 2} \\ \text{Group 3} \end{pmatrix} & \begin{pmatrix} 22 & 15 & 14 & 23 \\ 50 & 62 & 21 & 30 \\ 53 & 80 & 32 & 40 \end{pmatrix} \end{matrix}$$

$$B = \begin{matrix} & \begin{matrix} \text{Tamil} & \text{English} & \text{Science} & \text{Mathematics} \end{matrix} \\ \begin{matrix} \text{Group 1} \\ \text{Group 2} \\ \text{Group 3} \end{matrix} & \begin{pmatrix} 20 & 38 & 15 & 40 \\ 18 & 12 & 17 & 80 \\ 81 & 47 & 52 & 18 \end{pmatrix} \end{matrix}$$

Solution :

The total marks in both the examinations for all the three groups is the sum of the given matrices.

$$A + B = \begin{pmatrix} 22+20 & 15+38 & 14+15 & 23+40 \\ 50+18 & 62+12 & 21+17 & 30+80 \\ 53+81 & 80+47 & 32+52 & 40+18 \end{pmatrix} = \begin{pmatrix} 42 & 53 & 29 & 63 \\ 68 & 74 & 38 & 110 \\ 134 & 127 & 84 & 58 \end{pmatrix}$$

Example 3.59

$$\text{If } A = \begin{pmatrix} 1 & 3 & -2 \\ 5 & -4 & 6 \\ -3 & 2 & 9 \end{pmatrix} \text{ } B = \begin{pmatrix} 1 & 8 \\ 3 & 4 \\ 9 & 6 \end{pmatrix} \text{ find } A+B.$$

It is not possible to add A and B because they have different orders.

Example 3.60

$$\text{If } A = \begin{pmatrix} 7 & 8 & 6 \\ 1 & 3 & 9 \\ -4 & 3 & -1 \end{pmatrix} \text{ } B = \begin{pmatrix} 4 & 11 & -3 \\ -1 & 2 & 4 \\ 7 & 5 & 0 \end{pmatrix} \text{ then Find } 2A + B.$$

Solution :

Since A and B have same order 3×3 , $2A + B$ is defined.

$$\begin{aligned} \text{We have } 2A + B &= 2 \begin{pmatrix} 7 & 8 & 6 \\ 1 & 3 & 9 \\ -4 & 3 & -1 \end{pmatrix} + \begin{pmatrix} 4 & 11 & -3 \\ -1 & 2 & 4 \\ 7 & 5 & 0 \end{pmatrix} = \begin{pmatrix} 14 & 16 & 12 \\ 2 & 6 & 18 \\ -8 & 6 & -2 \end{pmatrix} + \begin{pmatrix} 4 & 11 & -3 \\ -1 & 2 & 4 \\ 7 & 5 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 18 & 27 & 9 \\ 1 & 8 & 22 \\ -1 & 11 & -2 \end{pmatrix} \end{aligned}$$

Example 3.61

If $A = \begin{pmatrix} 5 & 4 & -2 \\ \frac{1}{2} & \frac{3}{4} & \sqrt{2} \\ 1 & 9 & 4 \end{pmatrix}$ $B = \begin{pmatrix} -7 & 4 & -3 \\ \frac{1}{4} & \frac{7}{2} & 3 \\ 5 & -6 & 9 \end{pmatrix}$ then
Find $4A - 3B$.

Solution :

Since A, B are of the same order 3×3 , subtraction of $4A$ and $3B$ is defined.

$$\begin{aligned} 4A - 3B &= 4 \begin{pmatrix} 5 & 4 & -2 \\ \frac{1}{2} & \frac{3}{4} & \sqrt{2} \\ 1 & 9 & 4 \end{pmatrix} - 3 \begin{pmatrix} -7 & 4 & -3 \\ \frac{1}{4} & \frac{7}{2} & 3 \\ 5 & -6 & 9 \end{pmatrix} \\ &= \begin{pmatrix} 20 & 6 & -8 \\ 2 & 3 & 4\sqrt{2} \\ 4 & 36 & 16 \end{pmatrix} + \begin{pmatrix} 21 & -12 & 9 \\ -\frac{3}{4} & -\frac{21}{2} & -9 \\ -15 & 18 & -27 \end{pmatrix} \\ &= \begin{pmatrix} 41 & 4 & 1 \\ \frac{5}{4} & -\frac{15}{2} & 4\sqrt{2} - 9 \\ -11 & 54 & -11 \end{pmatrix} \end{aligned}$$

Example 3.62

Find the value of a, b, c, d, x, y from the following matrix equation.

$$\begin{pmatrix} d & 8 \\ 3b & a \end{pmatrix} + \begin{pmatrix} 3 & a \\ -2 & -4 \end{pmatrix} = \begin{pmatrix} 2 & 2a \\ b & 4c \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ -5 & 0 \end{pmatrix}$$

Solution :

First, we add the two matrices on both left, right hand sides to get

$$\begin{pmatrix} d+3 & 8+a \\ 3b-2 & a-4 \end{pmatrix} = \begin{pmatrix} 2 & 2a+1 \\ b-5 & 4c \end{pmatrix}$$

Equating the corresponding elements of the two matrices, we have

$$d + 3 = 2 \quad \text{gives } d = -1$$

$$8 + a = 2a + 1 \quad \text{gives } a = 7$$

$$3b - 2 = b - 5 \quad \text{gives } b = \frac{-3}{2}$$

$$\text{Substituting } a = 7 \text{ in } a - 4 = 4c \text{ gives } c = \frac{3}{4}$$

$$\text{Therefore, } a = 7, b = -\frac{3}{2}, c = \frac{3}{4}, d = -1$$

Example 3.63

If

$$A = \begin{pmatrix} 1 & 8 & 3 \\ 3 & 5 & 0 \\ 8 & 7 & 6 \end{pmatrix}, B = \begin{pmatrix} 8 & -6 & -4 \\ 2 & 11 & -3 \\ 0 & 1 & 5 \end{pmatrix}, C = \begin{pmatrix} 5 & 3 & 0 \\ -1 & -7 & 2 \\ 1 & 4 & 3 \end{pmatrix}$$

compute the following :

$$(i) 3A + 2B - C \quad \quad (ii) \frac{1}{2}A - \frac{3}{2}B$$

Solution :

$$i) 3A + 2B - C$$

$$\begin{aligned} &= 3 \begin{pmatrix} 1 & 8 & 3 \\ 3 & 5 & 0 \\ 8 & 7 & 6 \end{pmatrix} + 2 \begin{pmatrix} 8 & -6 & -4 \\ 2 & 11 & -3 \\ 0 & 1 & 5 \end{pmatrix} - \begin{pmatrix} 5 & 3 & 0 \\ -1 & -7 & 2 \\ 1 & 4 & 3 \end{pmatrix} \\ &= \begin{pmatrix} 3 & 24 & 9 \\ 9 & 15 & 0 \\ 24 & 21 & 18 \end{pmatrix} + \begin{pmatrix} 16 & -12 & -8 \\ 4 & 22 & -6 \\ 0 & 2 & 10 \end{pmatrix} + \begin{pmatrix} -5 & -3 & 0 \\ 1 & 7 & -2 \\ -1 & -4 & -3 \end{pmatrix} \\ &= \begin{pmatrix} 14 & 9 & 1 \\ 14 & 44 & -8 \\ 23 & 19 & 25 \end{pmatrix} \end{aligned}$$

$$ii) \frac{1}{2}A - \frac{3}{2}B$$

$$\begin{aligned} &= \frac{1}{2} (A - 3B) \\ &= \frac{1}{2} \left(\begin{pmatrix} 1 & 8 & 3 \\ 3 & 5 & 0 \\ 8 & 7 & 6 \end{pmatrix} - 3 \begin{pmatrix} 8 & -6 & -4 \\ 2 & 11 & -3 \\ 0 & 1 & 5 \end{pmatrix} \right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \left(\begin{pmatrix} 1 & 8 & 3 \\ 3 & 5 & 0 \\ 8 & 7 & 6 \end{pmatrix} + \begin{pmatrix} -24 & 18 & 12 \\ -6 & -33 & 9 \\ 0 & -3 & -15 \end{pmatrix} \right) \\
 &= \frac{1}{2} \begin{pmatrix} -23 & 26 & 15 \\ -3 & -28 & 9 \\ 8 & 4 & -9 \end{pmatrix} \\
 &= \begin{pmatrix} -\frac{23}{2} & 13 & \frac{15}{2} \\ -\frac{3}{2} & -14 & \frac{9}{2} \\ 4 & 2 & -\frac{9}{2} \end{pmatrix}
 \end{aligned}$$

EXERCISE 3.17

1. If $A = \begin{pmatrix} 1 & 9 \\ 3 & 4 \end{pmatrix}$, $B = \begin{pmatrix} 5 & 7 \\ 3 & 3 \end{pmatrix}$ then verify that

(i) $A + B = B + A$

ii) $A + (-A) = (-A) + A = O$

Solution :

Given $A = \begin{pmatrix} 1 & 9 \\ 3 & 4 \end{pmatrix}$, $B = \begin{pmatrix} 5 & 7 \\ 3 & 3 \end{pmatrix}$

- i) To verify : $A + B = B + A$

A & B are of same order

$$\therefore A + B = \begin{pmatrix} 6 & 16 \\ 6 & 7 \\ 9 & -3 \end{pmatrix}$$

$$B + A = \begin{pmatrix} 6 & 16 \\ 6 & 7 \\ 9 & -3 \end{pmatrix}$$

$$\therefore A + B = B + A$$

- ii) To verify : $A + (-A) = (-A) + A = O$

LHS : $A + (-A)$

$$= \begin{pmatrix} 1 & 9 \\ 3 & 4 \\ 8 & -3 \end{pmatrix} + \begin{pmatrix} -1 & -9 \\ -3 & -4 \\ -8 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} = O$$

RHS : $(-A) + A$

$$= \begin{pmatrix} -1 & -9 \\ -3 & -4 \\ -8 & 3 \end{pmatrix} + \begin{pmatrix} 1 & 9 \\ 3 & 4 \\ 8 & -3 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} = O$$

$$\therefore \text{LHS} = \text{RHS}$$

2. If $A = \begin{pmatrix} 4 & 3 & 1 \\ 2 & 3 & -8 \\ 1 & 0 & -4 \end{pmatrix}$, $B = \begin{pmatrix} 2 & 3 & 4 \\ 1 & 9 & 2 \\ -7 & 1 & -1 \end{pmatrix}$

and $C = \begin{pmatrix} 8 & 3 & 4 \\ 1 & -2 & 3 \\ 2 & 4 & -1 \end{pmatrix}$ then verify that

$$A + (B + C) = (A + B) + C$$

Solution :

$$B + C = \begin{pmatrix} 10 & 6 & 8 \\ 2 & 7 & 5 \\ -5 & 5 & -2 \end{pmatrix}$$

$$A + (B + C) = \begin{pmatrix} 4 & 3 & 1 \\ 2 & 3 & -8 \\ 1 & 0 & -4 \end{pmatrix} + \begin{pmatrix} 10 & 6 & 8 \\ 2 & 7 & 5 \\ -5 & 5 & -2 \end{pmatrix}$$

$$= \begin{pmatrix} 14 & 9 & 9 \\ 4 & 10 & -3 \\ -4 & 5 & -6 \end{pmatrix} \quad \dots\dots\dots(1)$$

$$A+B = \begin{pmatrix} 6 & 6 & 5 \\ 3 & 12 & -6 \\ -6 & 1 & -5 \end{pmatrix}$$

$$\therefore (A+B)+C = \begin{pmatrix} 6 & 6 & 5 \\ 3 & 12 & -6 \\ -6 & 1 & -5 \end{pmatrix} + \begin{pmatrix} 8 & 3 & 4 \\ 1 & -2 & 3 \\ 2 & 4 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} 14 & 9 & 9 \\ 4 & 10 & -3 \\ -4 & 5 & -6 \end{pmatrix} \quad \dots\dots\dots(2)$$

\therefore From (1) & (2) LHS = RHS

3. Find X and Y if $X + Y = \begin{pmatrix} 7 & 0 \\ 3 & 5 \end{pmatrix}$ and

$$X - Y = \begin{pmatrix} 3 & 0 \\ 0 & 4 \end{pmatrix}$$

Solution :

Given $X + Y = \begin{pmatrix} 7 & 0 \\ 3 & 5 \end{pmatrix} \quad \dots\dots\dots(1)$

$$X - Y = \begin{pmatrix} 3 & 0 \\ 0 & 4 \end{pmatrix} \quad \dots\dots\dots(2)$$

$$(1) + (2) \Rightarrow 2X = \begin{pmatrix} 10 & 0 \\ 3 & 9 \end{pmatrix}$$

$$\Rightarrow X = \begin{pmatrix} 5 & 0 \\ 3/2 & 9/2 \end{pmatrix}$$

$$(1) - (2) \Rightarrow 2Y = \begin{pmatrix} 4 & 0 \\ 3 & 1 \end{pmatrix}$$

$$\Rightarrow Y = \begin{pmatrix} 2 & 0 \\ 3/2 & 1/2 \end{pmatrix}$$

4. If $A = \begin{pmatrix} 0 & 4 & 9 \\ 8 & 3 & 7 \end{pmatrix}$, $B = \begin{pmatrix} 7 & 3 & 8 \\ 1 & 4 & 9 \end{pmatrix}$ find the

value of i) $B - 5A$ ii) $3A - 9B$

Solution :

Given

$$A = \begin{pmatrix} 0 & 4 & 9 \\ 8 & 3 & 7 \end{pmatrix}, B = \begin{pmatrix} 7 & 3 & 8 \\ 1 & 4 & 9 \end{pmatrix}$$

$$\begin{aligned} \text{i) } B - 5A &= \begin{pmatrix} 7 & 3 & 8 \\ 1 & 4 & 9 \end{pmatrix} - \begin{pmatrix} 0 & 20 & 45 \\ 40 & 15 & 35 \end{pmatrix} \\ &= \begin{pmatrix} 7 & -17 & -37 \\ -39 & -11 & -26 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \text{ii) } 3A - 9B &= \begin{pmatrix} 0 & 12 & 27 \\ 24 & 9 & 21 \end{pmatrix} - \begin{pmatrix} 63 & 27 & 72 \\ 9 & 36 & 81 \end{pmatrix} \\ &= \begin{pmatrix} -63 & -65 & -45 \\ 15 & -27 & -60 \end{pmatrix} \end{aligned}$$

5. Find the values of x, y, z if

$$\text{i) } \begin{pmatrix} x-3 & 3x-z \\ x+y+7 & x+y+z \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 6 \end{pmatrix}$$

$$\text{ii) } (x \ y - z \ z + 3) + (y \ 4 \ 3) = (4 \ 8 \ 16)$$

Solution :

$$\text{i) Given } \begin{pmatrix} x-3 & 3x-z \\ x+y+7 & x+y+z \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 6 \end{pmatrix}$$

$$\begin{aligned} \Rightarrow x-3 &= 1 & \left| \begin{array}{l} 3x-z=0 \\ 12-z=0 \\ z=12 \end{array} \right. & \left| \begin{array}{l} x+y+7=1 \\ \Rightarrow x+y=-6 \\ \Rightarrow 4+y=-6 \\ \Rightarrow y=-10 \end{array} \right. \end{aligned}$$

$$\text{ii) } (x \ y - z \ z + 3) + (y \ 4 \ 3) = (4 \ 8 \ 16)$$

$$\begin{aligned} \Rightarrow x+y &= 4 & \left| \begin{array}{l} y-z+4=8 \\ y-z=4 \\ y-10=4 \end{array} \right. & \left| \begin{array}{l} z+6=16 \\ z=10 \\ \therefore y=14 \end{array} \right. \end{aligned}$$

$$\therefore x = -10, y = 14, z = 10$$

6. Find x and y if $x \begin{pmatrix} 4 \\ -3 \end{pmatrix} + y \begin{pmatrix} -2 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$

Solution :

$$\text{Given } x \begin{pmatrix} 4 \\ -3 \end{pmatrix} + y \begin{pmatrix} -2 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$$

$$\Rightarrow 4x - 2y = 4 \quad -3x + 3y = 6$$

$$\Rightarrow 2x - y = 2 \quad \dots\dots\dots (1)$$

$$-x + y = 2 \quad \dots\dots\dots (2)$$

$$(1) \Rightarrow 2x - y = 2$$

$$(2) \Rightarrow \underline{-x + y = 2}$$

$$\text{Adding,} \quad \underline{x = 4}$$

Sub $x = 4$ in (2)

$$-4 + y = 2 \Rightarrow y = 6$$

$$\therefore x = 4, y = 6$$

the matrix equation

$$x \begin{pmatrix} 2x & 2 \\ 3 & x \end{pmatrix} + 2 \begin{pmatrix} 8 & 5x \\ 4 & 4x \end{pmatrix} = 2 \begin{pmatrix} x^2 + 8 & 24 \\ 10 & 6x \end{pmatrix}$$

Solution :

$$\text{Given } x \begin{pmatrix} 2x & 2 \\ 3 & x \end{pmatrix} + 2 \begin{pmatrix} 8 & 5x \\ 4 & 4x \end{pmatrix} = 2 \begin{pmatrix} x^2 + 8 & 24 \\ 10 & 6x \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} 2x^2 & 2x \\ 3x & x^2 \end{pmatrix} + \begin{pmatrix} 16 & 10x \\ 8 & 8x \end{pmatrix} = \begin{pmatrix} 2x^2 + 16 & 48 \\ 20 & 12x \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} 2x^2 + 16 & 12x \\ 3x + 8 & x^2 + 8x \end{pmatrix} = \begin{pmatrix} 2x^2 + 16 & 48 \\ 20 & 12x \end{pmatrix}$$

$$\begin{array}{l} \therefore 12x = 48 \Rightarrow x = 4 \\ 3x + 8 = 20 \Rightarrow 3x = 12 \\ \Rightarrow x = 4 \end{array} \left| \begin{array}{l} x^2 + 8x = 12x \\ x^2 - 4x = 0 \\ x(x - 4) = 0 \\ \Rightarrow x = 0, x = 4 \end{array} \right.$$

$$\therefore x = 4$$

8. Solve for x, y $\begin{pmatrix} x^2 \\ y^2 \end{pmatrix} + 2 \begin{pmatrix} -2x \\ y \end{pmatrix} = \begin{pmatrix} 5 \\ 8 \end{pmatrix}$

Solution :

$$\text{Given } \begin{pmatrix} x^2 \\ y^2 \end{pmatrix} + 2 \begin{pmatrix} -2x \\ y \end{pmatrix} = \begin{pmatrix} 5 \\ 8 \end{pmatrix}$$

$$\begin{array}{l} \Rightarrow x^2 - 4x = 5 \\ \Rightarrow x^2 - 4x - 5 = 0 \\ \Rightarrow (x - 5)(x + 1) = 0 \\ \Rightarrow x = 5, -1 \end{array} \left| \begin{array}{l} y^2 - 2y = 8 \\ y^2 - 2y - 8 = 0 \\ (y - 4)(y + 2) = 0 \\ \therefore y = 4, y = -2 \end{array} \right.$$

XVIII. MULTIPLICATION OF MATRICES

Key Points

- ✓ To multiply two matrices, the number of columns in the first matrix must be equal to the number of rows in the second matrix.
- ✓ Matrices are multiplied by multiplying the elements in a row of the first matrix by the elements in a column of the second matrix, and adding the results.
- ✓ Matrix multiplication is not commutative in general.
- ✓ Matrix multiplication is distributive over matrix addition.
- ✓ Matrix multiplication is always associative.
- ✓ If A is a square matrix of order $n \times n$ and I is the unit matrix of same order then $AI = IA = A$.
- ✓ $AB = 0$ does not necessarily imply that $A = 0$ or $B = 0$ or both $A, B = 0$

Example 3.64

If $A = \begin{pmatrix} 1 & 2 & 0 \\ 3 & 1 & 5 \end{pmatrix}$, $B = \begin{pmatrix} 8 & 3 & 1 \\ 2 & 4 & 1 \\ 5 & 3 & 1 \end{pmatrix}$ find AB.

Solution :

We observe that A is a 2×3 matrix and B is a 3×3 matrix, hence AB is defined and it will be of the order 2×3 .

$$\begin{aligned} \text{Given } A &= \begin{pmatrix} 1 & 2 & 0 \\ 3 & 1 & 5 \end{pmatrix}_{2 \times 3}, B = \begin{pmatrix} 8 & 3 & 1 \\ 2 & 4 & 1 \\ 5 & 3 & 1 \end{pmatrix}_{3 \times 3} \\ AB &= \begin{pmatrix} 1 & 2 & 0 \\ 3 & 1 & 5 \end{pmatrix} \times \begin{pmatrix} 8 & 3 & 1 \\ 2 & 4 & 1 \\ 5 & 3 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 8+4+0 & 3+8+0 & 1+2+0 \\ 24+2+25 & 9+4+15 & 3+1+5 \end{pmatrix} \\ &= \begin{pmatrix} 12 & 11 & 3 \\ 51 & 28 & 9 \end{pmatrix} \end{aligned}$$

Example 3.65

If $A = \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}$, $B = \begin{pmatrix} 2 & 0 \\ 1 & 3 \end{pmatrix}$ find AB and BA.
Check if $AB = BA$.

Solution :

We observe that A is a 2×2 matrix and B is a 2×2 matrix, hence AB is defined and it will be of the order 2×2 .

$$\begin{aligned} AB &= \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix} \times \begin{pmatrix} 2 & 0 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 4+1 & 0+3 \\ 2+3 & 0+9 \end{pmatrix} = \begin{pmatrix} 5 & 3 \\ 5 & 9 \end{pmatrix} \\ BA &= \begin{pmatrix} 2 & 0 \\ 1 & 3 \end{pmatrix} \times \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 4+0 & 2+0 \\ 2+3 & 1+9 \end{pmatrix} = \begin{pmatrix} 4 & 2 \\ 5 & 10 \end{pmatrix} \end{aligned}$$

Therefore, $AB \neq BA$

Example 3.66

If $A = \begin{pmatrix} 2 & -2\sqrt{2} \\ \sqrt{2} & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 2\sqrt{2} \\ -\sqrt{2} & 2 \end{pmatrix}$

Show that A and B satisfy commutative property

with respect to matrix multiplication.

Solution :

We have to show that $AB = BA$

$$\begin{aligned} \text{LHS: } AB &= \begin{pmatrix} 2 & -2\sqrt{2} \\ \sqrt{2} & 2 \end{pmatrix} \times \begin{pmatrix} 2 & 2\sqrt{2} \\ -\sqrt{2} & 2 \end{pmatrix} \\ &= \begin{pmatrix} 4+4 & 4\sqrt{2}-4\sqrt{2} \\ 2\sqrt{2}-2\sqrt{2} & 4+4 \end{pmatrix} \\ &= \begin{pmatrix} 8 & 0 \\ 0 & 8 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \text{RHS: } BA &= \begin{pmatrix} 2 & 2\sqrt{2} \\ -\sqrt{2} & 2 \end{pmatrix} \times \begin{pmatrix} 2 & -2\sqrt{2} \\ \sqrt{2} & 2 \end{pmatrix} \\ &= \begin{pmatrix} 4+4 & -4\sqrt{2}+4\sqrt{2} \\ -2\sqrt{2}+2\sqrt{2} & 4+4 \end{pmatrix} \\ &= \begin{pmatrix} 8 & 0 \\ 0 & 8 \end{pmatrix} \end{aligned}$$

Hence $\text{LHS} = \text{RHS}$ (ie) $AB = BA$

Example 3.67

Solve $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$

Solution :

$$\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}_{2 \times 2} \times \begin{pmatrix} x \\ y \end{pmatrix}_{2 \times 1} = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$$

By matrix multiplication $\begin{pmatrix} 2x+y \\ x+2y \end{pmatrix} = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$

Rewriting $2x + y = 4$ (1)

$x + 2y = 5$ (2)

(1) - 2 × (2) gives $2x + y = 4$
 $\frac{2x + 4y = 10}{-3y = -6}$ (-) gives $y = 2$

Substituting $y = 2$ in (1), $2x + 2 = 4$ gives $x = 1$

Therefore, $x = 1, y = 2$.

Example 3.68

If $A = \begin{pmatrix} 1 & -1 & 2 \end{pmatrix}$, $B = \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}$ and $C = \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix}$

show that $(AB)C = A(BC)$.

Solution :

LHS $(AB)C$

$$\begin{aligned} A &= \begin{pmatrix} 1 & -1 & 2 \end{pmatrix}_{1 \times 3} \times \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}_{3 \times 2} \\ &= (1 - 2 + 2 - 1 - 1 + 6) = (1 \ 4) \\ (AB)C &= (1 \ 4)_{1 \times 2} \times \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix} \\ &= (1 + 8 \ 2 - 4) = (9 \ -2) \quad \text{..... (1)} \end{aligned}$$

RHS $A(BC)$

$$\begin{aligned} BC &= \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}_{2 \times 2} \times \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix}_{2 \times 2} \\ &= \begin{pmatrix} 1 - 2 & 2 + 1 \\ 2 + 2 & 4 - 1 \end{pmatrix} = \begin{pmatrix} -1 & 3 \\ 4 & 3 \end{pmatrix} \\ A(BC) &= \begin{pmatrix} 1 & -1 & 2 \end{pmatrix}_{1 \times 3} \times \begin{pmatrix} -1 & 3 \\ 4 & 3 \\ 7 & -1 \end{pmatrix}_{3 \times 2} \end{aligned}$$

$$\begin{aligned} A(BC) &= (-1 - 4 + 14 \quad 3 - 3 - 2) \\ &= (9 \ -2) \quad \text{..... (2)} \end{aligned}$$

From (1) and (2), $(AB)C = A(BC)$.

Example 3.69

If $A = \begin{pmatrix} 1 & 1 \\ -1 & 3 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 2 \\ -4 & 2 \end{pmatrix}$, $C = \begin{pmatrix} -7 & 6 \\ 3 & 2 \end{pmatrix}$

verify that $A(B + C) = AB + AC$.

Solution :

LHS $A(B + C)$

$$B + C = \begin{pmatrix} 1 & 2 \\ -4 & 2 \end{pmatrix} + \begin{pmatrix} -7 & 6 \\ 3 & 2 \end{pmatrix} = \begin{pmatrix} -6 & 8 \\ -1 & 4 \end{pmatrix}$$

$$\begin{aligned} A(B + C) &= \begin{pmatrix} 1 & 1 \\ -1 & 3 \end{pmatrix} \times \begin{pmatrix} -6 & 8 \\ -1 & 4 \end{pmatrix} \\ &= \begin{pmatrix} -6 - 1 & 8 + 4 \\ 6 - 3 & -8 + 12 \end{pmatrix} = \begin{pmatrix} -7 & 12 \\ 3 & 4 \end{pmatrix} \quad \text{... (1)} \end{aligned}$$

RHS $AB + AC$

$$\begin{aligned} AB &= \begin{pmatrix} 1 & 1 \end{pmatrix} \times \begin{pmatrix} 1 & 2 \\ -4 & 2 \end{pmatrix} = \begin{pmatrix} 1 - 4 & 2 + 2 \end{pmatrix} \\ &= \begin{pmatrix} -3 & 4 \\ -13 & 4 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} AC &= \begin{pmatrix} 1 & 1 \\ -1 & 3 \end{pmatrix} \times \begin{pmatrix} -7 & 6 \\ 3 & 2 \end{pmatrix} = \begin{pmatrix} -7 + 3 & 6 + 2 \\ 7 + 9 & -6 + 6 \end{pmatrix} \\ &= \begin{pmatrix} -4 & 8 \\ 16 & 0 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \text{Therefore, } AB + AC &= \begin{pmatrix} -3 & 4 \\ -13 & 4 \end{pmatrix} + \begin{pmatrix} -4 & 8 \\ 16 & 0 \end{pmatrix} \\ &= \begin{pmatrix} -7 & 12 \\ 3 & 4 \end{pmatrix} \quad \text{..... (2)} \end{aligned}$$

From (1) and (2), $A(B + C) = AB + AC$.
Hence proved.

Example 3.70

If $A = \begin{pmatrix} 1 & 2 & 1 \\ 2 & -1 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & -1 \\ -1 & 4 \\ 0 & 2 \end{pmatrix}$ show that $(AB)^T = B^T A^T$.

Solution :LHS $(AB)^T$

$$\begin{aligned}
 AB &= \begin{pmatrix} 1 & 2 & 1 \\ 2 & -1 & 1 \end{pmatrix}_{2 \times 3} \times \begin{pmatrix} 2 & -1 \\ -1 & 4 \\ 0 & 2 \end{pmatrix}_{3 \times 2} \\
 &= \begin{pmatrix} 2-2+0 & -1+8+2 \\ 4+1+0 & -2-4+2 \end{pmatrix} = \begin{pmatrix} 0 & 9 \\ 5 & -4 \end{pmatrix} \\
 (AB)^T &= \begin{pmatrix} 0 & 9 \\ 5 & -4 \end{pmatrix}^T = \begin{pmatrix} 0 & 5 \\ 9 & -4 \end{pmatrix} \quad \dots\dots\dots (1)
 \end{aligned}$$

RHS $(B^T A^T)$

$$\begin{aligned}
 B^T &= \begin{pmatrix} 2 & -1 & 0 \\ -1 & 4 & 2 \end{pmatrix}, A^T = \begin{pmatrix} 1 & 2 \\ 2 & -1 \\ 1 & 1 \end{pmatrix} \\
 B^T A^T &= \begin{pmatrix} 2 & -1 & 0 \\ -1 & 4 & 2 \end{pmatrix}_{2 \times 3} = \begin{pmatrix} 1 & 2 \\ 2 & -1 \\ 1 & 1 \end{pmatrix}_{3 \times 2} \\
 &= \begin{pmatrix} 2-2+0 & 4+1+0 \\ -1+8+2 & -2-4+2 \end{pmatrix} \\
 B^T A^T &= \begin{pmatrix} 0 & 5 \\ 9 & -4 \end{pmatrix} \quad \dots\dots\dots (2)
 \end{aligned}$$

From (1) and (2), $(AB)^T = B^T A^T$.

Hence proved.

EXERCISE 3.18**1. Find the order of the product matrix AB if**

	(i)	(ii)	(iii)	(iv)	(v)
Orders of A	3×3	4×3	4×2	4×5	1×1
Orders of B	3×3	3×2	2×2	5×1	1×3

Solution :i) A is of order $= 3 \times 3$ B is of order $= 3 \times 3$ \therefore Order of AB $= 3 \times 3$ ii) $A \rightarrow 4 \times 3$, $B \rightarrow 3 \times 2$ \therefore Order of AB $= 4 \times 2$ iii) $A \rightarrow 4 \times 2$, $B \rightarrow 2 \times 2$ iv) $A \rightarrow 4 \times 5$, $B \rightarrow 5 \times 1$ \therefore Order of AB $= 4 \times 1$ v) $A \rightarrow 1 \times 1$, $B \rightarrow 1 \times 3$ \therefore Order of AB $= 1 \times 3$ **2. If A is of order $p \times q$ and B is of order $q \times r$ what is the order of AB and BA?****Solution :**Given A is of order $p \times q$ B is of order $q \times r$ \therefore Order of AB $= p \times r$

Order of BA is not defined

(\because No. of columns in B & no. of rows in A are not equal.)

3. A has 'a' rows and 'a + 3' columns. B has 'b' rows and '17-b' columns, and if both products AB and BA exist, find a, b?

Solution :

Given Order of A is $a \times (a + 3)$

Order of B is $b \times (17 - b)$

Product AB exist

$$\Rightarrow a + 3 = b$$

(No. of columns in A = No. of rows in B)

$$\Rightarrow a - b = -3 \dots\dots (1)$$

Product BA exist

$$\Rightarrow 17 - b = a$$

(No. of columns in B = No. of rows in A)

$$\Rightarrow a + b = 17 \dots\dots (2)$$

$$2a = 14$$

$$a = 7$$

Sub $a = 7$ in (1)

$$7 - b = -3$$

$$b = 10$$

$$\therefore a = 7, b = 10$$

4. If $A = \begin{pmatrix} 2 & 5 \\ 4 & 3 \end{pmatrix}$, $B = \begin{pmatrix} 1 & -3 \\ 2 & 5 \end{pmatrix}$ find AB, BA and check if $AB = BA$?

Solution :

$$\text{Given } A = \begin{pmatrix} 2 & 5 \\ 4 & 3 \end{pmatrix}, B = \begin{pmatrix} 1 & -3 \\ 2 & 5 \end{pmatrix}$$

$$AB = \begin{pmatrix} 2 & 5 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} 1 & -3 \\ 2 & 5 \end{pmatrix}$$

$$= \begin{pmatrix} 2 \times 1 + 5 \times 2 & 2 \times -3 + 5 \times 5 \\ 4 \times 1 + 3 \times 2 & 4 \times -3 + 3 \times 5 \end{pmatrix}$$

$$= \begin{pmatrix} 2 + 10 & -6 + 25 \\ 4 + 6 & -12 + 15 \end{pmatrix}$$

$$= \begin{pmatrix} 12 & 19 \\ 10 & 3 \end{pmatrix}$$

$$BA = \begin{pmatrix} 1 & -3 \\ 2 & 5 \end{pmatrix} \begin{pmatrix} 2 & 5 \\ 4 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} (1 \times 2) + (-3 \times 4) & 1 \times 5 + (-3) \times 3 \\ 2 \times 2 + 5 \times 4 & 2 \times 5 + 5 \times 3 \end{pmatrix}$$

$$= \begin{pmatrix} 2 - 12 & 5 - 9 \\ 4 + 20 & 10 + 15 \end{pmatrix}$$

$$= \begin{pmatrix} -10 & -4 \\ 24 & 25 \end{pmatrix} \dots\dots (2)$$

\therefore From (1) & (2) $AB \neq BA$

5. Given that $A = \begin{pmatrix} 1 & 3 \\ 5 & -1 \end{pmatrix}$, $B = \begin{pmatrix} 1 & -1 & 2 \\ 3 & 5 & 2 \end{pmatrix}$,

$$C = \begin{pmatrix} 1 & 3 & 2 \\ -4 & 1 & 3 \end{pmatrix} \text{ verify that}$$

$$A(B + C) = AB + AC$$

Solution :

Given

$$A = \begin{pmatrix} 1 & 3 \\ 5 & -1 \end{pmatrix}, B = \begin{pmatrix} 1 & -1 & 2 \\ 3 & 5 & 2 \end{pmatrix}, C = \begin{pmatrix} 1 & 3 & 2 \\ -4 & 1 & 3 \end{pmatrix}$$

To verify : $A(B + C) = AB + AC$

LHS : $A(B + C)$

$$\begin{aligned}
 &= \begin{pmatrix} 1 & 3 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} 2 & 2 & 4 \\ -1 & 6 & 5 \end{pmatrix} \\
 &= \begin{pmatrix} 1 \times 2 + 3 \times (-1) & 1 \times 2 + 3 \times 6 & 1 \times 4 + 3 \times 5 \\ 5 \times 2 + (-1) \times (-1) & 5 \times 2 - 1 \times 6 & 5 \times 4 - 1 \times 5 \end{pmatrix} \\
 &= \begin{pmatrix} 2-3 & 2+18 & 4+15 \\ 10+1 & 10-6 & 20-5 \end{pmatrix} \\
 &= \begin{pmatrix} -1 & 20 & 19 \\ 11 & 4 & 15 \end{pmatrix} \quad \dots\dots\dots (1)
 \end{aligned}$$

RHS : AB + AC

$$\begin{aligned}
 &= \begin{pmatrix} 1 & 3 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} 1 & -1 & 2 \\ 3 & 5 & 2 \end{pmatrix} + \begin{pmatrix} 1 & 3 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} 1 & 3 & 2 \\ -4 & 1 & 3 \end{pmatrix} \\
 &= \begin{pmatrix} 1+9 & -1+15 & 2+6 \\ 5-3 & -5-5 & 10-2 \end{pmatrix} + \begin{pmatrix} 1-12 & 3+3 & 2+9 \\ 5+4 & 15-1 & 10-3 \end{pmatrix} \\
 &= \begin{pmatrix} 10 & 14 & 8 \\ 2 & -10 & 8 \end{pmatrix} + \begin{pmatrix} -11 & 6 & 11 \\ 9 & 14 & 7 \end{pmatrix} \\
 &= \begin{pmatrix} -1 & 20 & 19 \\ 11 & 4 & 15 \end{pmatrix} \quad \dots\dots\dots (2)
 \end{aligned}$$

\therefore From (1) & (2) LHS = RHS

6. Show that the matrices

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix}, B = \begin{pmatrix} 1 & -2 \\ -3 & 1 \end{pmatrix} \text{ satisfy}$$

commutative property $AB = BA$

Solution :

$$\text{Given } A = \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix}, B = \begin{pmatrix} 1 & -2 \\ -3 & 1 \end{pmatrix}$$

$$\begin{aligned}
 AB &= \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} 1 & -2 \\ -3 & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1-6 & -2+2 \\ 3-3 & -6+1 \end{pmatrix} \\
 &= \begin{pmatrix} -5 & 0 \\ 0 & -5 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 BA &= \begin{pmatrix} 1 & -2 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix} \\
 &= \begin{pmatrix} 1-6 & -2+2 \\ -3+3 & -6+1 \end{pmatrix} \\
 &= \begin{pmatrix} -5 & 0 \\ 0 & -5 \end{pmatrix}
 \end{aligned}$$

$$\therefore AB = BA$$

\therefore Commutative property is true.

$$7. \text{ Let } A = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}, B = \begin{pmatrix} 4 & 0 \\ 1 & 5 \end{pmatrix}, C = \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix}$$

Show that i) $A(BC) = (AB)C$

ii) $(A - B)C = AC - BC$

iii) $(A - B)^T = A^T - B^T$

Solution :

$$\text{Given } A = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}, B = \begin{pmatrix} 4 & 0 \\ 1 & 5 \end{pmatrix}, C = \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix}$$

i) To show : $A(BC) = (AB)C$

$$\begin{aligned}
 BC &= \begin{pmatrix} 4 & 0 \\ 1 & 5 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 8+0 & 0+0 \\ 2+5 & 0+10 \end{pmatrix} \\
 &= \begin{pmatrix} 8 & 0 \\ 7 & 10 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 A(BC) &= \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 8 & 0 \\ 7 & 10 \end{pmatrix} = \begin{pmatrix} 8+14 & 0+20 \\ 8+21 & 0+30 \end{pmatrix} \\
 &= \begin{pmatrix} 22 & 20 \\ 29 & 30 \end{pmatrix} \quad \dots\dots\dots (1)
 \end{aligned}$$

$$\begin{aligned}
 AB &= \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 4 & 0 \\ 1 & 5 \end{pmatrix} \\
 &= \begin{pmatrix} 4+2 & 0+10 \\ 4+3 & 0+15 \end{pmatrix} \\
 &= \begin{pmatrix} 6 & 10 \\ 7 & 15 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 (AB)C &= \begin{pmatrix} 6 & 10 \\ 7 & 15 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} \\
 &= \begin{pmatrix} 12+10 & 0+20 \\ 14+15 & 0+30 \end{pmatrix} \\
 &= \begin{pmatrix} 22 & 20 \\ 29 & 30 \end{pmatrix} \quad \dots\dots\dots (2)
 \end{aligned}$$

\therefore From (1) & (2) LHS = RHS

ii) To show $(A - B)C = AC - BC$

$$\begin{aligned}
 (A - B)C &= \begin{pmatrix} -3 & 2 \\ 0 & -2 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} \\
 &= \begin{pmatrix} -6+2 & 0+4 \\ 0-2 & 0-4 \end{pmatrix} \\
 &= \begin{pmatrix} -4 & 4 \\ 0 & -4 \end{pmatrix} \quad \dots\dots\dots (1)
 \end{aligned}$$

$$\begin{aligned}
 AC &= \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} \\
 &= \begin{pmatrix} 2+2 & 0+4 \\ 2+3 & 0+6 \end{pmatrix} \\
 &= \begin{pmatrix} 4 & 4 \\ 5 & 6 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 BC &= \begin{pmatrix} 4 & 0 \\ 1 & 5 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} \\
 &= \begin{pmatrix} 8+0 & 0+0 \\ 2+5 & 0+10 \end{pmatrix} \\
 &= \begin{pmatrix} 8 & 0 \\ 7 & 10 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 \therefore AC - BC &= \begin{pmatrix} 4 & 4 \\ 5 & 6 \end{pmatrix} - \begin{pmatrix} 8 & 0 \\ 7 & 10 \end{pmatrix} \\
 &= \begin{pmatrix} -4 & 4 \\ -2 & -4 \end{pmatrix} \quad \dots\dots\dots (2)
 \end{aligned}$$

\therefore From (1) & (2), LHS = RHS

iii) To show : $(A - B)^T = A^T - B^T$

$$(A - B)^T = \begin{pmatrix} -3 & 2 \\ 0 & -2 \end{pmatrix}^T = \begin{pmatrix} -3 & 0 \\ 2 & -2 \end{pmatrix} \quad \dots\dots\dots (1)$$

$$\begin{aligned}
 A^T - B^T &= \begin{pmatrix} 1 & 1 \\ 2 & 3 \end{pmatrix} - \begin{pmatrix} 4 & 1 \\ 0 & 5 \end{pmatrix} \\
 &= \begin{pmatrix} -3 & 0 \\ 2 & -2 \end{pmatrix} \quad \dots\dots\dots (2)
 \end{aligned}$$

\therefore From (1) & (2)

$$(A - B)^T = A^T - B^T$$

8. If $A = \begin{pmatrix} \cos\theta & 0 \\ 0 & \cos\theta \end{pmatrix}$, $B = \begin{pmatrix} \sin\theta & 0 \\ 0 & \sin\theta \end{pmatrix}$
then show that $A^2 + B^2 = I$.

Solution :

$$\text{Given } A = \begin{pmatrix} \cos\theta & 0 \\ 0 & \cos\theta \end{pmatrix}, B = \begin{pmatrix} \sin\theta & 0 \\ 0 & \sin\theta \end{pmatrix}$$

To show : $A^2 + B^2 = I$

$$\begin{aligned}
 A^2 &= A \cdot A \\
 &= \begin{pmatrix} \cos\theta & 0 \\ 0 & \cos\theta \end{pmatrix} \begin{pmatrix} \cos\theta & 0 \\ 0 & \cos\theta \end{pmatrix} \\
 &= \begin{pmatrix} \cos^2\theta + 0 & 0 + 0 \\ 0 + 0 & 0 + \cos^2\theta \end{pmatrix} \\
 &= \begin{pmatrix} \cos^2\theta & 0 \\ 0 & \cos^2\theta \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 B^2 &= \begin{pmatrix} \sin\theta & 0 \\ 0 & \sin\theta \end{pmatrix} \begin{pmatrix} \sin\theta & 0 \\ 0 & \sin\theta \end{pmatrix} \\
 &= \begin{pmatrix} \sin^2\theta + 0 & 0 + 0 \\ 0 + 0 & 0 + \sin^2\theta \end{pmatrix} \\
 &= \begin{pmatrix} \sin^2\theta & 0 \\ 0 & \sin^2\theta \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}\therefore A^2 + B^2 &= \begin{pmatrix} \cos^2 \theta + \sin^2 \theta & 0 \\ 0 & \cos^2 \theta + \sin^2 \theta \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= I\end{aligned}$$

Hence proved.

9. If $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$ prove that $AA^T = I$

Solution :

Given $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$

To prove : $A.A^T = I$

LHS :

$$\begin{aligned}A.A^T &= \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} \cos^2 \theta + \sin^2 \theta & -\cos \theta \sin \theta + \sin \theta \cos \theta \\ -\sin \theta \cos \theta + \cos \theta \sin \theta & \sin^2 \theta + \cos^2 \theta \end{pmatrix} \\ &= I\end{aligned}$$

Hence proved.

10. Verify that $A^2 = I$ when $A = \begin{pmatrix} 5 & -4 \\ 6 & -5 \end{pmatrix}$

Solution :

Given $A = \begin{pmatrix} 5 & -4 \\ 6 & -5 \end{pmatrix}$

To prove : $A^2 = I$

$$\begin{aligned}A^2 &= A.A = \begin{pmatrix} 5 & -4 \\ 6 & -5 \end{pmatrix} \begin{pmatrix} 5 & -4 \\ 6 & -5 \end{pmatrix} \\ &= \begin{pmatrix} 25 - 24 & -20 + 20 \\ 30 - 30 & -24 + 25 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= I\end{aligned}$$

Hence proved.

11. If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ show that $A^2 - (a+d)A = (bc-ad)I$.

Solution :

Given $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

To prove : $A^2 - (a+d)A = (bc-ad)I$

$$\begin{aligned}A^2 &= A.A \\ &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{pmatrix}\end{aligned}$$

$$\begin{aligned}(a+d)A &= (a+d) \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\ &= \begin{pmatrix} a^2 + ad & ab + bd \\ ca + cd & ad + d^2 \end{pmatrix}\end{aligned}$$

$$\begin{aligned}A^2 - (a+d)A &= \begin{pmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{pmatrix} - \begin{pmatrix} a^2 + ad & ab + bd \\ ca + cd & ad + d^2 \end{pmatrix} \\ &= \begin{pmatrix} bc + ad & 0 \\ 0 & bc - ad \end{pmatrix} \\ &= (bc - ad) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= (bc - ad)I \\ &= \text{RHS}\end{aligned}$$

Hence proved.

12. If $A = \begin{pmatrix} 5 & 2 & 9 \\ 1 & 2 & 8 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 7 \\ 1 & 2 \\ 5 & -1 \end{pmatrix}$ verify
that $(AB)^T = B^T A^T$

Solution :

Given $A = \begin{pmatrix} 5 & 2 & 9 \\ 1 & 2 & 8 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 7 \\ 1 & 2 \\ 5 & -1 \end{pmatrix}$

To verify : $(AB)^T = B^T A^T$

$$\begin{aligned} AB &= \begin{pmatrix} 5 & 2 & 9 \\ 1 & 2 & 8 \end{pmatrix} \begin{pmatrix} 1 & 7 \\ 1 & 2 \\ 5 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 5 \times 1 + 2 \times 1 + 9 \times 5 & 5 \times 7 + 2 \times 2 + 9 \times -1 \\ 1 \times 1 + 2 \times 1 + 8 \times 5 & 1 \times 7 + 2 \times 2 + 8 \times -1 \end{pmatrix} \\ &= \begin{pmatrix} 5 + 2 + 45 & 35 + 4 - 9 \\ 1 + 2 + 40 & 7 + 4 - 8 \end{pmatrix} \\ &= \begin{pmatrix} 52 & 30 \\ 43 & 3 \end{pmatrix} \end{aligned}$$

$$\therefore (AB)^T = \begin{pmatrix} 52 & 43 \\ 30 & 3 \end{pmatrix} \quad \dots\dots\dots (1)$$

$$\begin{aligned} B^T A^T &= \begin{pmatrix} 1 & 1 & 5 \\ 7 & 2 & -1 \end{pmatrix} \begin{pmatrix} 5 & 1 \\ 2 & 2 \\ 9 & 8 \end{pmatrix} \\ &= \begin{pmatrix} 5 + 2 + 45 & 1 + 2 + 40 \\ 35 + 4 - 9 & 7 + 4 - 8 \end{pmatrix} \\ &= \begin{pmatrix} 52 & 43 \\ 30 & 3 \end{pmatrix} \quad \dots\dots\dots (2) \end{aligned}$$

\therefore From (1) & (2), $(AB)^T = B^T A^T$

Hence proved.

13. If $A = \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix}$ show that $A^2 - 5A + 7I_2 = 0$

Solution :

Given $A = \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix}$

To verify : $A^2 - 5A + 7I_2 = 0$

$$\begin{aligned} A^2 &= \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 9 - 1 & 3 + 2 \\ -3 - 2 & -1 + 4 \end{pmatrix} \\ &= \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \therefore A^2 - 5A + 7I_2 &= \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} - \begin{pmatrix} 15 & 5 \\ -5 & 10 \end{pmatrix} + 7 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} - \begin{pmatrix} 15 & 5 \\ -5 & 10 \end{pmatrix} + \begin{pmatrix} 7 & 0 \\ 0 & 7 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \\ &= 0 \end{aligned}$$

EXERCISE 3.19

Multiple Choice Questions :

1. A system of three linear equations in three variables is inconsistent if their planes

- (1) intersect only at a point
- (2) intersect in a line
- (3) coincides with each other
- (4) do not intersect

Hint :

Ans : (4)

System of equations is inconsistent if their planes do not intersect.

2. The solution of the system $x + y - 3z = -6$, $-7y + 7z = 7$, $3z = 9$ is

(1) $x = 1, y = 2, z = 3$

(2) $x = -1, y = 2, z = 3$

(3) $x = -1, y = -2, z = 3$

(4) $x = 1, y = 2, z = 3$

Hint :

Ans : (1)

(3) $\Rightarrow 3z = 9 \Rightarrow z = 3$

(2) $\Rightarrow -y + z = 1 \Rightarrow -y + 3 = 1$

$\Rightarrow -y = -2$

$y = 2$

(1) $\Rightarrow x + y - 3z = -6$

$\Rightarrow x + 2 - 9 = -6$

$\Rightarrow x = 7 - 6 = 1$

2

$x^2 - kx - 6$ then the value of k is

(1) 3 (2) 5 (3) 6 (4) 8

Hint :

Ans : (2)

$x^2 - 2x - 24 = (x - 6)(x + 4)$

$x^2 - kx - 6 = (x - 6)(x + 1)$

($\because x + 1$ is the only possible factor)

$= x^2 - 5x - 6$

$\therefore k = 5$

4. $\frac{3y-3}{y} \div \frac{7y-7}{3y^2}$ is

(1) $\frac{9y}{7}$

(2) $\frac{9y^3}{(21y-21)}$

(3) $\frac{21y^2-42y+21}{3y^3}$

(4) $\frac{7(y^2-2y+1)}{y^2}$

Hint :

Ans : (1)

$$\begin{aligned} &= \frac{3y-3}{y} \div \frac{7y-7}{3y^2} \\ &= \frac{3(y-1)}{y} \div \frac{7(y-1)}{3y^2} \\ &= \frac{9y}{7} \end{aligned}$$

5. $y^2 + \frac{1}{y^2}$ is not equal to

(1) $\frac{y^4+1}{y^2}$

(2) $\left(y + \frac{1}{y}\right)^2$

(3) $\left(y + \frac{1}{y}\right)^2 + 2$

(4) $\left(y + \frac{1}{y}\right)^2 - 2$

Hint :

Ans : (2)

$$y^2 + \frac{1}{y^2} \neq \left(y + \frac{1}{y}\right)^2$$

6. $\frac{x}{x^2-25} - \frac{8}{x^2+6x+5}$ gives

(1) $\frac{x^2-7x+40}{(x-5)(x+5)}$

(2) $\frac{x^2+7x+40}{(x-5)(x+5)(x+1)}$

(3) $\frac{x^2-7x+40}{(x^2-25)(x+1)}$

(4) $\frac{x^2+10}{(x^2-25)(x+1)}$

Hint :

Ans : (3)

$$\begin{aligned} &= \frac{x}{x^2-25} - \frac{8}{x^2+6x+5} \\ &= \frac{x}{(x+5)(x-5)} - \frac{8}{(x+5)(x+1)} \\ &= \frac{x(x+1) - 8(x-5)}{(x+5)(x-5)(x+1)} \\ &= \frac{x^2+x-8x+40}{(x+5)(x-5)(x+1)} \\ &= \frac{x^2-7x+40}{(x^2-25)(x+1)} \end{aligned}$$

7. The square root of $\frac{256x^8y^4z^{10}}{25x^6y^6z^6}$ is equal to

- (1) $\frac{16}{5} \left| \frac{x^2z^4}{y^2} \right|$ (2) $16 \left| \frac{y^2}{x^2z^4} \right|$
 (3) $\frac{16}{5} \left| \frac{y}{xz^2} \right|$ (4) $\frac{16}{5} \left| \frac{xz^2}{y} \right|$

Hint : Ans : (4)

$$\begin{aligned}
 &= \sqrt{\frac{256x^8y^4z^{10}}{25x^6y^6z^6}} \\
 &= \frac{16}{5} \left| \frac{x^4y^2z^5}{x^3y^3z^3} \right| \\
 &= \frac{16}{5} \left| \frac{xz^2}{y} \right|
 \end{aligned}$$

to make $x^2 + 64$ a perfect square

- (1) $4x^2$ (2) $16x^2$
 (3) $8x^2$ (4) $-8x^2$

Hint : Ans : (2)

$$\begin{aligned}
 &= x^4 + 64 \\
 &= (x^2)^2 + 8^2 \\
 &= (x^2)^2 + 8^2 + 2(x^2)(8) \\
 &= (x^2 + 8)^2, \text{ perfect square} \\
 &\therefore 16x^2 \text{ should be added}
 \end{aligned}$$

9. The solution of $(2x - 1)^2 = 9$ is equal to

- (1) -1 (2) 2
 (3) -1, 2 (4) None of these

Hint : Ans : (3)

$$\begin{aligned}
 (2x - 1)^2 &= 9 \Rightarrow 2x - 1 = \pm 3 \\
 &\Rightarrow 2x = 4, 2x = -2 \\
 x &= 2, x = -1
 \end{aligned}$$

10. The values of a and b if $4x^4 - 24x^3 + 76x^2 + ax + b$ is a perfect square are

- (1) 100, 120 (2) 10, 12
 (3) -120, 100 (4) 12, 10

Hint : Ans : (3)

$2x^2$	$4x^4 - 24x^3 + 76x^2 + ax + b$
$4x^2 - 6x$	$4x^4$ <hr style="border: 0; border-top: 1px solid black;"/> $-24x^3 + 76x^2$
$4x^2 - 12x + 10$	$-24x^3 + 36x^2$ <hr style="border: 0; border-top: 1px solid black;"/> $40x^2 + ax + b$ $40x^2 - 120x + 100$ $(+) \quad (-)$ <hr style="border: 0; border-top: 1px solid black;"/> 0

(\therefore Perfect square)

$$a + 120 = 0, \quad b - 100 = 0$$

$$a = -120 \quad b = 100$$

11. If the roots of the equation $qx^2 + px + r = 0$ are the squares of the roots of the equation $qx^2 + px + r = 0$, then q, p, r are in _____

- (1) A.P (2) G.P
 (3) Both A.P and G.P (4) none of these

Hint : Ans : (2)

Roots of $qx^2 + px + r = 0$ are squares of roots of $qx^2 + px + r = 0$

$$\Rightarrow \alpha^2 + \beta^2 = \frac{-p^2}{q^2} \quad \left| \quad \alpha^2 \beta^2 = \frac{r^2}{q^2} \right.$$

$$\text{and } \alpha + \beta = \frac{-p}{q}, \alpha\beta = \frac{r}{q}$$

$$\Rightarrow (\alpha + \beta)^2 - 2\alpha\beta = \frac{-p^2}{q^2}$$

$$\Rightarrow \frac{p^2}{q^2} - \frac{2r}{q} = \frac{-p^2}{q^2}$$

$$\Rightarrow \frac{2r}{q} = \frac{2p^2}{q^2}$$

$$\Rightarrow r = \frac{2p^2}{q^2}$$

$$\Rightarrow p^2 = qr$$

$\therefore q, p, r$ are in G.P.

12. Graph of a linear polynomial is a

- (1) straight line (2) circle
(3) parabola (4) hyperbola

Hint : **Ans : (1)**

line.

13. The number of points of intersection of the quadratic polynomial $x^2 + 4x + 4$ with the X axis is

- (1) 0 (2) 1 (3) 0 or 1 (4) 2

Hint : **Ans : (2)**

$$(x + 2)^2 = 0$$

$$\Rightarrow x + 2 = 0$$

$$\Rightarrow x = -2$$

\therefore The polynomial will meet x - axis at $(-2, 0)$

No. of points of intersection = 1.

14. For the given matrix $A = \begin{pmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \\ 9 & 11 & 13 & 15 \end{pmatrix}$

the order of the matrix A^T is

- (1) 2×3 (2) 3×2
(3) 3×4 (4) 4×3

Hint : **Ans : (3)**

A has 3 rows & 4 columns

\therefore A is of order 3×4

15. If A is a 2×3 matrix and B is a 3×4 matrix, how many columns does AB have?

- (1) 3 (2) 4 (3) 2 (4) 5

Hint : **Ans : (2)**

$A \rightarrow 2 \times 3, \quad B \rightarrow 3 \times 4$

\therefore AB is of order 2×4

\therefore No. of columns of $A \times B$ is 4.

16. If number of columns and rows are not equal in a matrix then it is said to be a

- (1) diagonal matrix
(2) rectangular matrix
(3) square matrix
(4) identity matrix

Hint : **Ans : (2)**

No. of rows \neq No. of columns

\Rightarrow Matrix is said to be rectangular

17. Transpose of a column matrix is

- (1) unit matrix (2) diagonal matrix
(3) column matrix (4) row matrix

Hint : **Ans : (4)**

Transpose of a column matrix is a row matrix

$$\text{Ex : If } A = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \quad A^T = (1 \quad 2 \quad 3)$$

18. Find the matrix X if $2X + \begin{pmatrix} 1 & 3 \\ 5 & 7 \end{pmatrix} = \begin{pmatrix} 5 & 7 \\ 9 & 5 \end{pmatrix}$

(1) $\begin{pmatrix} -2 & -2 \\ 2 & -1 \end{pmatrix}$

(2) $\begin{pmatrix} 2 & 2 \\ 2 & -1 \end{pmatrix}$

(3) $\begin{pmatrix} 1 & 2 \\ 2 & 2 \end{pmatrix}$

(4) $\begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix}$

Hint :

Ans : (2)

$$2X + \begin{pmatrix} 1 & 3 \\ 5 & 7 \end{pmatrix} = \begin{pmatrix} 5 & 7 \\ 9 & 5 \end{pmatrix}$$

$$\Rightarrow 2X = \begin{pmatrix} 4 & 4 \\ 4 & -2 \end{pmatrix}$$

$$\Rightarrow X = \begin{pmatrix} 2 & 2 \\ 2 & -1 \end{pmatrix}$$

19. Which of the following can be calculated from the given matrices

$$A = \begin{pmatrix} 3 & 4 \\ 5 & 6 \end{pmatrix}, B = \begin{pmatrix} 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

(i) A^2 (ii) B^2 (iii) AB (iv) BA

(1) (i) and (ii) only (2) (ii) and (iii) only

(3) (ii) and (iv) only (4) all of these

Hint :

Ans : (3)

A is of order 3×2

i) A^2 is not possible [$(3 \times 2)(3 \times 2)$

is not possible]

B is of order 3×3

ii) B^2 is possible

iii) AB is not defined (no. of columns in A \neq no. of rows in B)

iv) BA is of order 3×3

$\therefore BA$ is possible.

20. If $A = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 0 \\ 2 & -1 \\ 0 & 2 \end{pmatrix}$
and $C = \begin{pmatrix} 0 & 1 \\ -2 & 5 \end{pmatrix}$

Which of the following statements are correct?

(i) $AB + C = \begin{pmatrix} 5 & 5 \\ 5 & 5 \end{pmatrix}$

(ii) $BC = \begin{pmatrix} 0 & 1 \\ 2 & -3 \\ -4 & 10 \end{pmatrix}$

(iii) $BA + C = \begin{pmatrix} 2 & 5 \\ 3 & 0 \end{pmatrix}$

(iv) $(AB)C = \begin{pmatrix} -8 & 20 \\ -8 & 13 \end{pmatrix}$

(1) (i) and (ii) only (2) (ii) and (iii) only

(3) (iii) and (iv) only (4) all of these

Hint :

i) $AB + C = \begin{pmatrix} 1+4+0 & 0-2+6 \\ 3+4+0 & 0-2+2 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ -2 & 5 \end{pmatrix}$

$$= \begin{pmatrix} 5 & 4 \\ 7 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ -2 & 5 \end{pmatrix}$$

$$= \begin{pmatrix} 5 & 5 \\ 5 & 5 \end{pmatrix} \quad \text{Correct}$$

ii) $B \rightarrow 3 \times 2$, $C \rightarrow 2 \times 2$

$\therefore BC$ is of order 3×2

$$BC = \begin{pmatrix} 0+0 & 1+0 \\ 0+2 & 2-5 \\ 0-4 & 0+10 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 2 & -3 \\ -4 & 10 \end{pmatrix} \quad \text{Correct}$$

$$\text{iii) } BA + C \neq \begin{pmatrix} 2 & 5 \\ 3 & 0 \end{pmatrix}, (\because BA \text{ is of order } 3 \times 3)$$

$$\text{iv) } AB = \begin{pmatrix} 5 & 4 \\ 7 & 0 \end{pmatrix}$$

$$\begin{aligned} \therefore ABC &= \begin{pmatrix} 5 & 4 \\ 7 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -2 & 5 \end{pmatrix} \\ &= \begin{pmatrix} 0-8 & 5+20 \\ 0+0 & 7+0 \end{pmatrix} \\ &= \begin{pmatrix} -8 & 25 \\ 0 & 7 \end{pmatrix} \neq \begin{pmatrix} -8 & 20 \\ -8 & 13 \end{pmatrix} \end{aligned}$$

\therefore (i) & (ii) only are correct

UNIT EXERCISE - 3

1. Solve

$$\frac{1}{3}$$

Solution :

Given

$$\frac{1}{3}(x + y - 5) = y - z = 2x - 11 = 9 - (x + 2z)$$

$$\Rightarrow \frac{1}{3}(x + y - 5) = y - z$$

$$\Rightarrow x + y - 5 = 3y - 3z$$

$$\Rightarrow x - 2y + 3z = 5 \quad \dots\dots\dots (1)$$

$$\Rightarrow y - z = 2x - 11$$

$$\Rightarrow 2x - y + z = 11 \quad \dots\dots\dots (2)$$

Also, $2x - 11 = 9 - x - 2z$

$$3x + 2z = 20 \quad \dots\dots\dots (3)$$

$$(1) \Rightarrow x - 2y + 3z = 5$$

$$(2) \Rightarrow \times 2 \quad 4x - 2y + 2z = 22$$

$$\begin{array}{r} (-) \quad (+) \quad (-) \quad (-) \\ \hline -3x \qquad \qquad + z = -17 \quad \dots\dots\dots (4) \end{array}$$

Solving (3) & (4)

$$3x + 2z = 20$$

$$-3x + z = -17$$

$$\begin{array}{r} (-) \quad (+) \quad (-) \quad (-) \\ \hline 3z = 3 \\ z = 1 \end{array}$$

Sub $z = 1$ in (4)

$$-3x + 1 = -17$$

$$-3x = -18$$

$$x = 6$$

Sub $x = 6, z = 1$ in (1)

$$6 - 2y + 3 = 5$$

$$\Rightarrow -2y = -4$$

$$y = 2$$

$$\therefore x = 6, y = 2, z = 1.$$

2. One hundred and fifty students are admitted to a school. They are distributed over three sections A, B and C. If 6 students are shifted from section A to section C, the sections will have equal number of students. If 4 times of students of section C exceeds the number of students of section A by the number of students in section B, find the number of students in the three sections.

Solution :

Let the number of students in 3 sections.

A, B, C respectively be x, y, z

$$\therefore \text{By data given, } x + y + z = 150 \quad \text{--- (1)}$$

$$x - 6 = z + 6$$

$$x - z = 12 \quad \text{--- (2)}$$

Also, given $4z = x + y$

$$\Rightarrow x + y - 4z = 0 \quad \text{--- (3)}$$

Solving (1) & (2)

$$\begin{array}{r} x + y + z = 150 \\ x + y - 4z = 0 \\ \hline \text{Subtracting,} \quad 5z = 150 \\ z = 30 \end{array}$$

Sub. $z = 30$ in (2)

$$x - 30 = 12$$

$$x = 42$$

Sub. $x = 42, z = 30$ in (1)

$$42 + y + 30 = 150$$

$$\Rightarrow y = 150 - 72 \\ = 78$$

\therefore No. of students in Section A = 42

No. of students in Section B = 78

No. of students in Section C = 30

- 3. In a three-digit number, when the tens and the hundreds digit are interchanged the new number is 54 more than three times the original number. If 198 is added to the number, the digits are reversed. The tens digit exceeds the hundreds digit by twice as that of the tens digit exceeds the unit digit. Find the original number.**

Solution :

Let the original 3 digit number be

$$100x + 10y + z.$$

By data given

$$\begin{aligned} 100y + 10x + z &= 3(100x + 10y + z) + 54 \\ \Rightarrow 100y + 10x + z &= 300x + 30y + 3z + 54 \\ \Rightarrow 290x - 70y + 2z &= -54 \\ \Rightarrow 145x - 35y + z &= -27 \quad \text{--- (1)} \end{aligned}$$

When 198 is added to the number of digits are reversed

$$\Rightarrow (100x + 10y + z) + 198 = 100z + 10y + x$$

$$\Rightarrow 99x - 99z = -198$$

$$\Rightarrow x - z = -2$$

$$\Rightarrow x = z - 2 \quad \text{--- (2)}$$

10's digit exceeds 100's digit by twice as that of 10's digits exceed the unit's digit

$$\Rightarrow y - x = 2(y - z)$$

$$\Rightarrow y - x = 2y - 2z$$

$$\Rightarrow -x - y = -2z$$

$$\Rightarrow x + y = 2z$$

$$\Rightarrow z - 2 + y = 2z \quad (\text{from (2)})$$

$$y = 2z - z + 2$$

$$y = z + 2 \quad \text{--- (3)}$$

$$145(z - 2) - 35(z + 2) + z = -27$$

$$\Rightarrow 145z - 290 - 35z - 70 + z = -27$$

$$111z = 333$$

$$z = 3$$

$$\text{Sub. } z = 3 \text{ in (2)} \Rightarrow x = 3 - 2 = 1$$

$$\text{Sub. } z = 3 \text{ in (3)} \Rightarrow y = 3 + 2 = 5$$

$$x = 1, y = 5, z = 3$$

\therefore The original number is $100x + 10y + z$

$$= 100(1) + 10(5) + 3$$

$$= 100 + 50 + 3$$

$$= 153$$

4. Find the least common multiple of $xy(k^2 + 1) + k(x^2 + y^2)$ and $xy(k^2 - 1) + k(x^2 - y^2)$.

Solution :

$$\begin{aligned}
 &xy(k^2 + 1) + k(x^2 + y^2) \\
 &= kx^2 + (k^2 + 1)xy + ky^2 \\
 &= kx^2 + k^2xy + xy + ky^2 \\
 &= kx(x + ky) + y(x + ky) \\
 &= (kx + y)(x + ky) \\
 &xy(k^2 - 1) + k(x^2 - y^2) \\
 &= kx^2 + (k^2 - 1)xy - ky^2 \\
 &= kx^2 + k^2xy - xy - ky^2 \\
 &= kx(x + ky) - y(x + ky) \\
 &= (x + ky)(kx - y) \\
 &\text{LCM} = (x + ky)(kx + y)(kx - y) \\
 &= (x + ky)(k^2x^2 - y^2)
 \end{aligned}$$

5. Find the GCD of the following by division algorithm $2x^4 + 13x^3 + 27x^2 + 23x + 7$, $x^3 + 3x^2 + 3x + 1$, $x^2 + 2x + 1$

Solution :

$$\begin{array}{r}
 x + 1 \\
 x^2 + 2x + 1 \overline{) x^3 + 3x^2 + 3x + 1} \\
 \underline{x^3 + 2x^2 + x} \\
 x^2 + 2x + 1 \\
 \underline{x^2 + 2x + 1} \\
 0
 \end{array}$$

$$\begin{array}{r}
 2x^2 + 9x + 7 \\
 x^2 + 2x + 1 \overline{) 2x^4 + 13x^3 + 27x^2 + 23x + 7} \\
 \underline{2x^4 + 4x^3 + 2x^2} \\
 9x^3 + 25x^2 + 23x \\
 \underline{9x^3 + 18x^2 + 9x} \\
 7x^2 + 14x + 7 \\
 \underline{7x^2 + 14x + 7} \\
 0
 \end{array}$$

$$\therefore \text{G.C.D.} = x^2 + 2x + 1$$

6. Reduce the given Rational expressions to its lowest form

$$(i) \frac{x^{3a} - 8}{x^2a + 2xa + 4} \quad (ii) \frac{10x^3 - 25x^2 + 4x - 10}{-4 - 10x^2}$$

Solution :

$$\begin{aligned}
 (i) &\frac{x^{3a} - 8}{x^2a + 2xa + 4} \\
 &= \frac{(x^a)^3 - 2^3}{x^{2a} + 2x^a + 4} \\
 &= \frac{(x^a - 2)(x^{2a} + 2x^a + 4)}{x^{2a} + 2x^a + 4} \\
 &= x^a - 2 \\
 (ii) &\frac{10x^3 - 25x^2 + 4x - 10}{-4 - 10x^2} \\
 &= \frac{5x^2(2x - 5) + 2(2x - 5)}{-2(2 + 5x)} \\
 &= \frac{(5x^2 + 2)(2x - 5)}{-2(2 + 5x)} \\
 &= \frac{(2x - 5)}{-2} \\
 &= -x + \frac{5}{2}
 \end{aligned}$$

7. Simplify

$$\frac{\frac{1}{p} + \frac{1}{q+r}}{\frac{1}{p} - \frac{1}{q+r}} \times \left(1 + \frac{q^2 + r^2 - p^2}{2qr} \right)$$

Solution :

$$= \frac{\frac{1}{p} + \frac{1}{q+r}}{\frac{1}{p} - \frac{1}{q+r}} \times \left(1 + \frac{q^2 + r^2 - p^2}{2qr} \right)$$

$$\begin{aligned}
 &= \frac{\frac{q+r+p}{q+r-p} \times \left(\frac{q^2 + r^2 - p^2 + 2qr}{2qr} \right)}{p(q+r)} \\
 &= \frac{q+r+p}{q+r-p} \times \frac{(q+r)^2 - p^2}{2qr} \\
 &= \frac{q+r+p}{q+r-p} \times \frac{(q+r+p)(q+r-p)}{2qr} \\
 &= \frac{(p+q+r)^2}{2qr}
 \end{aligned}$$

8. Arul, Ravi and Ram working together can clean a store in 6 hours. Working alone, Ravi takes twice as long to clean the store as Arul does. Ram needs three times as long as Arul does. How long would it take each if they are working

Solution :

Let x, y, z be the working speed of Arul, Ravi and Ram respectively & W be the total work done.

$$x + y + z = \frac{W}{6} \quad \text{--- (1)}$$

By data given,

- * Ravi takes twice as Arul does

$$\Rightarrow \frac{W}{y} = 2 \left(\frac{W}{x} \right)$$

$$\Rightarrow \frac{1}{y} = \frac{2}{x}$$

$$\Rightarrow y = \frac{x}{2}$$

- * Ram takes thrice as Arul does

$$\Rightarrow \frac{W}{z} = 3 \left(\frac{W}{x} \right)$$

$$\Rightarrow \frac{1}{z} = \frac{3}{x}$$

$$\Rightarrow z = \frac{x}{3}$$

$$\therefore (1) \Rightarrow x + \frac{x}{2} + \frac{x}{3} = \frac{W}{6}$$

$$\Rightarrow \frac{6x + 3x + 2x}{6} = \frac{W}{6}$$

$$\Rightarrow W = 11x$$

$$\therefore x = \frac{W}{11}$$

$$\therefore y = \frac{x}{2} \Rightarrow y = \frac{W/11}{2} = \frac{W}{22} \text{ and}$$

$$z = \frac{x}{3} \Rightarrow z = \frac{W/11}{3} = \frac{W}{33}$$

$$\therefore \text{Arul alone does} = \frac{W}{x} = \frac{W}{W/11} = 11 \text{ hours}$$

$$\text{Ravi alone does} = \frac{W}{y} = \frac{W}{W/22} = 22 \text{ hours}$$

$$\text{Ram alone does} = \frac{W}{z} = \frac{W}{W/33} = 33 \text{ hours}$$

- 9 Find the square root of $289x^4 - 612x^3 + 970x^2 - 684x + 361$.

Solution :

$17x^2$	$17x^2 - 18x + 19$
$17x^2$	$289x^4 - 612x^3 + 970x^2 - 684x + 361$
	$289x^4$
$34x^2 - 18x$	$- 612x^3 + 970x^2$
	$- 612x^3 + 324x^2$
$34x^2 - 36x + 19$	$646x^2 - 684x + 361$
	$646x^2 - 684x + 361$
	0

$$\begin{aligned}
 \therefore \sqrt{289x^4 - 612x^3 + 970x^2 - 684x + 361} \\
 = |17x^2 - 18x + 19|
 \end{aligned}$$

10. Solve $\sqrt{y+1} + \sqrt{2y-5} = 3$

Solution :

Given $\sqrt{y+1} + \sqrt{2y-5} = 3$

$$\Rightarrow \sqrt{y+1} = 3 - \sqrt{2y-5}$$

Squaring on both sides

$$y+1 = 9 + 2y - 5 - 6\sqrt{2y-5}$$

$$\Rightarrow -y - 3 = -6\sqrt{2y-5}$$

$$\Rightarrow y + 3 = 6\sqrt{2y-5}$$

$$\Rightarrow (y+3)^2 = 36(2y-5)$$

$$\Rightarrow y^2 + 6y + 9 = 72y - 180$$

$$\Rightarrow y^2 - 66y + 189 = 0$$

$$\Rightarrow (y-63)(y-3) = 0$$

$$y = 63, 3$$

11. A boat takes 1.6 hours longer to go 36 kms up a river than down the river. If the speed of the water current is 4 km per hr, what is the speed of the boat in still water?

Solution :

Let the speed of the boat in still water be x km/hr.

Distance = 36 kms, Time difference = 1.6 hrs.

\therefore By data given,

$$\frac{36}{x-4} - \frac{36}{x+4} = \frac{8}{5} \quad (\because 1.6 \text{ hrs} = \frac{8}{5} \text{ hrs})$$

$$\Rightarrow 36 \left(\frac{1}{x-4} - \frac{1}{x+4} \right) = \frac{8}{5}$$

$$\Rightarrow \frac{x+4-x+4}{x^2-16} = \frac{8}{36 \times 5}$$

$$\Rightarrow \frac{8}{x^2-16} = \frac{8}{180}$$

$$\Rightarrow x^2 - 16 = 180$$

$$\Rightarrow x^2 = 196$$

$$\therefore x = 14$$

\therefore Speed of boat in still water = 14km / hr.

12. Is it possible to design a rectangular park of perimeter 320 m and area 4800 m²? If so find its length and breadth.

Solution :

Given perimeter of a rectangular park = 320 m

$$\text{Area} = 4800 \text{ m}^2$$

$$\therefore 2(l+b) = 320, \quad lb = 4800$$

$$\Rightarrow l+b = 160, \quad lb = 4800 \quad \dots\dots (2)$$

$$\therefore b + 160 - 2 \quad \dots\dots (1)$$

Sub (1) in (2)

$$l(160-l) = 4800$$

$$\Rightarrow 160l - l^2 = 4800$$

$$\Rightarrow l^2 - 160l + 4800 = 0$$

$$\Rightarrow (l-120)(l-40) = 0$$

$$\therefore l = 120, l = 40$$

$$\therefore l = 120, (1) \Rightarrow b = 160 - 120$$

$$\therefore \text{Length} = 120 \text{ m}$$

$$\text{Breadth} = 40 \text{ m}$$

13. At t minutes past 2 pm, the time needed to 3 pm is 3 minutes less than $\frac{t^2}{4}$. Find t .

Solution :

Given time needed by the minutes hand show

$$\frac{t^2}{4} - 3.$$

\therefore As per the data given,

$$\frac{t^2}{4} - 3 = 60 - t$$

$$\Rightarrow t^2 - 12 = 240 - 4t$$

$$\Rightarrow t^2 + 4t - 252 = 0$$

$$\Rightarrow (t+18)(t-14) = 0$$

$$\therefore t = -18, t = 14$$

$$\therefore t = 14 \text{ min.}$$

- 14. The number of seats in a row is equal to the total number of rows in a hall. The total number of seats in the hall will increase by 375 if the number of rows is doubled and the number of seats in each row is reduced by 5. Find the number of rows in the hall at the beginning.**

Solution :

Let the number of rows be x .

\therefore Number of seats in each row = x

\therefore Total number of seats in the hall = x^2

\therefore By the data given,

$$2x \times (x - 5) = x^2 + 375$$

$$\Rightarrow 2x^2 - 10x = x^2 + 375$$

$$\Rightarrow x^2 - 10x - 375 = 0$$

$$\Rightarrow (x - 25)(x + 15) = 0$$

$$\therefore x = 25, -15$$

\therefore No. of rows at the beginning = 25.

- 15. If α and β are the roots of the polynomial $f(x) = x^2 - 2x + 3$, find the polynomial whose roots are (i) $\alpha + 2$, $\beta + 2$**

(ii) $\frac{\alpha-1}{\alpha+1}, \frac{\beta-1}{\beta+1}$

Solution :

Given α, β are the roots of $f(x) = x^2 - 2x + 3$

$$\alpha + \beta = 2, \alpha\beta = 3$$

- i) To find the polynomial whose roots are $\alpha + 2$, $\beta + 2$

$$\text{Sum} = \alpha + \beta + 4$$

$$= 2 + 4$$

$$= 6$$

$$\text{Product} = (\alpha + 2)(\beta + 2)$$

$$= \alpha\beta + 2\alpha + 2\beta + 4$$

$$= \alpha\beta + 2(\alpha + \beta) + 4$$

$$= 3 + 2(2) + 4$$

$$= 11$$

\therefore The required polynomial is $x^2 - 6x + 11$.

- ii) To find the polynomial whose roots are

$$\frac{\alpha-1}{\alpha+1}, \frac{\beta-1}{\beta+1}$$

$$\text{Sum} = \frac{\alpha-1}{\alpha+1} + \frac{\beta-1}{\beta+1}$$

$$= \frac{\alpha\beta + \alpha - \beta - 1 + \alpha\beta - \alpha + \beta - 1}{(\alpha+1)(\beta+1)}$$

$$= \frac{2\alpha\beta - 2}{\alpha\beta + \alpha + \beta + 1}$$

$$= \frac{2(3) - 2}{3 + 2 + 1} = \frac{4}{6} = \frac{2}{3}$$

$$\text{Product} = \frac{\alpha-1}{\alpha+1} \times \frac{\beta-1}{\beta+1}$$

$$= \frac{\alpha\beta - \alpha - \beta - 1}{(\alpha+1)(\beta+1)} = \frac{\alpha\beta - (\alpha + \beta) + 1}{(\alpha+1)(\beta+1)}$$

$$= \frac{3 - 2 + 1}{3 + 2 + 1}$$

$$= \frac{2}{6}$$

$$= \frac{1}{3}$$

\therefore The required polynomial is

$$x^2 - \frac{2}{3}x + \frac{1}{3} = \frac{3x^2 - 2x + 1}{3}$$

and the equation is $3x^2 - 2x + 1 = 0$.

- 16. If -4 is a root of the equation $x^2 + px - 4 = 0$ and if the equation $x^2 + px + q = 0$ has equal roots, find the values of p and q .**

Solution :

Given -4 is a root of $x^2 + px - 4 = 0$

$$\begin{array}{l|l} \therefore \alpha + \beta = -p, & \alpha\beta = -4 \\ -4 + 1 = -p & \Rightarrow -4 \times \beta = -4 \\ \therefore p = 3 & \therefore \beta = 1 \end{array}$$

Also $x^2 + px + q = 0$ has equal notes.

$$\therefore a = 1, b = p, c = q$$

$$\Rightarrow b^2 - 4ac = 0$$

$$\Rightarrow p^2 - 4q = 0$$

$$\Rightarrow q - 4q = 0$$

$$\therefore p = 3, q = \frac{9}{4}$$

- 17. Two farmers Senthil and Ravi cultivate three varieties of grains namely rice, wheat and ragi. If the sale (in ₹) of three varieties of grains by both the farmers in the month of April is given by the matrix.**

April sale in ₹

$$A = \begin{pmatrix} \text{rice} & \text{wheat} & \text{ragi} \\ 500 & 1000 & 1500 \\ 2500 & 1500 & 500 \end{pmatrix} \begin{matrix} \text{Senthil} \\ \text{Ravi} \end{matrix}$$

and the May month sale (in ₹) is exactly twice as that of the April month sale for each variety.

- (i) What is the average sales of the months April and May.
 (ii) If the sales continue to increase in the same way in the successive months, what will be sales in the month of August?

Solution :

Given sale of April month is

$$A = \begin{pmatrix} 500 & 1000 & 1500 \\ 2500 & 1500 & 500 \end{pmatrix}$$

\therefore By given data, sale of May month is

$$B = \begin{pmatrix} 1000 & 2000 & 3000 \\ 5000 & 3000 & 1000 \end{pmatrix}$$

- i) Average sales of April & May

$$= \begin{pmatrix} \frac{1500}{2} & \frac{3000}{2} & \frac{4500}{2} \\ \frac{7500}{2} & \frac{4500}{2} & \frac{1500}{2} \end{pmatrix} \\ = \begin{pmatrix} 750 & 1500 & 2250 \\ 3750 & 2250 & 750 \end{pmatrix}$$

- ii) Sales in the month of August

$$= 16 \begin{pmatrix} 500 & 1000 & 1500 \\ 2500 & 1500 & 500 \end{pmatrix} \\ = \begin{pmatrix} 8000 & 16000 & 24000 \\ 40000 & 24000 & 8000 \end{pmatrix}$$

18. If $\cos\theta \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} + \sin\theta \begin{pmatrix} x & -\cos\theta \\ \cos\theta & x \end{pmatrix} = I_2$

find x.

Solution :

Given

$$\cos\theta \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} + \sin\theta \begin{pmatrix} x & -\cos\theta \\ \cos\theta & x \end{pmatrix} = I_2$$

$$\Rightarrow \begin{pmatrix} \cos^2\theta & \cos\theta\sin\theta \\ -\cos\theta\sin\theta & \cos^2\theta \end{pmatrix} + \begin{pmatrix} x\sin\theta & -\sin\theta\cos\theta \\ \sin\theta\cos\theta & x\sin\theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} \cos^2\theta + x\sin\theta & 0 \\ 0 & \cos^2\theta + x\sin\theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow \cos^2\theta + x\sin\theta = 1$$

$$\Rightarrow x\sin\theta = 1 - \cos^2\theta$$

$$\Rightarrow x\sin\theta = \sin^2\theta$$

$$\Rightarrow x = \sin\theta$$

19. Given $A = \begin{pmatrix} p & 0 \\ 0 & 2 \end{pmatrix}$, $B = \begin{pmatrix} 0 & -q \\ 1 & 0 \end{pmatrix}$, $C = \begin{pmatrix} 2 & -2 \\ 2 & 2 \end{pmatrix}$
and if $BA = C^2$, find p and q .

Solution : Given $BA = C^2$

$$\Rightarrow \begin{pmatrix} 0 & -q \\ 1 & 0 \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 2 & -2 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} 2 & -2 \\ 2 & 2 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} 0 & -2q \\ p & 0 \end{pmatrix} = \begin{pmatrix} 0 & -8 \\ 8 & 0 \end{pmatrix}$$

$$\therefore p = 8, \quad -2q = -8, \quad q = 4$$

20. $A = \begin{pmatrix} 3 & 0 \\ 4 & 5 \end{pmatrix}$, $B = \begin{pmatrix} 6 & 3 \\ 8 & 5 \end{pmatrix}$, $C = \begin{pmatrix} 3 & 6 \\ 1 & 1 \end{pmatrix}$

find the matrix D , such that $CD - AB = 0$

Solution : Given $CD - AB = 0$

$$\Rightarrow \begin{pmatrix} 3 & 6 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 3 & 0 \\ 4 & 5 \end{pmatrix} \begin{pmatrix} 6 & 3 \\ 8 & 5 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} a+c & b+d \end{pmatrix} = \begin{pmatrix} 64 & 37 \end{pmatrix}$$

$$\therefore 3a + 6c = 18 \quad \dots\dots (1)$$

$$a + c = 64 \quad \dots\dots (2)$$

$$(1) \Rightarrow a + 2c = 6$$

$$(3) \Rightarrow a + c = 64$$

$$\underline{\hspace{1.5cm}} \quad c = -58$$

$$a - 58 = 64$$

$$a = 122$$

$$3b + 6d = 9 \quad \dots\dots (3)$$

$$b + d = 37 \quad \dots\dots (4)$$

$$(3) \Rightarrow b + 2d = 3$$

$$(4) \Rightarrow b + d = 37$$

$$\underline{\hspace{1.5cm}} \quad d = -34$$

$$b - 34 = 37$$

$$b = 71$$

$$\therefore a = 122, b = 71, c = -58, d = -34$$

$$\therefore D = \begin{pmatrix} 122 & 71 \\ -58 & -34 \end{pmatrix}$$

PROBLEMS FOR PRACTICE

- Solve : $2x + 5y + 2z = -38$,
 $3x - 2y + 4z = 17$, $-6x + y - 7z = 12$
(Ans : $x = 3, y = -8, z = -2$)
- Solve : $3x - 9z = 33$, $7x - 4y - z = -15$,
 $4x + 6y + 5z = -6$
(Ans : $x = -1, y = 3, z = -4$)
- Solve : $x + 2y + z = 7$, $x + 3z = 11$, $2x - 3y = 1$
(Ans : $(2, 1, 3)$)
- Solve : $x - \frac{y}{5} = 6$, $y - \frac{z}{7} = 8$, $z - \frac{x}{2} = 10$
(Ans : $(8, 10, 14)$)
- Solve : $\frac{1}{3}(x + y - 5) = y - z = 2x - 11 = 9 - (x + 2z)$
- Oviya, Sankee, Mithu have a total of \$ 89 in their wallets. Oviya has \$6. less than Mithu. Sankee has 3 times Mithu has How many does each have ?
(Ans : $13, 57, 19$)
- Sum of 3 numbers is 10. Sum of the first number, twice the second number and 3 times the third is 29 and the sum of first, four times the second and nine times the third is 43, Find the numbers.
(Ans : $4, 3, 3$)
- In a shop, the following items were sold on 3 days.

	Rice	Oil	Sugar
Day 1	25	10	10
Day 2	16	6	4
Day 3	30	12	6

If the total values sold were Rs.820, Rs.480 and Rs.912 respectively. Find the cost of 1 kg of each item.

(Ans : 12, 40, 12)

9. Find the GCD of the following :

i) $x^4 - 27a^3x$, $(x - 3a)^2$ **(Ans : $x - 3a$)**

ii) $x^3 + 8x^2 - x - 8$, $x^3 + x^2 - x - 1$
(Ans : $x^2 - 1$)

iii) $x^2 - x - 2$, $x^2 + x - 6$, $3x^2 - 13x + 14$
(Ans : $x - 2$)

iv) $3(x^2 - 5x + 6)$, $4(x^2 - 4x + 3)$
(Ans : $x - 3$)

v) $6a^2 - 11a + 3$, $12a^2 + 5a - 3$
(Ans : $3a - 1$)

10. Find the LCM of the following :

i) $q^2 - 4$, $q^3 - 8$, $q^2 - 6q + 8$
(Ans : $(q^2 - 4)(q - 4)(q^2 + 2q + 4)$)

ii) $2(x - 1)^2$, $3(x^2 - 1)$
(Ans : $6(x - 1)^2(x + 1)$)

iii) $2m^2 - 18n^2$, $5m^2n + 15mn^2$, $m^3 + 27n^3$
(Ans : $10mn(m - 3n)(m^2 + 27n^3)$)

iv) $6b^2 - b - 1$, $3b^2 + 7b + 2$, $2b^2 + 3b - 2$
(Ans : $(3b + 1)(2b - 1)(b + 2)$)

11. Find the GCD by long division :

i) $3x^4 + 6x^3 - 12x^2 - 24x$, $4x^4 + 14x^3 + 8x^2 - 8x$
(Ans : $x(x^2 + 4x + 4)$)

ii) $x^4 + x^3 + 4x^2 + 4x$, $x^3 - 3x^2 + 4x - 12$
(Ans : $x^2 + 4$)

iii) $3x^2 + 13x + 10$, $3x^3 + 18x^2 + 33x + 18$
(Ans : $x + 1$)

12. If $(x + 3)(x - 2)$ is the GCD of $f(x) = (x + 3)(2x^2 - 3x + a)$ and $g(x) = (x - 2)(3x^2 + 7x - b)$, find a and b .

(Ans : $a = -2$, $b = 6$)

13. The HCF and LCM of 2 polynomials are $5x^2 + x$ and $(x^3 - 4x)(5x + 1)$ respectively. One of the polynomials is $5x^3 - 9x^2 - 2x$. Find the other.

(Ans : $x(x + 2)(5x + 1)$)

14. Find the LCM of the polynomials $2x^3 + 15x^2 + 2x - 35$, $x^3 + 8x^2 + 4x - 21$ whose GCD is $x + 7$.

(Ans : $x^3 + 8x^2 + 4x - 21)(2x^2 + x - 5)$)

15. Simplify :

i) $\frac{2x^2 + 3x + 1}{x^2 + x + 1} \times \frac{4x^2 + 5x + 1}{x^2 + x + 1} \times \frac{15x^2 + 8x + 1}{x^2 + x + 1}$
(Ans : 1)

ii) $\frac{x^4 + x^2 + 1}{x^2 + x + 1}$
(Ans : $x^2 - x + 1$)

iii) $\frac{a^2 - (b + c)^2}{c^2 - (a + b)^2} \div \frac{b^2 - (c + a)^2}{a^2 + ab + ca}$
(Ans : $\frac{(a - b - c)a}{(c - a - b)(b - a - c)}$)

16. Solve the following by factorisation :

i) $2x^2 + 6\sqrt{3}x - 60 = 0$
(Ans : $2\sqrt{3}$, $-5\sqrt{3}$)

ii) $\frac{4}{x} - 3 = \frac{5}{2x + 3}$ **(Ans : 1, -2)**

iii) $9x^2 - 9(a + b)x + (2a^2 + 5ab + 2b^2) = 0$
(Ans : $\frac{a + 2b}{3}$, $\frac{2a + b}{3}$)

$$\text{iv)} (5x - 2)(x + 1) = 3x(3x - 1)$$

$$\left(\text{Ans : } 1, \frac{1}{2} \right)$$

$$\text{v)} \frac{2x+3}{x+8} = \frac{3(x-4)}{x-2} \quad (\text{Ans : } 5, -18)$$

17. Solve the following by completing the square.

$$\text{i)} 5x^2 - 6x - 2 = 0 \quad \left(\text{Ans : } \frac{3 \pm \sqrt{19}}{5} \right)$$

$$\text{ii)} 2x^2 + x - 4 = 0 \quad \left(\text{Ans : } \frac{-1 \pm \sqrt{33}}{4} \right)$$

$$\text{iii)} 4x^2 + 17x = 15 \quad \left(\text{Ans : } \frac{3}{4}, -5 \right)$$

$$\text{iv)} 2 \left(\frac{2x+1}{x+3} \right) - 3 \left(\frac{x+3}{2x-1} \right) = 5$$

$$\left(\text{Ans : } -10, \frac{-1}{5} \right)$$

18. Solve the following by using quadratic formula.

$$\text{i)} \frac{1}{x+1} + \frac{2}{x+2} = \frac{4}{x+4} \quad (\text{Ans : } 2 \pm 2\sqrt{3})$$

$$\text{ii)} 9x^2 - 6ax + (a^2 - b^2) = 0 \quad \left(\text{Ans : } \frac{a+b}{3}, \frac{a-b}{3} \right)$$

$$\text{iii)} a(x^2 + 1) = x(a^2 + 1) \quad \left(\text{Ans : } a, \frac{1}{a} \right)$$

$$\text{iv)} (x-2)(2x+3) = 3(x-4)(x+8)$$

$$(\text{Ans : } -6, -4)$$

19. Divide 18 into two parts such that twice the sum of their squares is 5 times their product.

$$(\text{Ans : } 6, 12)$$

20. In a music hall, the number of seats in each row is 10 less than the number of rows. If there are 704 seats in the hall. Find the number of rows.

$$(\text{Ans : } 32)$$

21. The perimeter of a right triangle is 60 cm. Its hypotenuse is 25 cm. Find the area of the triangle.

$$(\text{Ans : } 150 \text{ cm}^2)$$

22. The speed of a boat in still water is 15 Km/hr. It goes 30 km upstream and return down stream to the original point in $4\frac{1}{2}$ hrs. Find the speed of the stream.

$$(\text{Ans : } 5 \text{ Km/hr})$$

23. An aeroplane left 30 min later than its scheduled time and in order to reach its

time, it has to increase its speed by 250 Km/hr from its usual speed. Find the original speed.

$$(\text{Ans : } 750 \text{ Km/hr})$$

24. The numerator of a fraction is 3 less than its denominator. If 2 is added to both the numerator and the denominator, the the sum of new fraction and original fraction is $\frac{29}{20}$. Find the original fraction.

$$\left(\text{Ans : } \frac{7}{10} \right)$$

25. A flock of swans contained x^2 members. As the clouds gathered. $10x$ went lake, and $\frac{1}{8}x^2$ flew away to a garden. The remaining three couples played about in the water. Howmany swans were there in that lake ?

$$(\text{Ans : } 144)$$

26. If one root of $x^2 - 3x + \phi = 0$ is twice the other. Find the value of ϕ .
(Ans : 2)
27. Find 'k' if the following equations have real & equal roots.
i) $2x^2 - 10x + k = 0$ (Ans : 25/2)
ii) $x^2 - 2x(1 + 3k) + 7(3 + 2k) = 0$
(Ans : 2 (or) -10/9)
iii) $(k + 4)x^2 + (k + 1)x + 1 = 0$
(Ans : 5, -3)
iv) $(p + 1)x^2 - 6(p + 1)x + 3(p + q) = 0$
(Ans : -1, 3)
28. Show that the roots of the equation $3p^2x^2 - 2pqx + q^2 = 0$ are not real.
29. If α and β are the roots of $3x^2 - 6x + 1 = 0$,
i) $\alpha^2\beta, \beta^2\alpha$ (Ans : $27x^2 - 18x + 1 = 0$)
ii) $2\alpha + \beta, 2\beta + \alpha$
(Ans : $3x^2 - 18x + 25 = 0$)
30. Draw the graph of $y = x^2 + 2x - 3 = 0$ and hence solve $x^2 - x - 6 = 0$ (Ans : 3, -2)
31. Draw the graph of $y = x^2 + 3x + 2$ and hence solve $x^2 + 2x + 4 = 0$ (Ans : not real)
32. Draw the graph of $y = (2x + 3)(x - 2)$
33. Draw the graph of $y = 2x^2$ and hence solve $2x^2 + x - 6 = 0$. (Ans : -2, 3/2)
34. If $A = \begin{pmatrix} -2 \\ 4 \\ 5 \end{pmatrix}$, $B = (1 \quad 3 \quad -6)$, verify that $(AB)^T = B^T A^T$
35. Construct a 4×3 matrix whose elements are $a_{ij} = \frac{|2i - 3j|}{2}$
36. If $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 4 \\ -1 & 1 \end{pmatrix}$ does $(A + B)^2 = A^2 + 2AB + B^2$ hold ?
(NO)
37. If $A = \begin{pmatrix} 2 & 3 & -1 \\ 3 & 0 & 2 \end{pmatrix}$, $B = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ Verify that $A(BC) = (AB)C$.
38. If $A = \begin{pmatrix} 4 & 6 & 2 \\ 0 & 1 & 5 \\ 0 & 3 & 2 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 1 & -1 \\ 3 & -1 & 4 \\ -1 & 2 & 1 \end{pmatrix}$,
Verify $(A - B)^T = A^T - B^T$.
39. If $B \cdot B^T = 9I$, where $B = \begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ x & 2 & y \end{pmatrix}$, find x, y.
(Ans : -2, -1)
40. If $A = \begin{pmatrix} 3 & 2 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & 5 \end{pmatrix}$, show that $A^2 - 7A + 10I = 0$

OBJECTIVE TYPE QUESTIONS

1. The value of a, b, c, d

$$\begin{pmatrix} d+1 & 10+a \\ 3b-2 & a-4 \end{pmatrix} = \begin{pmatrix} 2 & 2a+1 \\ b-5 & 4c \end{pmatrix} \text{ if}$$

$$a) 11, 7, 3, 1 \quad b) 9, \frac{-3}{2}, 1, \frac{5}{4}$$

$$c) -9, \frac{3}{2}, \frac{-5}{4}, -1 \quad d) 9, \frac{-3}{2}, \frac{5}{4}, 1$$

(Ans : (d))

2. If
- $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{pmatrix}$
- , then
- A^2
- is

$$a) \text{ a null matrix} \quad b) \text{ a unit matrix}$$

$$c) -A \quad d) A$$

(Ans : (b))

3. If
- $A = \begin{pmatrix} 1 & 2 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
- ,
- $B = \begin{pmatrix} 1 & -2 & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
-
- and
- $AB = I_3$
- , then
- $x + y$
- equals

$$a) 0 \quad b) -1 \quad c) 2 \quad d) \text{ none}$$

(Ans : (a))

4. If
- $A = \begin{pmatrix} 5 & x \\ y & 0 \end{pmatrix}$
- and
- $A = A^T$
- , then

$$a) x = 0, y = 5 \quad b) x + y = 5$$

$$c) x = y \quad d) \text{ none}$$

(Ans : (c))

5. Matrix
- $A = (a_{ij})_{m \times n}$
- is a square matrix if

$$a) m < n \quad b) m > n$$

$$c) m = 1 \quad d) m = n$$

(Ans : (d))

6. If
- $\frac{1}{\alpha}$
- is a root of
- $2x^2 - 5x + 7 = 0$
- , then the value of
- $7\alpha^2 - 5\alpha$
- is

$$a) 2 \quad b) -2 \quad c) 5 \quad d) -5$$

(Ans : (b))

7. If
- $x^2 + \frac{1}{x^2} = 23, x > 0$
- , then
- $x + \frac{1}{x}$
- is

$$a) 2 \quad b) 3 \quad c) 4 \quad d) 5$$

(Ans : (d))

8. Which one of the following is a root of the equation
- $2x^4 - 5x^3 - 3x^2 + 13x + 9 = 0$

$$a) 1 \quad b) -1 \quad c) 2 \quad d) 0$$

(Ans : (b))

9. For what value of k, will the system of equations
- $2x + 3y = k$
- and
- $(k - 1)x + (k + 2)y = 3k$
- has infinite solutions?

$$a) -7 \quad b) 5 \quad c) 7 \quad d) 0$$

(Ans : (c))

10. The discriminant
- Δ
- of the equation
- $3\sqrt{3}x^2 + 10x + \sqrt{3} = 0$

$$a) 64 \quad b) -64 \quad c) 81 \quad d) \text{ none}$$

(Ans : (a))

11. The LCM of
- $6x^2y, 9x^2yz, 12x^2y^2z$
- is

$$a) 36xy^2z^2 \quad b) 36x^2y^2z$$

$$c) 36xyz \quad d) 36xyz$$

(Ans : (b))

12. On dividing
- $\frac{x^2 - 25}{x + 3}$
- by
- $\frac{x + 5}{x^2 - 9}$
- equal to

$$a) (x - 5)(x - 3) \quad b) (x - 5)(x + 3)$$

$$c) (x + 5)(x - 3) \quad d) (x + 5)(x + 3)$$

(Ans : (a))

13. The solution set of the equation
- $(x - 3)^2 = 9$
- is

$$a) \{0, 3\} \quad b) \{3, 3\} \quad c) \{3, 6\} \quad d) \{0, 6\}$$

(Ans : (d))

14. The equation whose roots are
- $b + c$
- and
- $b - c$
- is

$$a) x^2 + 2bx + (b^2 - c^2) = 0$$

$$b) x^2 - 2bx + (b^2 - c^2) = 0$$

$$c) x^2 - 2bx - (b^2 - c^2) = 0$$

$$d) x^2 + 2bx - (b^2 - c^2) = 0$$

(Ans : (b))

15. The value of m if
- $10x^2 + mx - 10$
- leaves a remainder 2 when divided by
- $2x - 3$
- .

$$a) 7 \quad b) -7 \quad c) \frac{-80}{3} \quad d) \frac{80}{3}$$

(Ans : (b))


CHAPTER

4

GEOMETRY

I. CONGRUENCY AND SIMILARITY OF TRIANGLES

Points to Remember




$$\Delta ABC \cong \Delta PQR$$

$$\angle A = \angle P, \angle B = \angle Q, \angle C = \angle R.$$

$$AB = PQ, BC = QR, CA = RP$$

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{CA}{RP} = 1$$

Same shape and same size.



$$\Delta ABC \sim \Delta PQR$$


$$\angle A = \angle P, \angle B = \angle Q, \angle C = \angle R.$$

$$AB \neq PQ, BC \neq QR, CA \neq RP$$

$$\text{but } \frac{AB}{PQ} = \frac{BC}{QR} = \frac{CA}{RP} > 1 \text{ or } < 1$$

Same shape but not same size.

- ✓ **AA Criterion of similarity** If two angles of one triangle are respectively equal to two angles of another triangle, then the two triangles are similar, because the third angle in both triangles must be equal. Therefore, AA similarity criterion is same as the AAA similarity criterion.
- ✓ **SAS Criterion of similarity.** If one angle of a triangle is equal to one angle of another triangle and if the sides including them are proportional then the two triangles are similar.
- ✓ **SSS Criterion of similarity.** If three sides of a triangle are proportional to the three corresponding sides of another triangle, then the two triangles are similar.
- ✓ A perpendicular line drawn from the vertex of a right angled triangle divides two triangles similar to each other and also to original triangle.




$$\Delta ADB \sim \Delta BDC, \Delta ABC \sim \Delta ADB, \Delta ABC \sim \Delta BDC$$

- ✓ If two triangles are similar, then the ratio of the corresponding sides are equal to the ratio of their corresponding altitudes.

i.e. if $\Delta ABC \sim \Delta PQR$ then

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{CA}{RP} = \frac{AD}{PS} = \frac{BE}{QT} = \frac{CF}{RU}$$



- ✓ If two triangles are similar, then the ratio of the corresponding sides are equal to the ratio of the corresponding perimeters.

$\triangle ABC \sim \triangle DEF$ then


$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{CA}{FD} = \frac{AB + BC + CA}{DE + EF + FD}$$

- ✓ The ratio of the area of two similar triangles are equal to the ratio of the squares of their corresponding sides.

$$\frac{\text{area}(\triangle ABC)}{\text{area}(\triangle PQR)} = \frac{AB^2}{PQ^2} = \frac{BC^2}{QR^2} = \frac{AC^2}{PR^2}$$

- ✓ If two triangles have common vertex and their bases are on the same straight line, the ratio between their areas is equal to the ratio between the length of their bases.



Here, $\frac{\text{area}(\triangle ABD)}{\text{area}(\triangle BDC)} = \frac{AD}{DC}$



- ✓ Two triangles are said to be similar if their corresponding sides are proportional.
- ✓ The triangles are equiangular if the corresponding angles are equal.

Example 4.1

Show that $\triangle PST \sim \triangle PQR$



Solution :

i) In $\triangle PST$ and $\triangle PQR$,

$$\frac{PS}{PQ} = \frac{2}{2+1} = \frac{2}{3}, \frac{PT}{PR} = \frac{4}{4+2} = \frac{2}{3}$$

Thus, $\frac{PS}{PQ} = \frac{PT}{PR}$ and $\angle P$ is common


Therefore, by SAS similarity,
 $\triangle PST \sim \triangle PQR$

ii) In $\triangle PST$ and $\triangle PQR$,

$$\frac{PS}{PQ} = \frac{2}{2+3} = \frac{2}{5}, \frac{PT}{PR} = \frac{2}{2+3} = \frac{2}{5}$$

Thus, $\frac{PS}{PQ} = \frac{PT}{PR}$ and $\angle P$ is common


Therefore, by SAS similarity,
 $\triangle PST \sim \triangle PQR$

Example 4.2Is $\triangle ABC \sim \triangle PQR$?**Solution :**In $\triangle ABC$ and $\triangle PQR$,

$$\frac{PQ}{AB} = \frac{3}{6} = \frac{1}{2}, \quad \frac{QR}{BC} = \frac{4}{10} = \frac{2}{5}$$

$$\text{since } \frac{1}{2} \neq \frac{2}{5}, \quad \frac{PQ}{AB} \neq \frac{QR}{BC}$$

The corresponding sides are not proportional.

Therefore $\triangle ABC$ is not similar to $\triangle PQR$.Observe Fig.4.18 and find $\angle P$.**Solution :**In $\triangle BAC$ and $\triangle PRQ$,

$$\frac{AB}{RQ} = \frac{3}{6} = \frac{1}{2}$$

$$\frac{BC}{QP} = \frac{6}{12} = \frac{1}{2}; \quad \frac{CA}{PR} = \frac{3\sqrt{3}}{6\sqrt{3}} = \frac{1}{2}$$

$$\text{Therefore, } \frac{AB}{RQ} = \frac{BC}{QP} = \frac{CA}{PR}$$

By SSS similarity, we have $\triangle BAC \sim \triangle QRP$ $\angle P = \angle C$ (since the corresponding parts of similar triangle)

$$\angle P = \angle C = 180^\circ - (\angle A + \angle B) = 180^\circ - (90^\circ + 60^\circ)$$

$$\angle P = 180^\circ - 150^\circ = 30^\circ$$

Example 4.4

A boy of height 90cm is walking away from the base of a lamp post at a speed of 1.2m/sec. If the lamp post is 3.6m above the ground, find the length of his shadow cast after 4 seconds.

Solution :

Given, speed = 1.2 m/s,




Fig. 4.19

time = 4 seconds

distance = speed \times time

$$= 1.2 \times 4 = 4.8 \text{ m}$$

Let x be the length of the shadow after 4 seconds

$$\begin{aligned} \text{Since, } \triangle ABE \sim \triangle CDE, \quad \frac{BE}{DE} &= \frac{AB}{CD} \text{ gives } \frac{4.8 + x}{x} \\ &= \frac{3.6}{0.9} = 4 \text{ (since } 90\text{cm} = 0.9\text{m)} \end{aligned}$$


$$48 = x + 4x \text{ gives } 3x = 4.8 \text{ so, } x = 1.6 \text{ m}$$

The length of his shadow $DE = 1.6 \text{ m}$ **Example 4.5**

In Fig. $\angle A = \angle CED$ prove that $\triangle CAB \sim \triangle CED$. Also find the value of x .

Solution :In $\triangle CAB$ and $\triangle CED$, $\angle C$ is common,

$$\angle A = \angle CED$$



Therefore, $\triangle CAB \sim \triangle CED$ (By AA similarity)

Hence

$$\frac{CA}{CE} = \frac{AB}{DE} = \frac{CB}{CD}$$


$$\frac{AB}{DE} = \frac{CB}{CD} \text{ gives } \frac{9}{x} = \frac{10+2}{8} \text{ so, } x = \frac{8 \times 9}{12} = 6 \text{ cm.}$$

Example 4.6

In Fig. QA and PB are perpendiculars to AB. If $AO = 10$ cm, $BO = 6$ cm and $PB = 9$ cm. Find AQ.

Solution :

In $\triangle AOQ$ and $\triangle BOP$, $\angle OAQ = \angle OBP = 90^\circ$



$\angle AOQ = \angle BOP$ (Vertically opposite angles)

Therefore, by AA Criterion of similarity,

$\triangle AOQ \sim \triangle BOP$

$$\frac{AO}{BO} = \frac{OQ}{OP} = \frac{AQ}{BP}$$


$$\frac{10}{6} = \frac{AQ}{9} \text{ gives } AQ = \frac{10 \times 9}{6} = 15 \text{ cm}$$

Example 4.7

The perimeters of two similar triangles ABC and PQR are respectively 36 cm and 24 cm. If $PQ = 10$ cm, find AB.

Solution :

The ratio of the corresponding sides of similar triangles is same as the ratio of their perimeters,



Since $\triangle ABC \sim \triangle PQR$,

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR} = \frac{36}{24}$$

$$\frac{AB}{PQ} = \frac{36}{24} \text{ gives } \frac{AB}{10} = \frac{36}{24}$$

$$AB = \frac{36 \times 10}{24} = 15 \text{ cm}$$

Example 4.8

If $\triangle ABC$ is similar to $\triangle DEF$ such that $BC = 3$ cm, $EF = 4$ cm and area of $\triangle ABC = 54 \text{ cm}^2$. Find the area of $\triangle DEF$.

Solution :

Since the ratio of area of two similar triangles is equal to the ratio of the squares of any two corresponding sides, we have

$$\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DEF)} = \frac{BC^2}{EF^2} \text{ gives } \frac{54}{\text{Area}(\triangle DEF)} = \frac{3^2}{4^2}$$


$$\text{Area}(\triangle DEF) = \frac{16 \times 54}{9} = 96 \text{ cm}^2$$

Example 4.9

Two poles of height 'a' metres and 'b' metres are 'p' metres apart. Prove that the height of the point of intersection of the lines joining the top of each pole to the foot of the opposite pole is given by $\frac{ab}{a+b}$ metres.

Solution :

Let AB and CD be two poles of height 'a' metres and 'b' metres respectively such that the poles are 'p' metres apart. That is AC = p metres. Suppose the lines AD and BC meet at O, such that OL = h metres



Let CL = x and LA = y.

Then, $x + y = p$

In $\triangle ABC$ and $\triangle LOC$, we have $\angle CAB = \angle CLO$ [each equals to 90°]

$\angle C = \angle C$ [C is common]

$\triangle CAB \sim \triangle CLO$ [By AA similarity]

$$\frac{CA}{CL} = \frac{AB}{LO} \text{ gives } \frac{p}{x} = \frac{a}{h}$$

$$\text{so, } x = \frac{ph}{a} \quad \dots\dots\dots (1)$$

In $\triangle ALO$ and $\triangle ACD$, we have

$\angle ALO = \angle ACD$ [each equal to 90°]

$\angle A = \angle A$ [A is common]

$\triangle ALO \sim \triangle ACD$ [By AA similarity]

$$\frac{AL}{AC} = \frac{OL}{DC} \text{ gives}$$

$$\frac{y}{p} = \frac{h}{b} \text{ we get, } y = \frac{ph}{b} \quad \dots\dots\dots (2)$$

$$(1) + (2) \text{ gives } x + y = \frac{ph}{a} + \frac{ph}{b}$$

$$p = ph \left(\frac{1}{a} + \frac{1}{b} \right) \text{ (Since } x + y = p \text{)}$$

$$1 = h \left(\frac{a+b}{ab} \right)$$

$$\text{Therefore, } h = \frac{ab}{a+b}$$

Hence, the height of the intersection of the lines joining the top of each pole to the foot of the opposite pole is $\frac{ab}{a+b}$ metres.

Construction of similar triangles**Example 4.10**


Construct a triangle similar to a given triangle PQR with its sides equal to $\frac{3}{5}$ of the corresponding sides of the triangle PQR (scale factor $\frac{3}{5} < 1$)

Solution :

Given a triangle PQR we are required to construct another triangle whose sides are $\frac{3}{5}$ of the corresponding sides of the triangle PQR.

Steps of construction

1. Construct a $\triangle PQR$ with any measurement.



2. Draw a ray QX making an acute angle with QR on the side opposite to vertex P.
3. Locate 5 (the greater of 3 and 5 in $\frac{3}{5}$) points.

$Q_1, Q_2, Q_3, Q_4,$ and Q_5 on QX so that $QQ_1 = Q_1Q_2 = Q_2Q_3 = Q_3Q_4 = Q_4Q_5$

4. Join Q_5R and draw a line through Q_3 (the third point, 3 being smaller of 3 and 5 in $\frac{3}{5}$) parallel to Q_5R to intersect QR at R' .
5. Draw line through R' parallel to the line RP to intersect QP at P' . Then, $\Delta P'QR'$ is the required triangle each of whose sides is three-fifths of the corresponding sides of ΔPQR .

Example 4.11

Construct a triangle similar to a given triangle PQR with its sides equal to $\frac{7}{4}$ of the corresponding sides of the triangle PQR (scale factor $\frac{7}{4} > 1$)




Fig. 4.26

Given a triangle PQR , we are required to construct another triangle whose sides are $\frac{7}{4}$ of the corresponding sides of the triangle PQR .


Steps of construction

1. Construct a ΔPQR with any measurement.
2. Draw a ray QX making an acute angle with QR on the side opposite to vertex P .
3. Locate 7 (the greater of 4 and 7 in $\frac{7}{4}$) points. $Q_1, Q_2, Q_3, Q_4, Q_5, Q_6$ and Q_7 on QX so that $QQ_1 = Q_1Q_2 = Q_2Q_3 = Q_3Q_4 = Q_4Q_5 = Q_5Q_6 = Q_6Q_7$


4. Join Q_4 (the 4th point, 4 being smaller of 4 and 7 in $\frac{7}{4}$) to R and draw a line through Q_7 parallel to Q_4R , intersecting the extended line segment QR at R' .
5. Draw a line through R' parallel to RP intersecting the extended line segment QP at P' . Then $\Delta P'QR'$ is the required triangle each of whose sides is seven-fourths of the corresponding sides of ΔPQR .

EXERCISE 4.1

1. Check whether the which triangles are similar and find the value of x .



(i)



(ii)

Solution:

$$i) \frac{AE}{AC} = \frac{2}{2+3.5} = \frac{2}{5.5} = \frac{4}{11}$$

$$\frac{AD}{AB} = \frac{3}{3+5} = \frac{3}{8}$$

$$\therefore \frac{AE}{AC} \neq \frac{AD}{AB}$$

\therefore the 2 triangles are not similar.

- ii) Given $\angle PQB = 110^\circ \Rightarrow \angle PQC = 70^\circ = \angle QBA$

\therefore Corresponding angles are equal.

$$\therefore \frac{CQ}{QB} = \frac{PQ}{AB}$$


$$\Rightarrow \frac{3}{3} = \frac{x}{5}$$

$$\Rightarrow 1 = \frac{x}{5}$$

$$\therefore x = 5$$

2. A girl looks the reflection of the top of the lamp post on the mirror which is 66 m away from the foot of the lamppost. The girl whose height is 12.5 m is standing 2.5 m away from the mirror. Assuming the mirror is placed on the ground facing the sky and the girl, mirror and the lamp-post are in a same line, find the height of the lamp post.

Solution :



Given $AB = \text{height of girl} = 1.5 \text{ m}$
 $BC = \text{Dist. between girl \& Mirror} = 0.4 \text{ m}$
 $LP = \text{height of lamp post} = h$
 $CP = \text{dist. between Mirror \& Post} = 87.6 \text{ m}$

In $\triangle ABC$, $\triangle LPC$,
 $\angle B = \angle P = 90^\circ$
 $\angle ACB = \angle LCP$ (angle of incidence & angle of reflection)
 $\therefore \triangle ABC$ & $\triangle LPC$ are similar.

$$\therefore \frac{AB}{LP} = \frac{BC}{CP} \quad (\text{By AA similarity})$$

$$\Rightarrow \frac{1.5}{h} = \frac{0.4}{87.6}$$

$$\Rightarrow h = \frac{87.6 \times 1.5}{0.4}$$

$$= \frac{87.6 \times \frac{3}{2}}{\frac{4}{10}}$$

$$= 43.8 \times 3 \times \frac{5}{2}$$


$$= 21.9 \times 15$$

$$= 328.5 \text{ m}$$

\therefore Height of the lamp post = 328.5 m

3. A vertical stick of length 6 m casts a shadow 400 cm long on the ground and at the same time a tower casts a shadow 28 m long. Using similarity, find the height of the tower.

Solution :



In $\triangle ABC$ and $\triangle PQR$

$$\angle B = \angle Q = 90^\circ$$

$$\angle C = \angle R \quad (AC \parallel PR)$$

\therefore By AA similarity, $\triangle ABC \sim \triangle PQR$

$$\frac{AB}{PQ} = \frac{BC}{QR}$$

$$\Rightarrow \frac{6}{h} = \frac{4}{28}$$


$$\Rightarrow \frac{6}{h} = \frac{1}{7}$$

$$\Rightarrow h = 42 \text{ m}$$

\therefore Height of the tower = 42m.

4. Two triangles QPR and QSR, right angled at P and S respectively are drawn on the same base QR and on the same side of QR. If PR and SQ intersect at T, prove that $PT \times TR = ST \times TQ$.

Solution :



Consider ΔPQT and ΔSRT

i) $\angle P = \angle S = 90^\circ$

ii) $\angle PTQ = \angle STR$ (Vertically Opp. angle)

\therefore By AA similarity,


$$\Delta PQT \sim \Delta SRT$$

$$\therefore \frac{QT}{TR} = \frac{PT}{ST}$$

$$\Rightarrow PT \times TR = ST \times TQ$$

Hence proved.

5. In the adjacent figure, ΔABC is right angled at C and $DE \perp AB$. Prove that $\Delta ABC \sim \Delta ADE$ and hence find the lengths of AE and DE.



Solution :

In ΔABC & ΔADE ,

i) $\angle AED = \angle ACB = 90^\circ$

ii) $\angle A$ is common

\therefore By AA similarity,

$$\Delta ABC \sim \Delta ADE$$

Also,

$$AB^2 = AC^2 + BC^2$$

$$= 5^2 + 12^2$$

$$= 25 + 144$$

$$= 169$$

$$\therefore AB = 13$$


\therefore By similarity,

$$\frac{AD}{AB} = \frac{ED}{BC} = \frac{AE}{AC}$$

$$\Rightarrow \frac{3}{13} = \frac{DE}{12} = \frac{AE}{5}$$

$$\therefore AE = \frac{15}{13}, DE = \frac{36}{13}$$

6. In the adjacent figure, $\Delta ACB \sim \Delta APQ$. If $BC = 8$ cm, $PQ = 4$ cm, $BA = 6.5$ cm and $AP = 2.8$ cm, find CA and AQ.



Solution :


Given $\Delta ACB \sim \Delta APQ$

$$\therefore \frac{BC}{PQ} = \frac{AC}{AP} = \frac{AB}{AQ}$$

$$\Rightarrow \frac{8}{4} = \frac{AC}{2.8} = \frac{6.5}{AQ}$$

$$\frac{AC}{2.8} \quad \left| \quad \frac{6.5}{AQ} = 2 \right. \\ \Rightarrow AC = 5.6 \text{ cm} \quad \left| \quad \Rightarrow AQ = \frac{6.5}{2} = 3.25 \text{ cm} \right.$$

7. If figure OPRQ is a square and $\angle MLN = 90^\circ$. Prove that i) $\Delta LOP \sim \Delta QMO$
ii) $\Delta LOP \sim \Delta RPN$ iii) $\Delta QMO \sim \Delta RPN$
iv) $QR^2 = MQ \times RN$



Solution :

i) In ΔLOP , ΔQMO

a) $\angle OLP = \angle OQM = 90^\circ$

b) $\angle LOP = \angle OMQ$ (Corresponding angles)

\therefore By AA similarity,

$$\Delta LOP \sim \Delta QMO$$

ii) In $\triangle LOP$, $\triangle RPN$

a) $\angle OLP = \angle PRN = 90^\circ$

b) $\angle LPO = \angle PNR$ (Corresponding angles)

\therefore By AA similarity,

$$\triangle LOP \sim \triangle RPN$$


\therefore From (i) & (ii)

$$\triangle QMO \sim \triangle RPN$$

$$\begin{aligned} \therefore \frac{QM}{RP} &= \frac{QO}{RN} \Rightarrow \frac{QM}{QR} = \frac{QR}{RN} \quad (\because \text{Square}) \\ &\Rightarrow QR^2 = MQ \times RN \end{aligned}$$

8. If $\triangle ABC \sim \triangle DEF$ such that area of $\triangle ABC$ is 9cm^2 and the area of $\triangle DEF$ is 16cm^2 and $BC = 2.1\text{ cm}$. Find the length of EF .


Solution :



Given $\triangle ABC \sim \triangle DEF$

$$\begin{aligned} \therefore \frac{\text{Area of } \triangle ABC}{\text{Area of } \triangle DEF} &= \frac{BC^2}{EF^2} \\ \Rightarrow \frac{9}{16} &= \frac{(2.1)^2}{EF^2} \\ \Rightarrow EF^2 &= \frac{16 \times (2.1)^2}{9} \\ \therefore EF &= \frac{4 \times 2.1}{3} = 2.8\text{ cm} \end{aligned}$$

9. Two vertical poles of heights 6 m and 3 m are erected above a horizontal ground AC. Find the value of y .



Solution :

From the fig.

$$\triangle PAC \sim \triangle QBC$$

$$\angle PAC = \angle QBC \text{ (Corresponding angles)}$$

$$\angle C = \angle C$$

$$\therefore \frac{CB}{CA} = \frac{QB}{PA}$$

$$\Rightarrow \frac{CB}{CA} = \frac{y}{6} \quad \dots\dots\dots (1)$$

$$\angle RCA = \angle QBA \text{ (Corresponding angles)}$$

$$\angle A = \angle A$$

$$\therefore \frac{AB}{AC} = \frac{BQ}{RC} \Rightarrow \frac{AB}{BC} = \frac{y}{3} \quad \dots\dots(2)$$

Adding (1) & (2),

$$\frac{AB+BC}{AC} = \frac{y}{6} + \frac{y}{3}$$

$$\Rightarrow \frac{AC}{AC} = y \left(\frac{1}{6} + \frac{1}{3} \right)$$

$$\Rightarrow y \left(\frac{1+2}{6} \right) = 1$$

$$\Rightarrow y \left(\frac{1}{2} \right) = 1$$

$$\therefore y = 2\text{ m}$$

(or) Using formula $y = \frac{ab}{a+b}$


$$= \frac{6 \times 3}{6+3} = \frac{18}{9} = 2\text{ m}$$

10. Construct a triangle similar to a given triangle PQR with its sides equal to $\frac{2}{3}$

of the corresponding sides of the triangle

PQR (scale factor $\frac{2}{3}$).

Solution :




1. Construct a ΔPQR with any measurement.
2. Draw a ray QX making an acute angle with QR on the side opposite to vertex P .
3. Locate 3 points (greater of 2 and 3 in $\frac{2}{3}$) points.
 Q_1, Q_2, Q_3 on QX so that
 $QQ_1 = Q_1Q_2 = Q_2Q_3$
4. Join Q_3R and draw a line through Q_2 (3 being smaller of 2 and 3 in $\frac{2}{3}$) parallel to Q_3R to intersect QR at R' .
5. Draw line through R' parallel to the line RP intersecting the QP at P' . Then, $\Delta P'QR'$ is the required Δ .

11. Construct a triangle similar to a given triangle LMN with its sides equal to $\frac{4}{5}$

of the corresponding sides of the triangle

LMN (scale factor $\frac{4}{5}$).

Solution :




Steps of construction

1. Construct a ΔLMN with any measurement.
 2. Draw a ray MX making an acute angle with MN on the side opposite to vertex L .
 3. Locate 5 points (greater of 4 and 5 in $\frac{4}{5}$) points.
 M_1, M_2, M_3, M_4 & M_5 so that $MM_1 = M_1M_2 = M_2M_3 = M_3M_4 = M_4M_5$,
 4. Join M_5 to N and draw a line through M_4 (4 being smaller of 4 and 5 in $\frac{4}{5}$) parallel to M_5N to intersect MN at N' .
 5. Draw line through N' parallel to the line LN intersecting line segment ML to L' .
- Then, $L'M'N'$ is the required Δ .

12. Construct a triangle similar to a given triangle ABC with its sides equal to $\frac{6}{5}$

of the corresponding sides of the triangle ABC (scale factor $\frac{6}{5}$).

Solution :**Steps of construction**

1. Construct a $\triangle ABC$ with any measurement.
2. Draw a ray BX making an acute angle with BC on the side opposite to vertex A .

 $\frac{6}{5}$


points.

B_1, B_2, \dots, B_6 on BX so that $BB_1 = B_1B_2 = B_2B_3 = B_3B_4 = B_4B_5 = B_5B_6$,

4. Join B_4 (4 being smaller of 4 and 6 in $\frac{6}{4}$) to C and draw a line through B_6 parallel to B_4C to intersecting the extended line segment BC at C' .
5. Draw line through C' parallel to CA intersect the extended line segment BA to A' .

Then, $\triangle A'B'C'$ is the required \triangle .

-
- 13. Construct a triangle similar to a given triangle PQR with its sides equal to $\frac{7}{3}$ of the corresponding sides of the triangle PQR (scale factor $\frac{7}{3}$).**

Solution :**Steps of construction**

1. Construct a $\triangle PQR$ with any measurement.
2. Draw a ray PX making an acute angle with PR on the side opposite to vertex Q .
3. Locate 7 points (greater of 3 and 7 in $\frac{7}{3}$) points.

P_1, P_2, \dots, P_7 on PX so that $PP_1 = P_1P_2 = P_2P_3 = \dots = P_6P_7$,

4. Join P_3R (3 being smaller of 3 and 7 in $\frac{7}{3}$) and draw a line through P_7 parallel to P_3R to intersecting the extended line segment PR at R' .
 5. Draw line through R' parallel to QR intersect the extended line segment PQ to Q' .
- Then, $\triangle P'Q'R'$ is the required \triangle .
-

II. Thales Theorem and Angle Bisector Theorem


Theorem 1: Basic Proportionality Theorem (BPT) or Thales theorem

Statement

A straight line drawn parallel to a side of triangle intersecting the other two sides, divides the sides in the same ratio.

Proof

Given: In $\triangle ABC$, D is a point on AB and E is a point on AC.



To Prove : $\frac{AD}{DB} = \frac{AE}{EC}$

Construction : Draw a line $DE \parallel BC$


No.	Statement	Reason
1.	$\angle ABC = \angle ADE$ $= \angle 1$	Corresponding angles are equal because $DE \parallel BC$
2.	$\angle ACB = \angle AED$ $= \angle 2$	Corresponding angles are equal because $DE \parallel BC$
3.	$\angle DAE = \angle BAC$ $= \angle 3$	Both triangles have a common angle
	$\triangle ABC \sim \triangle ADE$	By AAA similarity
	$\frac{AB}{AD} = \frac{AC}{AE}$	Corresponding sides are proportional
	$\frac{AD + DB}{AD} = \frac{AE + EC}{AE}$	Split AB and AC using the points D and E.

4.	$1 + \frac{DB}{AD} = 1 + \frac{EC}{AE}$ $\frac{DB}{AD} = \frac{EC}{AE}$ $\frac{AD}{DB} = \frac{AE}{EC}$	On simplification Cancelling 1 on both sides Taking reciprocals
Hence proved		

Theorem 2: Converse of Basic Proportionality Theorem

Statement

If a straight line divides any two sides of a triangle in the same ratio, then the line must be parallel to the third side.



Given : In $\triangle ABC$, $\frac{AD}{DB} = \frac{AE}{EC}$

To Prove : $DE \parallel BC$

Construction : Draw $BF \parallel DE$

No.	Statement	Reason
1.	In $\triangle ABC$, $BF \parallel DE$	Construction
2.	$\frac{AD}{DB} = \frac{AE}{EC} \dots (1)$	Thales theorem (In $\triangle ABC$ taking D in AB and E in AC).
3.	$\frac{AD}{DB} = \frac{AF}{FC} \dots (2)$	Thales theorem (In $\triangle ABC$ taking F in AC)


4.	$\frac{AE}{EC} = \frac{AF}{FC}$ $\frac{AE}{EC} + 1 = \frac{AF}{FC} + 1$ $\frac{AE + EC}{EC} = \frac{AF + FC}{FC}$ $\frac{AC}{EC} = \frac{AC}{FC}$ $EC = FC$ <p>Therefore, F = C Thus DE BC</p>	<p>From (1) and (2)</p> <p>Adding 1 to both sides</p> <p>Cancelling AC on both sides</p> <p>F lies between E and C.</p> <p>Hence proved</p>
----	---	---

Theorem 3: Angle Bisector Theorem

Statement

angle divides the opposite side internally in the ratio of the corresponding sides containing the angle.

Proof



Given : In $\triangle ABC$, AD is the internal bisector

To Prove : $\frac{AB}{AC} = \frac{BD}{CD}$

Construction : Draw a line through C parallel to AB. Extend AD to meet line through C at E


No.	Statement	Reason
1.	$\angle AEC = \angle BAE = \angle 1$	Two parallel lines cut by a transversal make alternate angles equal.
2.	$\triangle ACE$ is isosceles $AC = CE \dots (1)$ $\triangle ABD \sim \triangle ECD$	In $\triangle ACE$, $\angle CAE = \angle CEA$
3.	$\frac{AB}{CE} = \frac{BD}{CD}$	By AA Similarity
4.	$\frac{AB}{AC} = \frac{BD}{CD}$	From (1) $AC = CE$. Hence proved.

Theorem 4: Converse of Angle Bisector Theorem

Statement

If a straight line through one vertex of a triangle divides the opposite side internally in the ratio of the other two sides, then the line bisects the angle internally at the vertex.

Proof



Given : ABC is a triangle. AD divides BC in the ratio of the sides containing the angles $\angle A$ to meet BC at D.

That is $\frac{AB}{AC} = \frac{BD}{DC} \dots\dots\dots (1)$

To prove : AD bisects $\angle A$ i.e. $\angle 1 = \angle 2$

Construction : Draw $CE \parallel DA$. Extend BA to meet at E.


No.	Statement	Reason
1.	$\angle BAD = \angle 1$ and $\angle DAC = \angle 2$	Assumption
2.	$\angle BAD = \angle AEC = \angle 1$	Since $DA \parallel CE$ and AC is transversal, corresponding angles are equal
3.	$\angle DAC = \angle ACE = \angle 2$	Since $DA \parallel CE$ and AC is transversal. Alternate angles are equal.
4.	$\frac{BA}{AE} = \frac{BD}{DC} \dots(2)$	In $\triangle BCE$ by Thales theorem
5.	$\frac{AB}{AC} = \frac{BD}{DC}$	From (1)
6.	$\frac{AB}{AC} = \frac{AE}{AC}$	From (1) and (2)
7.	$AC = AE \dots(3)$	Cancelling AB
8.	$\angle 1 = \angle 2$	$\triangle ACE$ is isosceles by (3)
9.	AD bisects $\angle A$	Since, $\angle 1 = \angle BAD = \angle 2 = \angle DAC$. Hence proved.

Example 4.12

In $\triangle ABC$ if $DE \parallel BC$, $AD = x$, $DB = x - 2$, and $EC = x - 1$ then find the lengths of the sides AB and AC .

Solution :

In $\triangle ABC$ we have $DE \parallel BC$



By Thales theorem, we have $\frac{AD}{DB} = \frac{AE}{EC}$

$$\frac{x}{x-2} = \frac{x+2}{x-1} \text{ gives } x(x-1) = (x-2)(x+2)$$

Hence, $x^2 - x = x^2 - 4$ so, $x = 4$

When $x = 4$, $AD = 4$, $DB = x - 2 = 2$,

$AE = x + 2 = 6$, $EC = x - 1 = 3$.

Hence, $AB = AD + DB = 4 + 2 = 6$,

$AC = AE + EC = 6 + 3 = 9$.

Therefore, $AB = 6$, $AC = 9$.


Example 4.13

D and E are respectively the points on the sides AB and AC of a $\triangle ABC$ such that $AB = 5.6$ cm, $AD = 1.4$ cm, $AC = 7.2$ cm and $AE = 1.8$ cm, show that $DE \parallel BC$.

Solution :

We have $AB = 5.6$ cm, $AD = 1.4$ cm, $AC = 7.2$ cm and $AE = 1.8$ cm

$$BD = AB - AD = 5.6 - 1.4 = 4.2 \text{ cm}$$



$$\text{and } EC = AC - AE = 7.2 - 1.8 = 5.4 \text{ cm.}$$

$$\frac{AD}{DB} = \frac{1.4}{4.2} = \frac{1}{3} \text{ and } \frac{AE}{EC} = \frac{1.8}{5.4} = \frac{1}{3}$$

$$\frac{AD}{DB} = \frac{AE}{EC}$$

Therefore, by converse of Basic Proportionality Theorem, we have DE is parallel to BC .

Hence proved.


Example 4.14

In the Fig. $DE \parallel AC$ and $DC \parallel AP$. Prove that

$$\frac{BE}{EC} = \frac{BC}{CP}.$$

Solution :

In ΔBPA , we have $DC \parallel AP$. By Basic Proportionality Theorem,



$$\text{We have } \frac{BC}{CP} = \frac{BD}{DA} \quad \dots\dots\dots (1)$$

In ΔBCA , we have $DE \parallel AC$. By Basic Proportionality Theorem,

$$\text{We have } \frac{BE}{EC} = \frac{BD}{DA} \quad \dots\dots\dots (2)$$


From (1) and (2) we get, $\frac{BE}{EC} = \frac{BC}{CP}$.
Hence proved.

Example 4.15

In the Fig., AD is the bisector of $\angle A$. If $BD = 4$ cm, $DC = 3$ cm and $AB = 6$ cm, find AC.

Solution :

In ΔABC , AD is the bisector of $\angle A$



Therefore by Angle Bisector Theorem

$$\frac{BD}{DC} = \frac{AB}{AC}$$

$$\frac{4}{3} = \frac{6}{AC} \text{ gives } 4AC = 18.$$


$$\text{Hence } AC = \frac{9}{2} = 4.5 \text{ cm}$$

Example 4.16

In the Fig. AD is the bisector of $\angle BAC$. If $AB = 10$ cm, $AC = 14$ cm and $BC = 6$ cm, find BD and DC.

Solution :

Let $BD = x$ cm, then $DC = (6 - x)$ cm



AD is the bisector of $\angle A$

Therefore by Angle Bisector Theorem

$$\frac{AB}{AC} = \frac{BD}{DC}$$

$$\frac{10}{14} = \frac{x}{6-x} \text{ gives } \frac{5}{7} = \frac{x}{6-x}$$


$$\text{So, } 12x = 30 \text{ we get, } x = \frac{30}{12} = 2.5 \text{ cm}$$


$$\text{Therefore, } BD = 2.5 \text{ cm, } DC = 6 - x$$

$$= 6 - 2.5 = 3.5 \text{ cm}$$

Construction of triangle**Example 4.17**

Construct a ΔPQR in which $PQ = 8$ cm, $\angle R = 60^\circ$ and the median RG from R to PQ is 5.8 cm. Find the length of the altitude from R to PQ .

Solution :



Construction

Step 1 : Draw a line segment $PQ = 8\text{cm}$.

Step 2 : At P, draw PE such that $\angle QPE = 60^\circ$.

Step 3 : At P, draw PF such that $\angle EPF = 90^\circ$.

Step 4 : Draw the perpendicular bisector to PQ, which intersects PF at O and PQ at G.

Step 5 : With O as centre and OP as radius draw a circle.

Step 6 : From G mark arcs of radius 5.8 cm on the circle. Mark them as R and S.

Step 7 : Join PR and RQ. Then $\triangle PQR$ is the required triangle.



Step 8 : From R draw a line RN perpendicular to LQ. LQ meets RN at M

Step 9 : The length of the altitude is $RM = 3.5\text{ cm}$.

Example 4.18

Construct a triangle $\triangle PQR$ such that $QR = 5\text{ cm}$, $\angle P = 30^\circ$ and the altitude from P to QR is of length 4.2 cm.

Solution :



Construction

Step 1 : Draw a line segment $QR = 5\text{cm}$.

Step 2 : At Q, draw QE such that $\angle RQE = 30^\circ$.

Step 3 : At Q, draw QF such that $\angle EQF = 90^\circ$.

Step 4 : Draw the perpendicular bisector XY to QR, which intersects QF at O and QR at G.

Step 5 : With O as centre and OQ as radius draw a circle.

Step 6 : From G mark an arc in the line XY at M, such that $GM = 4.2\text{ cm}$.

Step 7 : Draw AB through M which is parallel to QR.



Step 8 : AB meets the circle at P and S.

Step 9 : Join QP and RP. Then $\triangle PQR$ is the required triangle.

Example 4.19

Draw a triangle ABC of base BC = 8 cm, $\angle A = 60^\circ$ and the bisector of $\angle A$ meets BC at D such that BD = 6 cm.

Solution :

**Construction**

Step 1 : Draw a line segment BC = 8cm.

Step 2 : At B, draw BE such that $\angle CBE = 60^\circ$.

Step 3 : At B, draw BF such that $\angle EBF = 90^\circ$.

Step 4 : Draw the perpendicular bisector to BC, which intersects BF at O and BC at G.

Step 5 : With O as centre and OB as radius draw a circle.

Step 6 : From B mark an arcs of 6 cm on BC at D.

Step 7 : The perpendicular bisector intersects the circle at I. Join ID.

Step 8 : ID produced meets the circle at A. Now join AB and AC. Then $\triangle ABC$ is the required triangle.


EXERCISE 4.2

1. In $\triangle ABC$, D and E are points on the sides AB and AC respectively such that $DE \parallel BC$

(i) If $\frac{AD}{DB} = \frac{3}{4}$ and AC = 15 cm find AE.

(ii) If $AD = 8x - 7$, $DB = 5x - 3$, $AE = 4x - 3$ and $EC = 3x - 1$, find the value of x.

Solution :



- i) Given $\frac{AD}{DB} = \frac{3}{4}$, AC = 15

$$\begin{aligned} \therefore \text{By BPT } \frac{AD}{DB} &= \frac{AE}{EC} \\ \Rightarrow \frac{3}{4} &= \frac{x}{15-x} \\ \Rightarrow 4x &= 45 - 3x \\ \Rightarrow 7x &= 45 \end{aligned}$$

$$x = \frac{45}{7} = 6.428 \approx 6.43 \text{ cm}$$


- ii) Given $AD = 8x - 7$, $DB = 5x - 3$

$$AE = 4x - 3, EC = 3x - 1$$

$$\begin{aligned} \text{By BPT } \frac{AD}{DB} &= \frac{AE}{EC} \\ \Rightarrow \frac{8x-7}{5x-3} &= \frac{4x-3}{3x-1} \\ \Rightarrow (8x-7)(3x-1) &= (4x-3)(5x-3) \\ \Rightarrow 24x^2 - 29x + 7 &= 20x^2 - 27x + 9 \\ \Rightarrow 4x^2 - 2x - 2 &= 0 \\ \Rightarrow 2x^2 - x - 1 &= 0 \\ \therefore x &= 1, -\frac{1}{2} \\ x &= 1 \end{aligned}$$

2. ABCD is a trapezium in which $AB \parallel DC$ and P, Q are points on AD and BC respectively, such that $PQ \parallel DC$ if $PD = 18$ cm, $BQ = 35$ cm and $QC = 15$ cm, find AD.

Solution :



In trapezium ABCD, $AB \parallel DC \parallel PQ$

Join AC, meet PQ at R.

In $\triangle ACD$, $PR \parallel DC$

$$\begin{aligned} \therefore \text{By BPT } \frac{AP}{PD} &= \frac{AR}{RC} \\ \Rightarrow \frac{x}{18} &= \frac{AR}{RC} \quad \dots\dots\dots (1) \end{aligned}$$

In $\triangle ABC$, $RQ \parallel AB$

$$\begin{aligned} \therefore \text{By ABT } \frac{BQ}{QC} &= \frac{AR}{RC} \\ \Rightarrow \frac{35}{15} &= \frac{AR}{RC} \\ \Rightarrow \frac{7}{3} &= \frac{AR}{RC} \quad \dots\dots\dots (2) \end{aligned}$$

\therefore From (1) & (2)


$$\begin{aligned} \frac{x}{18} &= \frac{7}{3} \\ \Rightarrow x &= 42 \\ \therefore AD &= AP + PD \\ &= 42 + 18 \\ &= 60 \text{ m} \end{aligned}$$

3. In $\triangle ABC$, D and E are points on the sides AB and AC respectively. For each of the following cases show that $DE \parallel BC$

(i) $AB = 12$ cm, $AD = 8$ cm, $AE = 12$ cm and $AC = 18$ cm.

- (ii) $AB = 5.6$ cm, $AD = 1.4$ cm, $AC = 7.2$ cm and $AE = 1.8$ cm.

Solution :



In $\triangle ABC$, To Prove : $DE \parallel BC$

$$\begin{aligned} \text{i) } \frac{AD}{AB} &= \frac{8}{12} = \frac{2}{3} \\ \frac{AE}{AC} &= \frac{12}{18} = \frac{2}{3} \\ \frac{AD}{AB} &= \frac{AE}{AC} \end{aligned}$$

$$\begin{aligned} \text{ii) } \frac{AD}{AB} &= \frac{1.4}{5.6} = \frac{1}{4} \\ \frac{AE}{AC} &= \frac{1.8}{7.2} = \frac{1}{4} \\ \therefore \frac{AD}{AB} &= \frac{AE}{AC} \end{aligned}$$


\therefore By Converse of BPT, $DE \parallel BC$

4. In fig. if $PQ \parallel BC$ and $PR \parallel CD$ prove that

$$\text{(i) } \frac{AR}{AD} = \frac{AQ}{AB} \quad \text{(ii) } \frac{QB}{AQ} = \frac{DR}{AR}$$

Solution :

i) In $\triangle ABC$, $PQ \parallel BC$



$$\therefore \text{By } BPT \frac{AQ}{AB} = \frac{AP}{AC} \quad \dots\dots\dots (1)$$

In $\triangle ADC$, $PR \parallel DC$

$$\therefore \text{By } BPT \frac{AR}{AD} = \frac{AP}{AC} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2),

$$\frac{AQ}{AB} = \frac{AR}{AD}$$

$$ii) \text{ From (i) } \frac{AB}{AQ} = \frac{AD}{AR} \quad (\text{reciprocal})$$

$$\Rightarrow \frac{AB}{AQ} - 1 = \frac{AD}{AR} - 1$$


$$\Rightarrow \frac{AB - AQ}{AQ} = \frac{AD - AR}{AR}$$

$$\Rightarrow \frac{BQ}{AQ} = \frac{DR}{AR}$$

Hence proved.

5. Rhombus PQRB is inscribed in $\triangle ABC$ such that $\angle B$ is one of its angle. P, Q and R lie on AB, AC and BC respectively. If $AB = 12$ cm and $BC = 6$ cm, find the sides PQ, RB of the rhombus.

Solution :



Rhombus PQRS is inscribed in $\triangle ABC$.

Let the side of the rhombus be x .

$$\therefore AB = 12 \text{ cm } AP = 12 - x$$

$$BC = 6 \text{ cm } RC = 6 - x$$

In $\triangle ABC$, $PQ \parallel BC$

$$\therefore \frac{AP}{PB} = \frac{AQ}{QC} \quad \dots\dots\dots (1)$$

In $\triangle ABC$, $QR \parallel AB$

$$\therefore \frac{BR}{RC} = \frac{AQ}{QC} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\Rightarrow \frac{AP}{PB} = \frac{BR}{RC}$$

$$\Rightarrow \frac{12 - x}{x} = \frac{x}{6 - x}$$

$$\Rightarrow x^2 = (6 - x)(12 - x)$$


$$\Rightarrow x^2 = x^2 - 18x + 72$$

$$\Rightarrow 18x = 72$$

$$\Rightarrow x = 4 \text{ cm}$$

6. In trapezium ABCD, $AB \parallel DC$, E and F are points on non-parallel sides AD and BC respectively, such that $EF \parallel AB$. Show that $\frac{AE}{ED} = \frac{BF}{FC}$.

Solution :



In trapezium ABCD, $AB \parallel DC \parallel EF$

Join AC to meet EF at P

In $\triangle ADC$, $EP \parallel DC$

$$\therefore \text{By } BPT, \frac{AE}{ED} = \frac{AP}{PC} \quad \dots\dots\dots (1)$$

In $\triangle ABC$, $PR \parallel AB$

$$\therefore \text{By } BPT, \frac{BF}{FC} = \frac{AP}{PC} \quad \dots\dots\dots (2)$$

From (1) & (2)


$$\frac{AE}{ED} = \frac{BF}{FC}$$

Hence proved.

- 7. In figure $DE \parallel BC$ and $CD \parallel EF$. Prove that $AD^2 = AB \times AF$**

Solution :

In figure $DE \parallel BC$ and $CD \parallel EF$



$$\text{In } \triangle ACD, \text{ by } BPT, \frac{AF}{AD} = \frac{AE}{AC} \quad \dots\dots\dots (1)$$

$$\text{In } \triangle ABC, \text{ by } BPT, \frac{AD}{AB} = \frac{AE}{AC} \quad \dots\dots\dots (2)$$


\therefore From (1) & (2)

$$\frac{AF}{AD} = \frac{AD}{AB}$$

$$\Rightarrow AD^2 = AF \cdot AB$$

- 8. In $\triangle ABC$, AD is the bisector of $\angle A$ meeting side BC at D, if $AB = 10$ cm, $AC = 14$ cm and $BC = 6$ cm, find BD and DC**

Solution :



In $\triangle ABC$, AD is the bisector of $\angle A$.

$$\therefore \text{By } ABT, \frac{AB}{AC} = \frac{BD}{DC}$$

$$\Rightarrow \frac{10}{14} = \frac{x}{6-x}$$

$$\Rightarrow \frac{5}{7} = \frac{x}{6-x}$$

$$\Rightarrow 30 - 5x = 7x$$

$$\Rightarrow 12x = 30$$

$$x = \frac{5}{2} = 2.5$$

$$\therefore BD = 2.5 \text{ cm and } DC = 6 - x = 6 - 2.5$$

$$DC = 3.5 \text{ cm}$$


- 9. Check whether AD is bisector of $\angle A$ of $\triangle ABC$ in each of the following**

(i) $AB = 5$ cm, $AC = 10$ cm, $BD = 1.5$ cm

(ii) $AB = 4$ cm, $AC = 6$ cm, $BD = 1.6$ cm and $CD = 2.4$ cm.

Solution :

i)




$$\frac{AB}{AC} = \frac{5}{10} = \frac{1}{2}, \quad \frac{BD}{DC} = \frac{1.5}{3.5} = \frac{3}{7}$$

$$\therefore \frac{AB}{AC} \neq \frac{BD}{DC}$$

\therefore AD is not the bisector of $\angle A$.

ii)



$$\frac{AB}{AC} = \frac{4}{6} = \frac{2}{3}, \quad \frac{BD}{DC} = \frac{1.6}{2.4} = \frac{2}{3}$$

$$\therefore \frac{AB}{AC} = \frac{BD}{DC}$$


\therefore By Converse of ABT,

AD is the bisector of $\angle A$.

11. ABCD is a quadrilateral in which AB = AD, the bisector of $\angle BAC$ and $\angle CAD$ in-

E and F respectively. Prove that EF || BD.

Solution :



In $\triangle ACD$, AF is the angle bisector

$$\therefore \text{By ABT, } \frac{AD}{AC} = \frac{DF}{FC} \quad \dots\dots\dots (1)$$

In $\triangle ABC$, AE is the angle bisector

$$\begin{aligned} \therefore \text{By ABT, } \frac{AB}{AC} &= \frac{BE}{EC} \\ \Rightarrow \frac{AD}{AC} &= \frac{BE}{EC} \quad \dots (2) \quad (\text{Given } AB = AD) \end{aligned}$$

\therefore From (1) & (2),

$$\frac{BE}{EC} = \frac{DF}{FC}$$

\therefore By Converse of BPT,



$$EF \parallel BD$$

Hence proved.

12. Construct a $\triangle PQR$ which the base PQ = 4.5 cm, $\angle R = 35^\circ$ and the median from R to PQ is 6 cm.

Solution :

Rough Diagram



Construction

Step 1 : Draw a line segment $PQ = 4.5\text{ cm}$.

Step 2 : At P, draw PE such that $\angle QPE = 35^\circ$.

Step 3 : At P, draw PF such that $\angle EPF = 90^\circ$.


Step 4 : Draw the perpendicular bisector to PQ , meets PF at O and PQ at G .

Step 5 : With O as centre and OP as radius draw a circle.

Step 6 : From G mark arcs of 6 cm on the circle at R and S .


Step 7 : Join PR , RQ . Then $\triangle PQR$ is the required \triangle .

Step 8 : Join RG , which is the median.



$\angle P = 40^\circ$ and the median PG from P to QR is 4.4 cm . Find the length of the altitude from P to QR .

Solution :

**Construction**

Step 1 : Draw a line segment $PQ = 5\text{ cm}$.

Step 2 : At P, draw PE such that $\angle QPE = 40^\circ$.

Step 3 : At P, draw PF such that $\angle EPF = 90^\circ$.

Step 4 : Draw the perpendicular bisector to PQ , meets PF at O and PQ at G .

Step 5 : With O as centre and OP as radius draw a circle.


Step 6 : From G mark arcs of 4.4 cm on the circle radius 4.4 cm .


Step 7 : Join PR , RQ . Then $\triangle PQR$ is the required \triangle .

Step 8 : Length of altitude is $RM = 3\text{ cm}$

14. Construct a $\triangle PQR$ such that $QR = 6.5\text{ cm}$, $\angle P = 60^\circ$ and the altitude from P to QR is of length 4.5 cm .

Solution :



**Construction**

Step 1 : Draw a line segment $QR = 6.5$ cm.

Step 2 : At Q, draw QE such that $\angle RQE = 60^\circ$.

Step 3 : At Q, draw QF such that $\angle EQF = 90^\circ$.

Step 4 : Draw the perpendicular bisector XY to QR intersects QF at O & QR at G.

Step 5 : With O as centre and OQ as radius draw a circle.

Step 6 : XY intersects QR at G. On XY, from G, mark arc M such that $GM = 4.5$ cm.



Step 7 : Draw AB, through M which is parallel to QR.

Step 8 : AB meets the circle at P and S.

Step 9 : Join QP, RP. Then $\triangle PQR$ is the required \triangle .

15. Construct a $\triangle ABC$ such that $AB = 5.5$ cm, $\angle C = 25^\circ$ and the altitude from C to AB is 4 cm.

Solution :

**Construction**

Step 1 : Draw a line segment $AB = 5.5$ cm.

Step 2 : At A, draw AE such that $\angle BAE = 25^\circ$.

Step 3 : At A, draw AF such that $\angle EAF = 90^\circ$.

Step 4 : Draw the perpendicular bisector XY to AB intersects AF at O & AB at G.

Step 5 : With O as centre and OA as radius draw a circle.


Step 6 : XY intersects AB at G. On XY, from G, mark arc M such that $GM = 4$ cm.

Step 7 : Draw PQ, through M parallel to AB meets the circle at C and D.

Step 8 : Join AC, BC. Then $\triangle ABC$ is the required \triangle .

16. Draw a triangle ABC of base BC = 5.6 cm, $\angle A = 40^\circ$ and the bisector of $\angle A$ meets BC at D such that CD = 4 cm.

Solution :




Construction

- Step 1 : Draw a line segment BC = 5.6 cm.
 Step 2 : At B, draw BE such that $\angle CBE = 40^\circ$.
 Step 3 : At B, draw BF such that $\angle CBF = 90^\circ$.
 Step 4 : Draw the perpendicular bisector to BC meets BF at O & BC at G.
 Step 5 : With O as centre and OB as radius draw a circle.
 Step 6 : From B, mark an arc of 4 cm on BC at D.
 Step 7 : The \perp r bisector meets the circle at I & Join ID.
 Step 8 : ID produced meets the circle at A. Join AB & AC.
 Step 9 : Then $\triangle ABC$ is the required triangle.

17. Draw $\triangle PQR$ such that PQ = 6.8 cm, vertical angle is 50° and the bisector of the vertical angle meets the base at D where PD = 5.2 cm.

Solution :



Construction


- Step 1 : Draw a line segment PQ = 6.8 cm.
 Step 2 : At P, draw PE such that $\angle QPE = 50^\circ$.
 Step 3 : At P, draw PF such that $\angle QPF = 90^\circ$.
 Step 4 : Draw the perpendicular bisector to PQ meets PF at O and PQ at G.
 Step 5 : With O as centre and OP as radius draw a circle.
 Step 6 : From P mark an arc of 5.2 cm on PQ at D.
 Step 7 : The perpendicular bisector meets the circle at R. Join PR and QR.
 Step 8 : Then $\triangle PQR$ is the required triangle.

III. Pythagoras Theorem :

Theorem 5 : Pythagoras Theorem

Statement

In a right angle triangle, the square on the hypotenuse is equal to the sum of the squares on the other two sides.



Proof

Given : In $\triangle ABC$, $\angle A = 90^\circ$

To prove : $AB^2 + AC^2 = BC^2$

Construction : Draw $AD \perp BC$

No.	Statement	Reason
1.	Compare $\triangle ABC$ and $\triangle ABD$ $\angle B$ is common $\angle BAC = \angle BDA = 90^\circ$ Therefore, $\triangle ABC \sim \triangle ABD$ $\frac{AB}{BD} = \frac{BC}{AB}$ $AB^2 = BC \times BD$ (1)	Given $\angle BAC = 90^\circ$ and by construction By AA similarity
2.	Compare $\triangle ABC$ and $\triangle ADC$ $\angle C$ is common $\angle BAC = \angle ADC = 90^\circ$ Therefore $\triangle ABC \sim \triangle ADC$ $\frac{BC}{AC} = \frac{AC}{DC}$ $AC^2 = BC \times DC$ (2)	Given $\angle BAC = 90^\circ$ and by construction $\angle CDA = 90^\circ$ By AA similarity

Converse of Pythagoras Theorem

Statement

If the square of the longest side of a triangle is equal to sum of squares of other two sides, then the triangle is a right angle triangle.

Example 4.20


An insect 8 m away initially from the foot of a lamp post which is 6 m tall, crawls towards it moving through a distance. If its distance from the top of the lamp post is equal to the distance it has moved, how far is the insect away from the foot of the lamp post?

Solution :

Distance between the insect and the foot of the lamp post

$$BD = 8 \text{ m}$$

The height of the lamp post, $AB = 6 \text{ m}$



After moving a distance of $x \text{ m}$, let the insect be at C

Let, $AC = CD = x$. Then $BC = BD - CD = 8 - x$

In $\triangle ABC$, $\angle B = 90^\circ$

$$AC^2 = AB^2 + BC^2 \text{ gives } x^2 = 6^2 + (8 - x)^2$$

$$x^2 = 36 + 64 - 16x + x^2$$

$$16x = 100 \text{ then } x = 6.25$$


$$\text{Then, } BC = 8 - x = 8 - 6.25 = 1.75 \text{ m}$$

Therefore the insect is 1.75 m away from the foot of the lamp post.

Example 4.21

P and Q are the mid-points of the sides CA and CB respectively of a $\triangle ABC$, right angled at C. Prove that $4 (AQ^2 + BP^2) = 5 AB^2$.

Solution :



Since, $\triangle AQC$ is a right triangle at C,

$$AQ^2 = AC^2 + QC^2 \quad \dots\dots\dots (1)$$

Also, $\triangle BPC$ is a right triangle at C,

$$BP^2 = BC^2 + CP^2 \quad \dots\dots\dots (2)$$

From (1) and (2), $AQ^2 + BP^2 =$

$$AC^2 + QC^2 + BC^2 + CP^2$$

$$\begin{aligned} 4 (AQ^2 + BP^2) &= 4AC^2 + 4QC^2 + 4BC^2 + 4CP^2 \\ &= 4AC^2 + (2QC)^2 + 4BC^2 + (2CP)^2 \\ &= 4AC^2 + BC^2 + 4BC^2 + AC^2 \\ (\text{Since P and Q are mid points}) \\ &= 5 (AC^2 + BC^2) \end{aligned}$$

$$4 (AQ^2 + BP^2) = 5 AB^2$$


(By Pythagoras Theorem)

Example 4.22

What length of ladder is needed to reach a height of 7 ft along the wall when the base of the ladder is 4 ft from the wall? Round off your answer to the next tenth place.

Solution :

Let x be the length of the ladder. $BC = 4$ ft, $AC = 7$ ft.



By Pythagoras theorem we have, $AB^2 = AC^2 + BC^2$

$$x^2 = 7^2 + 4^2 \text{ gives } x^2 = 49 + 16$$

$$x^2 = 65. \text{ Hence, } x = \sqrt{65}$$

The number $\sqrt{65}$ is between 8 and 8.1.


$$8^2 = 64 < 65 < 65.61 = 8.1^2$$

Therefore, the length of the ladder is ap-

Example 4.23

An Aeroplane leaves an airport and flies due north at a speed of 1000 km/hr. At the same time, another aeroplane leaves the same airport and flies due west at a speed of 1200 km/hr. How far apart will be the two planes after $1\frac{1}{2}$ hours?

Solution :



Let the first aeroplane starts from O and goes upto A towards north, (Distance = Speed \times time)

$$\text{where } OA = \left(1000 \times \frac{3}{2} \right) \text{ km} = 1500 \text{ km}$$

Let the second aeroplane starts from O at the same time and

goes upto B towards west,

$$\text{where } OB = \left(1200 \times \frac{3}{2}\right) \text{ km} = 1800 \text{ km}$$

The required distance to be found is BA.

In right angled triangle AOB, $AB^2 = OA^2 + OB^2$

$$AB^2 = (1500)^2 + (1800)^2 = 100^2 (15^2 + 18^2)$$


$$= 100^2 \times 549 = 100^2 \times 9 \times 61$$

$$AB = 100 \times 3 \times \sqrt{61} = 300\sqrt{61} \text{ kms.}$$

EXERCISE 4.3

1. A man goes 18 m due east and then 24 m due north. Find the distance of his current position from the starting point.

Solution:




P → Starting Point

By Pythagoras Theorem,


$$\begin{aligned} PN &= \sqrt{18^2 + 24^2} \\ &= \sqrt{324 + 576} \\ &= \sqrt{900} \\ &= 30 \text{ m} \end{aligned}$$

∴ Distance of his current position from the starting point = 30 m

2. There are two paths that one can choose to go from Sarah's house to James house. One way is to take C street, and the other way requires to take A street and then B street. How much shorter is the direct path along C street? (Using figure).



Solution:



Path - 1 (Direct C Street)

$$\begin{aligned} SJ &= \sqrt{(1.5)^2 + 2^2} \\ &= \sqrt{2.25 + 4} \\ &= \sqrt{6.25} \\ &= 2.5 \text{ miles} \end{aligned}$$

Path = 2 (B Street & then A Street)


$$\begin{aligned} SP + PJ &= 1.5 + 2 \\ &= 3.5 \text{ miles} \end{aligned}$$

$$\therefore \text{Required} = 3.5 - 2.5$$

$$= 1 \text{ mile}$$

∴ 1 mile is shorter along C Street.

3. To get from point A to point B you must avoid walking through a pond. You must walk 34 m south and 41 m east. To the nearest meter, how many meters would be saved if it were possible to make a way through the pond?

Solution:

Path - 1 (Through pond)

$$\begin{aligned}
 AB &= \sqrt{34^2 + 41^2} \\
 &= \sqrt{1156 + 1681} \\
 &= \sqrt{2837} \\
 &= 53.26 \text{ m}
 \end{aligned}$$

Path - 2 (South & then East)


Total dist. covered

$$\begin{aligned}
 &= 34 + 41 \\
 &= 75 \text{ m}
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{Reqd. time saving} &= 75 - 53.26 \\
 &= 21.74 \text{ m}
 \end{aligned}$$

4. In the rectangle WXYZ, $XY + YZ = 17$ cm, and $XZ + YW = 26$ cm.

Calculate the length and breadth of the rectangle?

**Solution:**

$$\text{Given } xy + yz = 17$$

$$\Rightarrow l + b = 17 \quad \dots\dots (1)$$

$$xz + yw = 26$$

$$\Rightarrow \sqrt{l^2 + b^2} + \sqrt{l^2 + b^2} = 26$$

$$\Rightarrow 2\sqrt{l^2 + b^2} = 26$$

$$\therefore l^2 + b^2 = 169$$

$$\Rightarrow l^2 + (17 - l)^2 = 169 \quad (\text{From (1)})$$

$$\Rightarrow 2l^2 - 34l + 289 = 169$$

$$\Rightarrow 2l^2 - 34l + 120 = 0$$

$$\Rightarrow l^2 - 17l + 60 = 0$$

$$\Rightarrow (l - 12)(l - 5) = 0$$

$$\therefore l = 12, l = 5$$


But $l = 12$ only

$$\therefore (1) \Rightarrow b = 17 - 12$$

$$= 5$$

$$\therefore \text{Length} = 12 \text{ cm, Breadth} = 5 \text{ m}$$

5. The hypotenuse of a right triangle is 6 m more than twice of the shortest side. If the third side is 2 m less than the hypotenuse, find the sides of the triangle.

Solution:Let x be the shortest side. z be the hypotenuse & y be the 3rd side.

$$\begin{aligned}
 \text{By data given, } z &= 2x + 6, & y &= z - 2 \\
 & & &= 2x + 6 - 2 \\
 & & &= 2x + 4
 \end{aligned}$$

In $\triangle ABC$,


By Pythagoras theorem,

$$x^2 + y^2 = z^2$$

$$\begin{aligned}
 \Rightarrow x^2 + (2x + 4)^2 &= (2x + 6)^2 \\
 \Rightarrow x^2 + 4x^2 + 16x + 16 &= 4x^2 + 24x + 36 \\
 \Rightarrow x^2 - 8x - 20 &= 0 \\
 \Rightarrow (x - 10)(x + 2) &= 0 \\
 \therefore x &= 10 \text{ m} \\
 \therefore y &= 2(10) + 4 = 24 \text{ m} \\
 z &= 2(10) + 6 = 26 \text{ m} \\
 \therefore \text{The length of 3 sides are} & \\
 10\text{m, } 24\text{m, } 26\text{m} &
 \end{aligned}$$

6. 5 m long ladder is placed leaning towards a vertical wall such that it reaches the wall at a point 4m high. If the foot of the ladder is moved 1.6 m towards the wall, then find the distance by which the top of the ladder would slide upwards on the wall.

Solution:



$$\begin{aligned}
 \text{In } \triangle ABC, AB &= \sqrt{5^2 - 4^2} \\
 &= \sqrt{25 - 16} \\
 &= \sqrt{9} \\
 &= 3
 \end{aligned}$$

$$\text{Given } AD = 1.6\text{m}$$

$$\begin{aligned}
 \Rightarrow DB &= 3 - 1.6 \\
 &= 1.4 \text{ m}
 \end{aligned}$$

Now, In $\triangle DEB$,


$$\begin{aligned}
 EB &= \sqrt{5^2 - (1.4)^2} \\
 &= \sqrt{25 - 1.96} \\
 &= \sqrt{23.04} \\
 &= 4.8
 \end{aligned}$$

$$\therefore EC = EB - BC = 4.8 - 4 = 0.8 \text{ m}$$

\therefore When the foot of the ladder moves 1.6 m towards the wall, the top of the ladder will slide upwards at a dist of 0.8 m

7. The perpendicular PS on the base QR of a $\triangle PQR$ intersects QR at S, such that $QS = 3 \text{ SR}$. Prove that $2PQ^2 = 2PR^2 + QR^2$.

Solution:



$$\text{Given } QS = 3 \cdot SR$$

$$\text{To Prove : } 2PQ^2 = 2PR^2 + QR^2$$

$$= 3SR + SR$$

$$QR = 4SR \Rightarrow SR = \frac{1}{4} QR$$

$$\text{In } \triangle PQS, PQ^2 = PS^2 + QS^2 \quad \dots (1)$$

$$\text{In } \triangle PRS, PR^2 = PS^2 + SR^2 \quad \dots (2)$$

$$\begin{aligned}
 (1) - (2) \Rightarrow PQ^2 - PR^2 &= QS^2 - SR^2 \\
 &= (3 \cdot SR)^2 - SR^2 \\
 &= 8 \cdot SR^2 \\
 &= 8 \left(\frac{1}{4} QR \right)^2 \\
 &= 8 \left(\frac{1}{16} QR^2 \right)
 \end{aligned}$$


$$PQ^2 - PR^2 = \frac{QR^2}{2}$$

$$\Rightarrow 2PQ^2 - 2PR^2 = QR^2$$


$$\Rightarrow 2PQ^2 = 2PR^2 + QR^2$$

Hence proved.

8. In the adjacent figure, ABC is a right angled triangle with right angle at B and points D, E trisect BC. Prove that $8AE^2 = 3AC^2 + 5AD^2$



Solution:



$$\therefore BD = DE = EC = x \text{ (take)}$$

$$\therefore BD = x, BE = 2x, BC = 3x$$

$$\begin{aligned} \text{In } \triangle ABD, AD^2 &= AB^2 + BD^2 \\ &= AB^2 + x^2 \end{aligned}$$

$$\begin{aligned} \text{In } \triangle ABE, AE^2 &= AB^2 + BE^2 \\ &= AB^2 + (2x)^2 \\ &= AB^2 + 4x^2 \end{aligned}$$

$$\begin{aligned} \text{In } \triangle ABC, AC^2 &= AB^2 + BC^2 \\ &= AB^2 + (3x)^2 \\ &= AB^2 + 9x^2 \end{aligned}$$

$$\text{To prove : } 8AE^2 = 3AC^2 + 5AD^2$$

RHS :

$$\begin{aligned} 3AC^2 + 5AD^2 &= 3(AB^2 + 9x^2) + 5(AB^2 + x^2) \\ &= 3AB^2 + 27x^2 + 5AB^2 + 5x^2 \\ &= 8AB^2 + 32x^2 \\ &= 8(AB^2 + 4x^2) \end{aligned}$$

2

$$= \text{LHS}$$

Hence proved.

IV. Circles and Tangents

Key Points


- ✓ If a line touches the given circle at only one point, then it is called tangent to the circle.
- ✓ A tangent at any point on a circle and the radius through the point are perpendicular to each other.
- ✓ No tangent can be drawn from an interior point of the circle.
- ✓ Only one tangent can be drawn at any point on a circle.
- ✓ Two tangents can be drawn from any exterior point of a circle.
- ✓ The lengths of the two tangents drawn from an exterior point to a circle are equal,
- ✓ If two circles touch externally the distance between their centers is equal to the sum of their radii.
- ✓ If two circles touch internally, the distance between their centers is equal to the difference of their radii.
- ✓ The two direct common tangents drawn to the circles are equal in length.

Theorem 6 : Alternate Segment theorem**Statement**

If a line touches a circle and from the point of contact a chord is drawn, the angles between the tangent and the chord are respectively equal to the angles in the corresponding alternate segments.

Proof

Given : A circle with centre at O, tangent AB touches the circle at P and PQ is a chord. S and T are two points on the circle in the opposite sides of chord PQ.



To prove : (i) $\angle QPB = \angle PSQ$ and

(ii) $\angle QPA = \angle PTQ$

Construction : Draw the diameter POR. Draw QR, QS and PS.

No.	Statement	Reason
1.	$\angle RPB = 90^\circ$ Now, $\angle RPQ + \angle QPB = 90^\circ$...(1)	Diameter RP is perpendicular to tangent AB.
2.	In $\triangle RPQ$, $\angle PQR = 90^\circ$...(2)	Angle in a semi-circle is 90°
3.	$\angle QRP + \angle RPQ = 90^\circ$(3)	In a right angled triangle, sum of the two acute angles is 90° .


4.	$\angle RPQ + \angle QPB = \angle QRP + \angle RPQ$ $\angle QPB = \angle QRP$(4)	From (1) and (3).
5.	$\angle QRP = \angle PSQ$(5)	Angles in the same segment are equal.
6.	$\angle QPB = \angle PSQ$(6)	From (4) and (5) ; Hence (i) is proved.
7.	$\angle QPB + \angle QPA = 180^\circ$(7)	Linear pair of angles.
8.	$\angle PSQ + \angle PTQ = 180^\circ$(8)	Sum of opposite angles of a cyclic quadrilateral is 180° .
9.	$\angle QPB + \angle QPA = \angle PSQ + \angle PTQ$	From (7) and (8).
10.	$\angle QPB + \angle QPA = \angle QPB + \angle PTQ$	$\angle QPB = \angle PSQ$ from (6)
		This completes the proof.

Example 4.24

Find the length of the tangent drawn from a point whose distance from the centre of a circle is 5 cm and radius of the circle is 3 cm.

Solution :

Given $OP = 5$ cm, radius $r = 3$ cm



To find the length of tangent PT.

In right angled $\triangle OTP$,

$OP^2 = OT^2 + PT^2$ (by Pythagoras theorem)

$5^2 = 3^2 + PT^2$ gives $PT^2 = 25 - 9 = 16$

Length of the tangent $PT = 4$ cm


Example 4.25

PQ is a chord of length 8 cm to a circle of radius 5 cm. The tangents at P and Q intersect at a point T. Find the length of the tangent TP.

Solution :

Let $TR = y$. Since, OT is perpendicular bisector of PQ.

$$PR = QR = 4 \text{ cm}$$



$$\text{In } \triangle ORP, OP^2 = OR^2 + PR^2$$

$$OR^2 = OP^2 - PR^2$$

$$OR^2 = 5^2 - 4^2 = 25 - 16 = 9 \Rightarrow OR = 3 \text{ cm}$$

$$OT = OR + RT = 3 + y \quad \dots\dots\dots (1)$$

$$\text{In } \triangle PRT, TP^2 = TR^2 + PR^2 \quad \dots\dots\dots (2)$$

and $\triangle OPT$ we have, $OT^2 = TP^2 - OP^2$

$$OT^2 = (TR^2 + PR^2) + OP^2$$

(substitute for TP^2 from (2))

$$(3+y)^2 = y^2 + 4^2 + 5^2 \text{ (substitute for OT from (1))}$$

$$9 + 6y + y^2 = y^2 + 16 + 25$$

$$\text{Therefore } y = TR = \frac{16}{3}$$


$$6y = 41 - 9 \text{ we get } y = \frac{16}{3}$$

$$\text{From (2), } TP^2 = TR^2 + PR^2$$

$$TP^2 = \left(\frac{16}{3}\right)^2 + 4^2 = \frac{256}{9} + 16 = \frac{400}{9} \text{ so, } TP = \frac{20}{3} \text{ cm}$$

Example 4.26

In Figure O is the centre of a circle. PQ is a chord and the tangent PR at P makes an angle of 50° with PQ. Find $\angle POQ$

**Solution :**

$\angle OPQ = 90^\circ - 50^\circ = 40^\circ$ (angle between the radius and tangent is 90°)

$OP = OQ$ (Radii of a circle are equal)


$\angle OPQ = \angle OQP = 40^\circ$ ($\triangle OPQ$ is isosceles)

$$\angle POQ = 180^\circ - \angle OPQ - \angle OQP$$

$$0 \quad 0 \quad 0 \quad 0$$

Example 4.27

In Fig., $\triangle ABC$ is circumscribing a circle. Find the length of BC.

**Solution :**

$AN = AM = 3 \text{ cm}$ (Tangents drawn from same external point are equal)


$BN = BL = 4 \text{ cm}$

$CL = CM = AC - AM = 9 - 3 = 6 \text{ cm}$

Gives $BC = BL + CL = 4 + 6 = 10 \text{ cm}$

Example 4.28

If radii of two concentric circles are 4 cm and 5 cm then find the length of the chord of one circle which is a tangent to the other circle.



Solution :

OA = 4 cm, OB = 5 cm; also $OA \perp BC$.

$$OB^2 = OA^2 + AB^2$$

$$5^2 = 4^2 + AB^2 \text{ gives } AB^2 = 9$$

Therefore $AB = 3$ cm

$BC = 2AB$ hence $BC = 2 \times 3 = 6$ cm



CONSTRUCTION OF A TANGENT TO A CIRCLE

Example 4.29

Draw a circle of radius 3 cm. Take a point P on this circle and draw a tangent at P.

Solution :

Given, radius $r = 3$ cm

**Construction**

Step 1 : Draw a circle with centre at O of radius 3 cm.

Step 2 : Take a point P on the circle. Join OP.

Step 3 : Draw perpendicular line TT' to OP which


Step 4 : TT' is the required tangent.


Example 4.30

Draw a circle of radius 4 cm. At a point L on it draw a tangent to the circle using the alternate segment.

Solution :

Given, radius = 4 cm



**Construction**

radius 4 cm.

Step 2 : Take a point L on the circle. Through L draw any chord LM.

Step 3 : Take a point M distinct from L and N on the circle, so that L, M and N are in anti-clockwise direction. Join LN and NM.

Step 4 : Through L draw a tangent TT' such that $\angle TLM = \angle MNL$.

Step 5 : TT' is the required tangent.



Example 4.31

Draw a circle of diameter 6 cm from a point P, which is 8 cm away from its centre. Draw the two tangents PA and PB to the circle and measure their lengths.

Solution :

Given, diameter (d) = 6 cm, we find radius

$$(r) = \frac{6}{2} = 3 \text{ cm.}$$

**Construction**

Step 1 : With centre at O, draw a circle of radius 3 cm.

Step 2 : Draw a line OP of length 8 cm.

Step 3 : Draw a perpendicular bisector of OP, which cuts OP at M.

Step 4 : With M as centre and MO as radius, draw a circle which cuts previous circle at A and B.

Step 5 : Join AP and BP. AP and BP are the required tangents. Thus length of the tangents are $PA = PB = 7.4 \text{ cm}$.


Verification : In the right angle triangle OAP , $PA^2 = OP^2 - OA^2 = 64 - 9 = 55$

$$PA = \sqrt{55} = 7.4 \text{ cm (approximately) .}$$

Concurrency Theorems

Key Points

- ✓ A cevian is a line segment that extends from one vertex of a triangle to the opposite side.
- ✓ A median is a cevian that divides the opposite side into two congruent(equal) lengths.
- ✓ An altitude is a cevian that is perpendicular to the opposite side.
- ✓ An angle bisector is a cevian that bisects the corresponding angle.
- ✓ Ceva's Theorem : Let ABC be a triangle and let D,E,F be points on lines BC, CA, AB respectively. Then the cevians AD, BE, CF are concurrent if and only if $\frac{BD}{DC} \times \frac{CE}{EA} \times \frac{AF}{FB} = 1$ where the lengths are directed.
- ✓ Menelaus Theorem : A necessary and sufficient condition for points P, Q, R on the respective sides BC, CA, AB (or their extension) of a triangle ABC to be collinear is that $\frac{BP}{PC} \times \frac{CQ}{QA} \times \frac{AR}{RB} = 1$ where all segments in the formula are directed segments.



Example 4.32

Show that in a triangle, the medians are concurrent.

Solution :

Medians are line segments joining each vertex to the midpoint of the corresponding opposite sides.

Thus medians are the cevians where D, E, F are midpoints of BC, CA and AB respectively.

Since D is a mid point of

$$BC, BD = DC \text{ so } \frac{BD}{DC} = 1 \quad \dots\dots\dots (1)$$

Since, E is a midpoint of


$$CA, CE = EA \text{ so } \frac{CE}{EA} = 1 \quad \dots\dots\dots (2)$$

Since, F is a midpoint of AB,

$$AB, AF = FB \text{ so } \frac{AF}{FB} = 1 \quad \dots\dots\dots (3)$$

Thus, multiplying (1), (2) and (3) we get,

$$\frac{BD}{DC} \times \frac{CE}{EA} \times \frac{AF}{FB} = 1 \times 1 \times 1 = 1$$




And so, Ceva's theorem is satisfied.

Hence the Medians are concurrent.

Example 4.33

In Fig., ABC is a triangle with $\angle B = 90^\circ$, $BC = 3$ cm and $AB = 4$ cm. D is point on AC such that $AD = 1$ cm and E is the midpoint of AB. Join D and E and extend DE to meet CB at F. Find BF.



Solution :

Consider $\triangle ABC$. Then D, E and F are respective points on the sides CA, AB and BC. By construction D, E, F are collinear.

$$\text{By Menelaus' theorem } \frac{AE}{EB} \times \frac{BF}{FC} \times \frac{CD}{DA} = 1 \quad \dots (1)$$

By assumption, $AE = EB = 2$, $DA = 1$ and

$$FC = FB + BC = BF + 3$$

By Pythagoras theorem, $AC^2 = AB^2 + BC^2$

$$= 16 + 9 = 25. \text{ Therefore } AC = 5 \text{ and So, } CD = AC - AD = 5 - 1 = 4.$$

Substituting the values of FC, AE, EB, DA, CD in (1),

$$\text{we get, } \frac{2}{2} \times \frac{BF}{BF+3} \times \frac{4}{1} = 1$$

$$4BF = BF + 3$$

$$4BF - BF = 3 \text{ therefore } BF = 1$$


Example 4.34

Suppose AB, AC and BC have lengths 13, 14 and 15 respectively. If $\frac{AF}{FB} = \frac{2}{5}$ and $\frac{CE}{EA} = \frac{5}{8}$. Find BD and DC.

Solution :

Given that AB = 13, AC = 14 and BC = 15

Let BD = x and DC = y



Using Ceva's theorem, we have,

$$\frac{BD}{DC} \times \frac{CE}{EA} \times \frac{AF}{FB} = 1 \quad \dots\dots\dots (1)$$

Substitute the values of $\frac{AF}{FB}$ and $\frac{CE}{EA}$ in (1),

$$\text{We have } \frac{BD}{DC} \times \frac{5}{8} \times \frac{2}{5} = 1$$

$$\frac{x}{y} \times \frac{10}{40} = 1 \text{ we get, } \frac{x}{y} \times \frac{1}{4} = 1.$$

$$\text{Hence, } x = 4y \quad \dots\dots\dots (2)$$

$$BC = BD + DC = 15$$

$$\text{so, } x + y = 15 \quad \dots\dots\dots (3)$$


From (2), using $x = 4y$ in (3) we get, $4y + y = 15$ gives $5y = 15$ then $y = 3$

Substitute $y = 3$ in (3) we get, $x = 12$.

Hence BD = 12, DC = 3.

Example 3.7

In a garden containing several trees, three particular trees P, Q, R are located in the following way, BP = 2 m, CQ = 3 m, RA = 10 m, PC = 6 m, QA = 5 m, RB = 2 m, where A, B, C are points such that P lies on BC, Q lies on AC and R lies on AB. Check whether the trees P, Q, R lie on a same straight line.



Solution :

By Menelaus's theorem, the trees P, Q, R will be collinear (lie on same straight line)

$$\text{if } \frac{BP}{PC} \times \frac{CQ}{QA} \times \frac{RA}{RB} = 1 \quad \dots\dots\dots (1)$$

Given BP = 2 m, CQ = 3 m, RA = 10 m, PC = 6 m, QA = 5 m and RB = 2 m

Substituting these values in (1) we get,


$$\frac{BP}{PC} \times \frac{CQ}{QA} \times \frac{RA}{RB} = \frac{2}{6} \times \frac{3}{5} \times \frac{10}{2} = \frac{60}{60} = 1$$

Hence the trees P, Q, R lie on a same straight line.

EXERCISE 4.4

1. The length of the tangent to a circle from a point P, which is 25 cm away from the centre is 24 cm. What is the radius of the circle?

Solution:



Given $OP = 25$ cm, $PT = 24$ cm


Radius & Tangent are Perpendicular

$$\begin{aligned}\therefore OT &= \sqrt{OP^2 - PT^2} \\ &= \sqrt{2^2 - 2^2} \\ &= \sqrt{625 - 576} \\ &= \sqrt{49} \\ &= 7 \text{ cm}\end{aligned}$$

\therefore Radius = 7 cm

2. $\triangle LMN$ is a right angled triangle with $\angle L = 90^\circ$. A circle is inscribed in it. The lengths of the sides containing the right angle are 6 cm and 8 cm. Find the radius of the circle.

Solution:



$$\begin{aligned}MN &= \sqrt{ML^2 + LN^2} \\ &= \sqrt{64 + 36} \\ &= \sqrt{100} \\ &= 10\end{aligned}$$

Area of $\triangle MLN$ = Area of $\triangle MOL$ +

Area of $\triangle NOL$ + Area of $\triangle MON$

$$\begin{aligned}\Rightarrow \frac{1}{2} \times ML \times LN &= \frac{1}{2} \times LM \times r + \frac{1}{2} \times LN \times r \\ &\quad + \frac{1}{2} \times MN \times r\end{aligned}$$

$$\Rightarrow 8 \times 6 = 8r + 6r + 10r$$

$$\Rightarrow 24r = 48$$

$$r = 2 \text{ cm}$$

3. A circle is inscribed in $\triangle ABC$ having sides 8 cm, 10 cm and 12 cm as shown in figure, Find AD, BE and CF.


Solution:

From the fig,

$$x + y = 12$$

$$y + z = 8$$

$$z + x = 10$$



Adding, $2(x + y + z) = 30$

$$\Rightarrow x + y + z = 15$$

$$\Rightarrow 12 + z = 15$$

$$\therefore z = 3$$

$$\Rightarrow y + 3 = 8$$


$$\Rightarrow y = 5$$

$$\therefore x + 5 = 12$$

$$\Rightarrow x = 7$$


$$\therefore AD = 7 \text{ cm, } BE = 5 \text{ cm, } CF = 3 \text{ cm}$$

4. PQ is a tangent drawn from a point P to a circle with centre O and QOR is a diameter of the circle such that $\angle POR = 120^\circ$. Find $\angle OPQ$.

Solution:

Given $\angle POR = 120^\circ$
 $\Rightarrow \angle POQ = 60^\circ$ (linear pair)
 Also $\angle OQP = 90^\circ$ (Radius \perp tangent)
 $\therefore \angle OPQ = 90^\circ - 60^\circ$
 $= 30^\circ$

5. A tangent ST to a circle touches it at B . AB is a chord such that $\angle ABT = 65^\circ$. Find $\angle AOB$, where “O” is the centre of the circle.

Solution:


Given $\angle TBA = 65^\circ \Rightarrow \angle APB = 65^\circ$
 (angles in alternate segment).
 $\therefore \angle AOB = 2\angle APB = 2(65^\circ) = 130^\circ$ (circumference)
 (Angle subtended at the centre is twice the angle subtended at any point on the remaining

6. In figure, O is the centre of the circle with radius 5 cm. T is a point such that $OT = 13$ cm and OT intersects the circle E, if AB is the tangent to the circle at E, find the length of AB .

Solution:

In the figure, given $OP = 5$, $OT = 13$

$$\begin{aligned}\therefore PT &= \sqrt{13^2 - 5^2} \\ &= \sqrt{169 - 25} \\ &= \sqrt{144} \\ &= 12 \\ &= TQ\end{aligned}$$



Also, $OE = 5 \Rightarrow ET = 13 - 5 = 8$

Let $AP = AE = x \Rightarrow TA = 12 - x$

\therefore In $\triangle AET$, $\angle AET = 90^\circ$

$$\therefore x^2 + 8^2 = (12 - x)^2$$

$$\Rightarrow x^2 + 8^2 = 144 + x^2 - 24x$$

$$\Rightarrow 64 = 144 - 24x$$


$$\Rightarrow 24x = 80$$

$$x = \frac{10}{3}$$

$$\therefore AB = 2x$$

$$\text{Length of tangent } AB = \frac{20}{3} \text{ cm}$$

7. In two concentric circles, a chord of length 16 cm of larger circle becomes a tangent to the smaller circle whose radius is 6 cm. Find the radius of the larger circle.

Solution:

Given the chord AB of larger circle is a tangent for the smaller circle & OT is radius.

OT is perpendicular to AB.

$\therefore AT = TB = 8 \text{ cm}, OT = 6 \text{ cm}$


\therefore In ΔOBT ,

$$\begin{aligned} OB &= \sqrt{8^2 + 6^2} \\ &= \sqrt{64 + 36} \\ &= \sqrt{100} \\ &= 10 \text{ cm} \end{aligned}$$

\therefore Radius of the larger circle = 10 cm

8. Two circles with centres O and O' of radii 3 cm and 4 cm, respectively intersect at two points P and Q, such that OP and O'P are tangents to the two circles. Find the length of the common chord PQ.

Solution:



Given $OP = 4 \text{ cm}$ (radius of 1st circle)

$O'P = 3 \text{ cm}$ (radius of 2nd circle)

Clearly $OP \perp O'P$ (tangent & radius are \perp)

$$\begin{aligned} \therefore OO' &= \sqrt{4^2 + 3^2} \\ &= \sqrt{16 + 9} \\ &= \sqrt{25} \\ &= 5 \end{aligned}$$

Let R be a point of PQ such that

$OR = x$ & $O'R = 5 - x$

Also, $\Delta OPO' \sim \Delta OQO'$ & $\Delta OPR \sim \Delta OQR$ (by similarity)

$\therefore \angle ORP = 90^\circ$

\therefore In ΔORP , $PR^2 = 16 - x^2$

In $\Delta O'RP$, $PR^2 = 9 - (5 - x)^2$

$\therefore 16 - x^2 = 9 - (5 - x)^2$

$\Rightarrow 10x = 32$

$$\therefore x = \frac{16}{5}$$

$$\therefore PR = \sqrt{16 - \frac{256}{25}}$$

$$= \sqrt{\frac{144}{25}}$$

$$= \frac{12}{5}$$

$$= 2.4$$


$\therefore PQ = 2(PR)$

$$= 2(2.4)$$

$$= 4.8 \text{ cm}$$

9. Show that the angle bisectors of a triangle are concurrent.

Solution:



Consider a ΔABC & let the angular bisectors of A and B meet at 'O'.

From O, draw perpendicular OD, OE, OF to BC, CA, AB respectively.

Now $\Delta BOD \equiv \Delta BOF$

($\because \angle ODB = \angle OFB = 90^\circ$)

$\therefore OD = OF$ ($\angle OBD = \angle OBF$)

Similarly in ΔOAE & ΔOAF , we can prove

$$OE = OF$$

$$\therefore OD = OE = OF$$

Now, join OC,

Consider $\triangle OCD$, $\triangle OCE$

Here i) $\angle ODC = \angle OEC = 90^\circ$ & OC is common

$$\text{ii) } OD = OE$$

$$\therefore \triangle OCD = \triangle OCE$$


$$\therefore \angle OCD = \angle OCE$$

CO is angle bisector of $\angle C$.

\therefore Angle bisectors of a triangle are concurrent.

10. In $\triangle ABC$, with $B = 90^\circ$, $BC = 6$ cm and $AB = 8$ cm, D is a point on AC such that $AD = 2$ cm and E is the midpoint of AB. Join D to E and extend it to meet at F. Find BF.

Solution:



Given In $\triangle ABC$, $AB = 8$ cm, $BC = 6$ cm

$$\therefore AC = \sqrt{64 + 36} = \sqrt{100} = 10$$

Also $AD = 2 \Rightarrow CD = 8$ cm

E is the mid point of AB

$$\Rightarrow AE = EB = 4 \text{ cm}$$

By Menelaus Theorem,

$$\frac{AE}{EB} \times \frac{BF}{FC} \times \frac{CD}{DA} = 1$$


$$\Rightarrow \frac{4}{4} \times \frac{BF}{BF+6} \times \frac{8}{2} = 1$$

$$\Rightarrow 4BF = BF + 6$$

$$\Rightarrow 3BF = 6$$

$$\therefore BF = 2 \text{ cm}$$

11. An artist has created a triangular stained glass window and has one strip of small length left before completing the window. She needs to figure out the length of left out portion based on the lengths of the other sides as shown in the figure.



Solution:

By applying Ceva's theorem, the Cevians AD, BE and CF intersect at exactly one point if and only if

$$BD \times CE \times AF = DC \times EA \times FB$$

$$\Rightarrow 3 \times 4 \times 5 = 10 \times 3 \times FB$$


$$\Rightarrow 60 = 30 \times FB$$

$$\therefore FB = 2 \text{ cm}$$

12. Draw a tangent at any point R on the circle of radius 3.4 cm and centre at P?

Solution:

Rough Diagram



Construction

Step 1 : Draw a circle with centre at P of radius 3.4 cm.


Step 2 : Take a point R on the circle and Join PR.

Step 3 : Draw perpendicular line TT' to PR which passes through R.

Step 4 : TT' is the required tangent.

13. Draw a circle of radius 4.5 cm. Take a point on the circle. Draw the tangent at that point using the alternate segment theorem.

Solution:

**Construction**

Step 1 : With O as the centre, draw a circle of radius 4.5 cm.

Step 2 : Take a point L on the circle. Through L draw any chord LM.


Step 3 : Take a point M distinct from L and N on the circle, so that L, M and N are in anti-clockwise direction. Join LN and NM.

Step 4 : Through L draw a tangent TT' such that $\angle TLM = \angle MNL$.

Step 5 : TT' is the required tangent.

14. Draw the two tangents from a point which is 10 cm away from the centre of a circle of radius 5 cm. Also, measure the lengths of the tangents.

Solution:

**Construction**

Step 1 : With centre at O, draw a circle of radius 5 cm.

Step 2 : Draw a line OP = 10 cm.

Step 3 : Draw a perpendicular bisector of OP, which cuts OP at M.

Step 4 : With M as centre and MO as radius, draw a circle which cuts previous circle at A and B.


Step 5 : Join AP and BP. AP and BP are the required tangents. Thus length of the tangents are $PA = PB = 8.7$ cm.

Verification : In the right angle triangle $\triangle OAP$,

$$\begin{aligned} PA^2 &= \sqrt{OP^2 - OA^2} \\ &= \sqrt{100 - 25} = \sqrt{75} = 8.7 \text{ cm} \end{aligned}$$

15. Take a point which is 11 cm away from the centre of a circle of radius 4 cm and draw the two tangents to the circle from that point.

Solution:



Construction

Step 1 : With centre at O, draw a circle of radius 4 cm.

Step 2 : Draw a line $OP = 11$ cm.

Step 3 : Draw a perpendicular bisector of OP, which cuts OP at M.

Step 4 : With M as centre and MO as radius, draw a circle which cuts previous circle at A and B.


Step 5 : Join AP and BP. They are the required tangents $AP = BP = 10.3$ cm.

Verification : In the right angle triangle $\triangle OAP$,

$$\begin{aligned} AP &= \sqrt{OP^2 - OA^2} \\ &= \sqrt{121 - 16} = \sqrt{105} = 10.3 \text{ cm} \end{aligned}$$

16. Draw the two tangents from a point which is 5 cm away from the centre of a circle of diameter 6 cm. Also, measure the lengths of the tangents.

Solution:



Construction

Step 1 : With centre at O, draw a circle of radius 3 cm.

Step 2 : Draw a line $OP = 5$ cm.

Step 3 : Draw a perpendicular bisector of OP, which cuts OP at M.

Step 4 : With M as centre and OM as radius, draw a circle which cuts previous circle at A and B.


Step 5 : Join AP and BP. They are the required tangents $AP = BP = 4$ cm.

Verification :

$$\begin{aligned} AP &= \sqrt{OP^2 - OA^2} \\ &= \sqrt{5^2 - 3^2} \\ &= \sqrt{25 - 9} \\ &= \sqrt{16} = 4 \text{ cm} \end{aligned}$$

17. Draw a tangent to the circle from the point P having radius 3.6 cm, and centre at O. Point P is at a distance 7.2 cm from the centre.

Solution:

**Construction**

Step 1 : Draw a circle of radius 3.6 cm. with centre at O.

Step 2 : Draw a line $OP = 7.2$ cm.

Step 3 : Draw a perpendicular bisector of OP , which cuts it M .

Step 4 : With M as centre and OM as radius, draw a circle which cuts previous circle at A and B .

Step 5 : Join AP and BP . They are the required tangents $AP = BP = 6.3$ cm.

Verification :

$$\begin{aligned}
 AP &= \sqrt{OP^2 - OA^2} \\
 &= \sqrt{(7.2)^2 - (3.6)^2} \\
 &= \sqrt{51.84 - 12.96} \\
 &= \sqrt{38.88} = 6.3 \text{ (approx)}
 \end{aligned}$$

EXERCISE 4.5**Multiple choice questions**

1. If in triangles $\triangle ABC$ and $\triangle EDF$, $\frac{AB}{DE} = \frac{BC}{FD}$ then they will be similar, when
- 1) $\angle B = \angle E$
 - 2) $\angle A = \angle D$
 - 3) $\angle B = \angle D$
 - 4) $\angle A = \angle F$

Hint :**Ans : (3)**

$$\triangle ABC \sim \triangle EDF \text{ if } \frac{AB}{DE} = \frac{BC}{FD} \text{ and}$$

$$\angle B = \angle D$$

$$\angle A = \angle F$$

$$\angle C = \angle E$$

2. In $\triangle LMN$, $L = 60^\circ$, $M = 50^\circ$. If $\triangle LMN \sim \triangle PQR$ then the value of $\angle R$ is


- 1) 40°
- 2) 70°
- 3) 30°
- 4) 110°

Hint :**Ans : (2)**

$$\begin{aligned}
 \angle R &= 180^\circ - (\angle L + \angle M) \\
 &= 180^\circ - (60^\circ + 50^\circ) \\
 &= 180^\circ - 110^\circ \\
 &= 70^\circ
 \end{aligned}$$

3. If $\triangle ABC$ is an isosceles triangle with $\angle C = 90^\circ$ and $AC = 5$ cm, then AB is

- 1) 2.5 cm
- 2) 5 cm
- 3) 10 cm
- 4) $5\sqrt{2}$ cm

Hint :**Ans : (4)**


$$\triangle ABC \text{ is isosceles} \Rightarrow \angle B = \angle A = 45^\circ$$

$$\therefore \sin 45^\circ = \frac{5}{AB}$$

$$\Rightarrow \frac{1}{\sqrt{2}} = \frac{5}{AB}$$

$$\Rightarrow AB = 5\sqrt{2} \text{ cm}$$

4. In a given figure $ST \parallel QR$, $PS = 2$ cm and $SQ = 3$ cm. Then the ratio of the area of ΔPQR to the area of ΔPST is



- 1) 25 : 4 2) 25 : 7 3) 25 : 11 4) 25 : 13

Hint :

Ans : (1)

Where $PQ = PS + SQ$

$$\frac{\text{Area of } \Delta PQR}{\text{Area of } \Delta PST} = \frac{PQ^2}{PS^2}$$

$$= \frac{25}{4}$$

\therefore Ratio = 25 : 4

5. The perimeters of two similar triangles ΔABC and ΔPQR are 36 cm and 24 cm respectively. If $PQ = 10$ cm, then the length of AB is

- 1) $6\frac{2}{3}$ cm 2) $\frac{10\sqrt{6}}{3}$ cm
3) $\frac{2}{3}$ cm 4) 15 cm

Hint :

Ans : (4)

$$\frac{\text{Perimeter of } \Delta ABC}{\text{Perimeter of } \Delta PQR} = \frac{36}{24} = \frac{AB}{PQ}$$

$$\Rightarrow \frac{3}{2} = \frac{AB}{10}$$


$$\Rightarrow AB = 15 \text{ cm}$$

6. If in ΔABC , $DE \parallel BC$, $AB = 3.6$ cm, $AC = 2.4$ cm and $AD = 2.1$ cm then the length of AE is

- 1) 1.4 cm 2) 1.8 cm
3) 1.2 cm 4) 1.05 cm

Hint :

Ans : (1)



By BPT, $\frac{AD}{AB} = \frac{AE}{AC}$

$$\Rightarrow \frac{2.1}{3.6} = \frac{AE}{2.4}$$

$$\Rightarrow AE = 2.4 \times \frac{2.1}{3.6}$$

$$= \frac{2}{3} \times 2.1$$

$$= 2 \times 0.7$$


$$= 1.4 \text{ cm}$$

7. In a ΔABC , AD is the bisector of $\angle BAC$. If $AB = 8$ cm, $BD = 6$ cm and $DC = 3$ cm. The length of the side AC is

- 1) 6 cm 2) 4 cm 3) 3 cm 4) 8 cm

Hint :

Ans : (2)




By ABT, $\frac{8}{x} = \frac{6}{3}$

$$\Rightarrow \frac{8}{x} = 2$$

$$\Rightarrow x = 4 \text{ cm}$$

8. In the adjacent figure $\angle BAC = 90^\circ$ and $AD \perp BC$ then

- 1) $BD \cdot CD = BC^2$ 2) $AB \cdot AC = BC^2$
 3) $BD \cdot CD = AD^2$ 4) $AB \cdot AC = AD^2$



Hint :

Ans : (3)

$$\Delta DBA \sim \Delta DAC$$

$$\therefore \frac{BD}{AD} = \frac{AD}{CD}$$

$$\Rightarrow AD^2 = BD \times CD$$


9. Two poles of heights 6 m and 11 m stand vertically on a plane ground. If the distance

tance between their tops?

- 1) 13 m 2) 14 m 3) 15 m 4) 12.8 m

Hint :

Ans : (1)




$$\therefore AD = \sqrt{12^2 + 5^2} = \sqrt{169} = 13 \text{ m}$$

10. In the given figure, $PR = 26$ cm, $QR = 24$

cm, $\angle PAQ = 90^\circ$, $PA = 6$ cm and $QA = 8$ cm. Find $\angle PQR$

- 1) 80° 2) 85° 3) 75° 4) 90°



Hint :

Ans : (4)

In ΔPAQ , $PA = 6$, $QA = 8$

$$\begin{aligned} \Rightarrow PQ &= \sqrt{6^2 + 8^2} \\ &= \sqrt{100} \\ &= 10 \end{aligned}$$

Also, in ΔPQR , $PQ^2 + QR^2 = 10^2 + 24^2$

$$= 676$$

$$= 26^2$$

$$= PR^2$$


$$\therefore \angle Q = 90^\circ$$

11. A tangent is perpendicular to the radius at the

- 1) centre 2) point of contact
 3) infinity 4) chord

Hint :

Ans : (2)




A tangent is perpendicular to the radius at the point of contact.

12. How many tangents can be drawn to the circle from an exterior point?

- 1) one 2) two 3) infinite 4) zero

Hint :

Ans : (2)




Two tangents can be drawn to the circle from an external point.

13. The two tangents from an external points P to a circle with centre at O are PA and PB. If $\angle APB = 70^\circ$ then the value of $\angle AOB$ is
 1) 100° 2) 110° 3) 120° 4) 130°


Hint :

Ans : (2)



$$\begin{aligned} OA &\perp AP, OB \perp BP \\ \therefore \angle AOB + 90^\circ + 90^\circ + 70^\circ &= 360^\circ \\ \Rightarrow \angle AOB &= 360^\circ - 250^\circ \\ &= 110^\circ \end{aligned}$$

14. In figure CP and CQ are tangents to a circle with centre at O. ARB is another tangent touching the circle at R. If CP = 11 cm and BC = 7 cm, then the length of BR is
 1) 6 cm 2) 5 cm 3) 8 cm 4) 4 cm

**Hint :**

Ans : (4)

$$\begin{aligned} CP &= CQ = 11 \text{ cm}, BC = 7 \\ \therefore BQ &= 11 - 7 \\ &= 4 \\ \therefore BR &= BQ = 4 \text{ cm} \end{aligned}$$

15. In figure if PR is tangent to the circle at P and O is the centre of the circle, then $\angle POQ$ is

1) 120° 2) 100° 3) 110° 4) 90°


Hint :

$$\begin{aligned} \angle RPQ &= 60^\circ \Rightarrow \angle QOP = 60^\circ \text{ (O' is on the circle)} \\ \Rightarrow \angle QOP &= 2(60^\circ) \\ &= 120^\circ \end{aligned}$$

UNIT EXERCISE - 4

1. In the figure, if $BD \perp AC$ and $CE \perp AB$,

$$(i) \triangle AEC \sim \triangle ADB \quad (ii) \frac{CA}{AB} = \frac{CE}{DB}$$

**Solution :**

In $\triangle AEC$, $\triangle ADB$

i) $\angle A$ is common

ii) $\angle AEC = \angle ADB = 90^\circ$

\therefore By AA similarity, $\triangle AEC \sim \triangle ADB$


\therefore Corresponding sides are proportional.

$$(ie) \frac{CA}{AB} = \frac{CE}{DB}$$

Hence proved.

2. In the given figure $AB \parallel CD \parallel EF$. If $AB =$

6 cm, $CD = x$ cm, $EF = 4$ cm, $BD = 5$ cm and $DE = y$ cm. Find x and y .



Solution :

By example 4.9,

$$x = \frac{ab}{a+b} = \frac{6 \times 4}{6+4} = \frac{24}{10} = \frac{12}{5}$$

In $\triangle ABE$,

$$\Rightarrow \frac{ED}{EB} = \frac{CD}{AB}$$

$$\Rightarrow \frac{y}{y+5} = \frac{x}{6}$$

$$\Rightarrow \frac{y}{y+5} = \frac{2}{5 \times 6}$$


$$\Rightarrow \frac{y}{y+5} = \frac{2}{5}$$

$$\Rightarrow 5y = 2y + 10$$


$$\Rightarrow 3y = 10$$

$$\Rightarrow y = \frac{10}{3}$$

3. O is any point inside a triangle ABC. The bisector of $\angle AOB$, $\angle BOC$ and $\angle COA$ meet the sides AB, BC and CA in point D, E and F respectively. Show that $AD \times BE \times CF = DB \times EC \times FA$.



Solution :



By using Ceva's Theorem,


$$\frac{AD}{DB} \times \frac{BE}{EC} \times \frac{AF}{FC} = 1$$

$$\Rightarrow AD \times BE \times AF = DB \times EC \times FC$$

Hence proved.

4. In the figure, ABC is a triangle in which $AB = AC$. Points D and E are points on the side AB and AC respectively such that $AD = AE$. Show that the points B, C, E and D lie on a same circle.

Solution :



$$\begin{aligned} \text{In } \triangle ABC, AB = AC &\Rightarrow \angle ACB = \angle ABC \\ &\Rightarrow \angle ECB = \angle DBC \quad \dots\dots\dots (1) \end{aligned}$$

Also given $AD = AE$

$$\therefore BD = CE$$

$$\therefore DE \parallel BC$$

$$\therefore \angle DBC + \angle BDE = 180^\circ$$

(Sum of 2 adjacent angles is 180°)

$$\Rightarrow \angle ECB + \angle BDE = 180^\circ \quad (\text{From (1)})$$


\therefore Opposite angles are supplementary.

\therefore The points B, C, E & D are concyclic.

Hence proved.

5. Two trains leave a railway station at the same time. The first train travels due west and the second train due north. The first train travels at a speed of 20 km/hr and the second train travels at 30 km/hr. After 2 hours, what is the distance between them?

Solution :



Given speed of 1st train = 20 Km/hr

∴ Speed of 2nd train = 30 Km/hr

$$ON = 30 \times 2 = 60 \text{ Km} = y$$


∴ Distance between 2 trains after 2 hrs.

$$\begin{aligned} Z &= \sqrt{x^2 + y^2} \\ &= \sqrt{40^2 + 60^2} \\ &= \sqrt{1600 + 3600} \\ &= \sqrt{5200} \\ &= \sqrt{40 \times 13} \\ &= 20\sqrt{13} \text{ Km} \end{aligned}$$

6. D is the mid point of side BC and $AE \perp BC$. If $BC = a$, $AC = b$, $AB = c$, $ED = x$, $AD = p$ and $AE = h$, prove that

$$\begin{aligned} (i) \quad b^2 &= p^2 + ax + \frac{a^2}{4} \quad (ii) \quad c^2 = p^2 - ax + \frac{a^2}{4} \\ (iii) \quad b^2 + c^2 &= 2p^2 + \frac{a^2}{2} \end{aligned}$$

Solution :



$$\begin{aligned} i) \quad AC^2 &= b^2 = AE^2 + EC^2 \\ &= h^2 + \left(x + \frac{a}{2}\right)^2 \\ &= (p^2 - x^2) + \left(x^2 + \frac{a^2}{4} + ax\right) \\ &= p^2 + \frac{a^2}{4} + ax \end{aligned}$$

Hence proved

$$\begin{aligned} ii) \quad AB^2 &= c^2 = AE^2 + BE^2 \\ &= h^2 + \left(\frac{a}{2} - x\right)^2 \\ &= p^2 - x^2 + \frac{a^2}{4} + x^2 - ax \\ &= p^2 - ax + \frac{a^2}{4} \end{aligned}$$

Hence proved


iii) Adding (i) & (ii)

$$\begin{aligned} b^2 + c^2 &= \left(p^2 + \frac{a^2}{4} + ax\right) + \left(p^2 - ax + \frac{a^2}{4}\right) \\ &= 2p^2 + 2 \frac{a^2}{4} \\ &= 2p^2 + \frac{a^2}{2} \end{aligned}$$

Hence proved

7. A man whose eye-level is 2 m above the ground wishes to find the height of a tree. He places a mirror horizontally on the ground 20 m from the tree and finds that if he stands at a point C which is 4 m from the mirror B, he can see the reflection of the top of the tree. How height is the tree?

Solution :



Assume that the man & tree are standing up on a straight line

$$\therefore \angle PBC = \angle TBR$$

$$\triangle PCB \sim \triangle TRB$$

\therefore Corresponding sides are proportional

$$(ie) \frac{TR}{PC} = \frac{RB}{BC}$$

$$\Rightarrow \frac{x}{2} = \frac{20}{4}$$


$$\Rightarrow \frac{x}{2} = 5$$

$$\Rightarrow x = 10 \text{ m}$$

\therefore Height of the tree = 10 m

8. An emu which is 8 ft tall is standing at the foot of a pillar which is 30 ft high. It walks away from the pillar. The shadow of the emu falls beyond emu. What is the relation between the length of the shadow and the distance from the emu to the pillar?

Solution :



PR = 30 ft = height of pillar

RU = 8 ft = height of emu

UW = x = Shadow of emu,

y \rightarrow distance between pillar & bird

$$\triangle EUW \sim \triangle PWR$$


$$\therefore \frac{x}{x+y} = \frac{8}{30} \Rightarrow 15x = 4x + 4y$$

$$\Rightarrow x = \frac{4}{11}y$$

\therefore Length of the shadow = $\frac{4}{11}$ (Distance between pillar & emu)

9. Two circles intersect at A and B. From a point P on one of the circles lines PAC and PBD are drawn intersecting the second circle at C and D. Prove that CD is parallel to the tangent at P.

Solution :



PT is a tangent at P.

$\therefore \angle TPB = \angle PAB$ (angles in alternate segment)

$\angle PAB = \angle ACD$ (Exterior angle of a cyclic quad. ABCD is equal to interior opp. angles)


$$\therefore \angle TPB = \angle ACD$$

\therefore Alternate interior angles are equal

\therefore CD and PT are parallel.

10. Let ABC be a triangle and D,E,F are points on the respective sides AB, BC, AC (or their extensions). Let $AD : DB = 5 : 3$, $BE : EC = 3 : 2$ and $AC = 21$. Find the length of the line segment CF.

Solution :




By Ceva's theorem,


$$\begin{aligned} \frac{AD}{DB} \times \frac{BE}{EC} \times \frac{CF}{FA} &= 1 \\ \Rightarrow \frac{5}{3} \times \frac{3}{2} \times \frac{x}{21-x} &= 1 \\ \Rightarrow \frac{x}{21-x} &= \frac{2}{5} \\ \Rightarrow 5x &= 42 - 2x \\ \Rightarrow 7x &= 42 \\ \therefore x &= 6 \\ \therefore CF &= 6 \end{aligned}$$

PROBLEMS FOR PRACTICE

- PQ is a diameter of a circle and PA is a chord. The tangent at A meets PQ produced at B. If $\angle QPA = 35^\circ$, find $\angle QBA$.
(Ans : 20°)
- The medians AD and BE of a $\triangle ABC$ intersect at G. The line through G parallel to BD intersects AC at K. Prove that $AC = 6EK$.
- In the figure, $\angle BED = \angle BDE$ and E is the middle point of BC. Prove that $\frac{AF}{CF} = \frac{AD}{BE}$



- In $\triangle ABC$, $B = 90^\circ$, AD, CE are two medians drawn from A and C respectively. If $AC = 5$ cm, $AD = \frac{3\sqrt{5}}{2}$ find CE.
(Ans : $2\sqrt{5}$ cm)
- In $\triangle ABC$, $\angle C = 90^\circ$, P and Q are the points of the sides CA and CB respectively which divide the sides in the ratio 2 : 1. Prove that $9AQ^2 = 9AC^2 + 4BC^2$.
- A vertical row of trees 12 m long casts a shadow 8 m long on the ground. At the same time, a tower casts a shadow 12 m long on the ground. Find the height of the tower.
(Ans : 60m)
- The area of 2 similar triangles $\triangle ABC$, $\triangle PQR$ are 25 cm^2 , 49 cm^2 respectively. If $QR = 9.8 \text{ cm}$, find BC.
(Ans : 7 cm)
- In $\triangle ABC$, $B = 90^\circ$, D is the mid point of BC. Prove that $AC^2 = 4AD^2 - 3AB^2$.
- Two isosceles triangles have equal vertical angles and their areas are in the ratio of 16 : 25. Find the ratio of their corresponding heights.
(Ans : 4 : 5)
- In the fig, $AD = 3$ cm, $AE = 5$ cm, $BD = 4$ cm, $CE = 4$ cm, $CF = 2$ cm, $BF = 2.5$ cm, Find the pair of parallel lines & hence their lengths.



(Ans : $\frac{28}{9}$ cm ; 7 cm)

11. In an equilateral $\triangle ABC$, E is any point on BC such that $BE = \frac{1}{4} BC$. Prove that $16 AE^2 = 13 AB^2$.
12. In $\triangle PQR$, Qx is the bisector of $\angle Q$ meeting PR in x. If $PQ : QR = 3 : 5$, $XR = 15$ cm, Find PR.
(Ans : 29 cm)
13. $\triangle ABC \sim \triangle DEF$, If $BC = 2$ cm, $EF = 5$ cm and area of $\triangle DEF = 50$, Find area of $\triangle ABC$
(Ans : 8)
14. In $\triangle ABC$, D is a point on AB, $CD = 9$ cm, $DM = 6$ cm, $BC = 12$ cm, $\angle CAB = \angle BCD$. Find the perimeter of $\triangle ADC$.

(Ans 45 cm)

15. A vertical pillar is bent at a height of 2.4 m and its upper end touches the ground at a distance of 1.8 m from its other end on the ground. Find the height of the pillar.

(Ans : 5.4 m)

16. Construct a $\triangle PQR$, such that $QR = 5.1$ cm, $\angle P = 60^\circ$, altitude from P to QR is 3.2 cm.
17. Construct a $\triangle ABC$ having base 6 cm, vertical angle 60° and median in rough the vertex is 4 cm.
18. Construct a \triangle of base 5 cm and vertical angle 120° such that the bisector of the vertical angle meets the base at a point 3 cm.
19. Draw the tangents to a circle whose diameter is 10 cm from a point 13 cm from its centre.
20. Draw a circle of diameter 8 cm. Take a
the circle, draw a tangent to the circle at the point P.

CHAPTER 5

COORDINATE GEOMETRY

I. AREA OF A TRIANGLE AND QUADRILATERAL :

Key Points

- ✓ The area of ΔABC is the absolute value of the expression

$$= \frac{1}{2} \{ (x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)) \}$$

- ✓ The vertices $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ of ΔABC are said to be “taken in order” if A, B, C are taken in counter-clock wise direction.
- ✓ The following pictorial representation helps us to write the above formula very easily.

$$\frac{1}{2} \left\{ \begin{array}{c} x_1 \quad x_2 \quad x_3 \quad x_1 \\ \swarrow \quad \searrow \quad \swarrow \quad \searrow \\ y_1 \quad y_2 \quad y_3 \quad y_1 \end{array} \right\}$$

$$\text{Area of } \Delta ABC = \frac{1}{2} \{ (x_1y_2 + x_2y_3 + x_3y_1) - (x_2y_1 + x_3y_2 + x_1y_3) \} \text{ sq. units.}$$

- ✓ Three distinct points $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ will be collinear if and only if area of $\Delta ABC = 0$.
- ✓ Area of the quadrilateral $ABCD = \frac{1}{2} \{ (x_1 - x_3)(y_2 - y_4) - (x_2 - x_4)(y_1 - y_3) \}$ sq. units.

Example 5.1

Find the area of the triangle whose vertices are $(-3, 5)$, $(5, 6)$ and $(5, -2)$.

Solution :


Plot the points in a rough diagram and take them in counter-clockwise order.

Let the vertices be $A(-3, 5)$, $B(5, -2)$, $C(5, 6)$

$$\begin{array}{ccc} \downarrow & \downarrow & \downarrow \\ (x_1, y_1) & (x_2, y_2) & (x_3, y_3) \end{array}$$

The area ΔABC is

$$\begin{aligned} &= \frac{1}{2} \{ (x_1y_2 + x_2y_3 + x_3y_1) - (x_2y_1 + x_3y_2 + x_1y_3) \} \\ &= \frac{1}{2} \{ (6 + 30 + 25) - (25 - 10 - 18) \} \\ &= \frac{1}{2} \{ 61 + 3 \} \\ &= \frac{1}{2} (64) = 32 \text{ sq. units} \end{aligned}$$

**Example 5.2**

Show that the points P(-1.5, 3), Q(6, -2), R(-3, 4) are collinear.

Solution :

The points are P(-1.5, 3), Q(6, -2), R(-3, 4)

Area of ΔPQR

$$\begin{aligned}
 &= \frac{1}{2} \{ (x_1y_2 + x_2y_3 + x_3y_1) - (x_2y_1 + x_3y_2 + x_1y_3) \} \\
 &= \frac{1}{2} \{ (3 + 24 - 9) - (18 + 6 - 6) \} \\
 &= \frac{1}{2} \{ 18 - 18 \} = 0
 \end{aligned}$$

Therefore, the given points are collinear.

Example 5.3

If the area of the triangle formed by the vertices A(-1, 2), B(k, -2) and C(7, 4) (taken in order) is 22 sq. units, find the value of k.

Solution :

The vertices are A(-1, 2), B(k, -2) and C(7, 4)

Area of triangle ABC is 22 sq. units

$$\begin{aligned}
 \frac{1}{2} \{ (x_1y_2 + x_2y_3 + x_3y_1) - (x_2y_1 + x_3y_2 + x_1y_3) \} &= 22 \\
 \frac{1}{2} \{ (2 + 4k + 14) - (2k - 14 - 4) \} &= 22 \\
 2k + 34 &= 44 \\
 \text{gives } 2k &= 10 \text{ so } k = 5
 \end{aligned}$$

Example 5.4

If the points P(-1, -4), Q(b, c) and R(5, -1) are collinear and if $2b + c = 4$, then find the values of b and c.

Solution :

Since the three points P(-1, -4), Q(b, c) and R(5, -1) are collinear,

Area of triangle PQR = 0

$$\begin{aligned}
 \frac{1}{2} \{ (x_1y_2 + x_2y_3 + x_3y_1) - (x_2y_1 + x_3y_2 + x_1y_3) \} &= 0 \\
 \frac{1}{2} \{ (-c - b - 20) - (-4b + 5c + 1) \} &= 0 \\
 -c - b - 20 + 4b - 5c - 1 &= 0 \\
 b - 2c &= 7 \quad \text{--- (1)}
 \end{aligned}$$

$$\begin{aligned}
 \text{Also, } 2b + c &= 4 \quad \text{--- (2)} \\
 \text{(from given information)}
 \end{aligned}$$


Solving (1) and (2) we get $b = 3$, $c = -2$

Example 5.5

The floor of a hall is covered with identical tiles which are in the shapes of triangles. One such triangle has the vertices at (-3, 2), (-1, -1) and (1, 2). If the floor of the hall is completely covered by 110 tiles, find the area of the floor.

Solution :

Vertices of one triangular tile are at (-3, 2), (-1, -1) and (1, 2)



Area of this tile

$$= \frac{1}{2} \{ (3 - 2 + 2) - (-2 - 1 - 6) \} \text{ sq. units}$$

$$= \frac{1}{2} (12) = 6 \text{ sq. units}$$

Since the floor is covered by 110 triangle shaped identical tiles,

$$\text{Area of floor} = 110 \times 6 = 660 \text{ sq. units}$$


Find the area of the quadrilateral formed by the points (8, 6), (5, 11), (-5, 12) and (-4, 3).

Solution :

Before determining the area of quadrilateral, plot the vertices in a graph.

Let the vertices be A(8,6), B(5,11), C(-5, 12) and D(-4, 3)

Therefore, area of the quadrilateral ABCD



$$= \frac{1}{2} \left\{ (x_1y_2 + x_2y_3 + x_3y_4 + x_4y_1) - (x_2y_1 + x_3y_2 + x_4y_3 + x_1y_4) \right\}$$

$$= \frac{1}{2} \{ (88 + 60 - 15 - 24) - (30 - 55 - 48 + 24) \}$$

$$= \frac{1}{2} \{ 109 + 49 \}$$

$$= \frac{1}{2} \{ 158 \} = 79 \text{ sq. units}$$

Example 5.7

The given diagram shows a plan for constructing a new parking lot at a campus. It is estimated that such construction would cost ₹1300 per square feet. What will be the total cost for making the parking lot?

Solution :

vertices are at A(2,2), B(5,5), C(4,9) and D(1,7)

Therefore, Area of parking lot

$$= \frac{1}{2} \{ (10 + 45 + 28 + 2) - (10 + 20 + 9 + 14) \}$$

$$= \frac{1}{2} \{ 85 - 53 \}$$

$$\text{Use : } \frac{1}{2} \left\{ \begin{array}{cccc} 2 & 5 & 4 & 1 \\ 2 & 5 & 9 & 7 \\ & & & 2 \end{array} \right\}$$

$$= \frac{1}{2} (32) = 16 \text{ sq. units.}$$

So, area of parking lot = 16 sq. feet

Construction rate per square feet = ₹1300

Therefore, total cost for constructing the parking lot = $16 \times 1300 = ₹20800$

EXERCISE 5.1

1. Find the area of the triangle formed by the points


(i) (1, -1), (-4, 6) and (-3, -5)

(ii) (-10, -4), (-8, -1) and (-3, -5)

Solution:

- i) Given points are (1, -1), (-4, 6), (-3, -5)

Taking A, B, C in anti clockwise direction.



∴ Area of triangle ABC

$$= \frac{1}{2} \left\{ \begin{array}{ccc} 1 & -4 & -3 \\ -1 & 6 & -5 \end{array} \right\}$$

$$= \frac{1}{2} \{ (6 + 20 + 3) - (4 - 18 - 5) \}$$


$$= \frac{1}{2} [29 + 19]$$

$$= \frac{1}{2} (48)$$

$$= 24 \text{ sq. units}$$

- (ii) Given points are (-10, -4), (-8, -1), (-3, -3)

Taking in anticlock direction



Let A (-10, -4), B (-3, -5), C (-8, -1)

∴ Area of triangle ABC

$$= \frac{1}{2} \left\{ \begin{array}{ccc} -10 & -3 & -8 \\ -4 & -5 & -1 \end{array} \right\}$$

$$= \frac{1}{2} [(50 + 3 + 32) - (12 + 40 + 10)]$$

$$= \frac{1}{2} [85 - 62]$$

$$= \frac{23}{2}$$

$$= 11.5 \text{ sq. units}$$

2. Determine whether the sets of points are collinear ?

(i) $\left\{ -\frac{1}{2}, 3 \right\}$, (-5, 6) and (-8, 8)

(ii) (a, b + c), (b, c + a) and (c, a + b)

Solution :

Area of triangle formed by 3 points

$$= \frac{1}{2} \left[\begin{array}{ccc} -1/2 & -5 & -8 \\ 3 & 6 & 8 \end{array} \right]$$

$$= \frac{1}{2} [(-3 - 40 - 24) - (-15 - 48 - 4)]$$

$$= \frac{1}{2} [-67 - (-67)]$$

$$= \frac{1}{2} (0) = 0$$

∴ The 3 points are collinear.

- ii) Given points are A (a, b + c), B (b, c + a), C (c, a + b)

Area of triangle formed by 3 points

$$= \frac{1}{2} \left[\begin{array}{ccc} a & b & c \\ b+c & c+a & a+b \end{array} \right]$$

$$\begin{aligned}
 &= \frac{1}{2}[(ac + a^2 + ab + b^2 + bc + c^2) \\
 &\quad - (b^2 + bc + c^2 + ac + a^2 + ab)] \\
 &= \frac{1}{2}[(a^2 + b^2 + c^2 + ab + bc + ca) \\
 &\quad - (a^2 + b^2 + c^2 + ab + bc + ca)] \\
 &= \frac{1}{2}[0] \\
 &\therefore \text{The 3 points are collinear.}
 \end{aligned}$$

3. Vertices of given triangles are taken in order and their areas are provided below. In each case, find the value of 'p'.

S.No.	Vertices	Area (sq. units)
(i)	(0, 0), (p, 8), (6, 2)	20
(ii)	(p, p), (5, 6), (5, -2)	32

Area of triangle formed by 3 points

$$\begin{aligned}
 &= \frac{1}{2} \left[\begin{array}{ccc} 0 & p & 6 \\ 0 & 8 & 2 \\ 0 & 0 & 0 \end{array} \right] \\
 &\Rightarrow [0 + 2p + 0] - [0 + 48 + 0] = 40 \\
 &\Rightarrow 2p - 48 = 40 \\
 &\Rightarrow 2p = 88 \\
 &\therefore p = 44
 \end{aligned}$$

ii) Given vertices are (p, p), (5, 6), (5, -2),
Area = 32 sq. units.

Area of triangle formed by 3 points

$$\begin{aligned}
 &= \frac{1}{2} \left[\begin{array}{ccc} p & 5 & 5 \\ p & 6 & -2 \\ p & p & p \end{array} \right] = 32 \\
 &\Rightarrow (6p - 10 + 5p) - (5p + 30 - 2p) = 64 \\
 &\Rightarrow (11p - 10) - (3p + 30) = 64 \\
 &\Rightarrow 8p = 104 \\
 &\quad p = \frac{104}{8} \\
 &\quad p = 13
 \end{aligned}$$

4. In each of the following, find the value of 'a' for which the given points are collinear.

(i) (2, 3), (4, a) and (6, -3) (ii) (a, 2 - 2a), (-a+1, 2a) and (-4 - a, 6 - 2a)

Solution :

i) Given, 3 points (2, 3), (4, a), (6, -3) are collinear.

$$\text{ie, } \frac{1}{2} \begin{bmatrix} 2 & 4 & 6 & 2 \\ 3 & a & -3 & 3 \end{bmatrix} = 0$$

$$\Rightarrow (2a - 12 + 18) - (12 + 6a - 6) = 0$$

$$\Rightarrow (2a + 6) - (6a + 6) = 0$$

$$\Rightarrow -4a = 0$$

$$\Rightarrow a = 0$$

ii) Given 2 points (a, 2 - 2a), (-a + 1, 2a),
(-4 - a, 6 - 2a) are collinear.

\therefore Area of triangle formed by 3 points is 0.

$$\text{ie, } \frac{1}{2} \left[\begin{array}{ccc} a & -a+1 & -4-a \\ 2-2a & 2a & 6-2a \\ a & 2-2a & 2-2a \end{array} \right] = 0$$

$$\begin{aligned}
 &\Rightarrow [2a^2 + (-a+1)(6-2a) + (-4-a)(2-2a)] \\
 &\quad - [(2-2a)(-a+1) + 2a(-4-a) + a(6-2a)] = 0
 \end{aligned}$$

$$\Rightarrow (2a^2 + 2a^2 - 8a + 6 + 2a^2 + 6a - 8)$$

$$- [2a^2 - 4a + 2 - 2a^2 - 8a + 6a - 2a^2] = 0$$

$$\Rightarrow (6a^2 - 2a - 2) - (-2a^2 - 6a + 2) = 0$$

$$\Rightarrow 8a^2 + 4a - 4 = 0$$

$$\Rightarrow 2a^2 + a - 1 = 0$$

$$\Rightarrow a = -1, \frac{1}{2}$$

5. Find the area of the quadrilateral whose vertices are at

- (i) $(-9, -2)$, $(-8, -4)$, $(2, 2)$ and $(1, -3)$ (ii) $(-9, 0)$, $(-8, 6)$, $(-1, -2)$ and $(-6, -3)$


Solution :

- (i) Given vertices of quadrilateral are $(-9, -2)$, $(-8, -4)$, $(2, 2)$, $(1, -3)$

First we plot the points in the plane

Let A $(-9, -2)$, B $(-8, -4)$, C $(1, -3)$, D $(2, 2)$

- 2) \therefore Area of quadrilateral



$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -9 & -8 & 1 & 2 & -9 \\ -2 & -4 & -3 & 2 & -2 \end{vmatrix} \\
 &= \frac{1}{2} [(36 + 24 + 2 - 4) - (16 - 4 - 6 - 18)] \\
 &= \frac{1}{2} [58 - (-12)] \\
 &= \frac{1}{2} [70] \\
 &= 35 \text{ sq. units}
 \end{aligned}$$


- ii) Given vertices of quadrilateral are $(-9, 0)$, $(-8, 6)$, $(-1, -2)$, $(-6, -3)$

First, we plot the points in the plane.

Let A $(-8, 6)$, B $(-9, 0)$, C $(-6, -3)$, D $(-1, -2)$

Area of quadrilateral

$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -8 & -9 & -6 & -1 & -8 \\ 6 & 0 & -3 & -2 & 6 \end{vmatrix} \\
 &= \frac{1}{2} [(0 + 27 + 12 - 6) - (-54 + 0 + 3 + 16)] \\
 &= \frac{1}{2} [33 - (-35)] \\
 &= \frac{1}{2} [68] \\
 &= 34 \text{ sq. units}
 \end{aligned}$$



6. Find the value of k, if the area of a quadrilateral is 28 sq. units, whose vertices are $(-4, -2)$, $(-3, k)$, $(3, -2)$ and $(2, 3)$

Solution :

$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -4 & -3 & 3 & 2 & -4 \\ -2 & k & -2 & 3 & -2 \end{vmatrix} = 28 \\
 \Rightarrow & (-4k + 6 + 9 - 4) - (6 + 3k - 4 - 12) = 56 \\
 \Rightarrow & (11 - 4k) - (3k - 10) = 56 \\
 \Rightarrow & 21 - 7k = 56 \\
 & \therefore 7k = -35 \\
 & k = -5
 \end{aligned}$$

7. If the points A $(-3, 9)$, B (a, b) and C $(4, -5)$ are collinear and if $a + b = 1$, then find a and b.

Solution :

Given, A $(-3, 9)$, B (a, b) , C $(4, -5)$ are collinear and $a + b = 1$. — (1)

Area of triangle formed by 3 points = 0.

$$\text{ie) } \frac{1}{2} \begin{vmatrix} -3 & a & 4 & -3 \\ 9 & b & -5 & 9 \end{vmatrix} = 0$$

$$\Rightarrow (-3b - 5a + 36) - (9a + 4b + 15) = 0$$

$$\Rightarrow -5a - 3b + 36 - 9a - 4b - 15 = 0$$

$$\Rightarrow -14a - 7b + 21 = 0$$

$$\Rightarrow 2a + b - 3 = 0$$

$$\Rightarrow 2a + 1 - a - 3 = 0 \quad (\text{from (1)})$$

$$\therefore \Rightarrow a = 2 \quad b = -1$$


8. Let P(11,7), Q(13.5, 4) and R(9.5, 4) be the mid-points of the sides AB, BC and AC respectively of $\triangle ABC$. Find the coordinates of the vertices A, B and C. Hence find the area of $\triangle ABC$ and compare this with area of $\triangle PQR$.

Solution :

In $\triangle ABC$, given that P, Q, R are the mid points of AB, BC, CA respectively.

$$P(11, 7), Q(13.5, 4), R(9.5, 4)$$

Let A (x_1, y_1) , B (x_2, y_2) , C (x_3, y_3) be the vertices.



$$\therefore \frac{x_1 + x_2}{2} = 11, \quad \frac{y_1 + y_2}{2} = 7$$

$$\Rightarrow x_1 + x_2 = 22, \quad y_1 + y_2 = 14 \quad \dots(1)$$

$$\frac{x_2 + x_3}{2} = \frac{27}{2}, \quad \frac{y_2 + y_3}{2} = 4$$

$$\Rightarrow x_2 + x_3 = 27, \quad y_2 + y_3 = 8 \quad \dots(2)$$

$$\frac{x_1 + x_3}{2} = \frac{19}{2}, \quad \frac{y_1 + y_3}{2} = 4$$

$$\Rightarrow x_1 + x_3 = 19, \quad y_1 + y_3 = 8 \quad \dots(3)$$

\therefore Adding (1), (2) and (3)

$$2(x_1 + x_2 + x_3) = 68, \quad 2(y_1 + y_2 + y_3) = 30$$

$$x_1 + x_2 + x_3 = 34, \quad y_1 + y_2 + y_3 = 15$$

$$22 + x_3 = 34, \quad 14 + y_3 = 15$$

$$x_3 = 12, \quad y_3 = 1$$

$$\therefore C(12, 1)$$

$$x_2 = 15, \quad y_2 = 7$$

$$\therefore B \text{ is } (15, 7)$$

$$(1) \Rightarrow x_1 + 15 = 22, \quad y_1 + 7 = 14$$

$$x_1 = 7, \quad y_1 = 7$$

$$\therefore A \text{ is } (7, 7)$$

Area of $\triangle ABC$

$$= \frac{1}{2} \begin{vmatrix} 7 & 15 & 12 & 7 \\ 7 & 7 & 1 & 7 \end{vmatrix}$$

$$= \frac{1}{2} [(49 + 15 + 84) - (105 + 84 + 7)]$$

$$= \frac{1}{2} [148 - 196]$$


$$= \frac{1}{2} (-48)$$

$$= 24 \quad (\because \text{Area can't be -ve})$$

Area of ΔPQR

$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} 11 & \frac{27}{2} & \frac{19}{2} & 11 \\ 7 & 4 & 4 & 7 \end{vmatrix} \\
 &= \frac{1}{2} [(44 + 54 + 66.5) - (94.5 + 38 + 44)] \\
 &= \frac{1}{2} [164.5 - 176.5] \\
 &= \frac{1}{2} [-12] \\
 &= 6 \quad (\because \text{Area can't be } -ve) \\
 &\therefore \text{Area of } \Delta ABC = 4 \text{ (Area of } \Delta PQR).
 \end{aligned}$$

9. In the figure, the quadrilateral swimming pool shown is surrounded by concrete patio. Find the area of the patio.

**Solution :**

Required area of the patio = Area of portion ABCD – Area of portion EFGH


$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -4 & 8 & 6 & -10 & -4 \\ -8 & -4 & 10 & 6 & -8 \end{vmatrix} - \frac{1}{2} \begin{vmatrix} -3 & 6 & 3 & -6 & -3 \\ -5 & -2 & 7 & 4 & -5 \end{vmatrix} \\
 &= \frac{1}{2} [(16 + 80 + 36 + 80) - (-64 - 24 - 100 - 24)] \\
 &\quad - \frac{1}{2} [(6 + 42 + 12 + 30) - (-30 - 6 - 42 - 12)] \\
 &= \frac{1}{2} [212 - (-212)] - \frac{1}{2} [90 - (-90)] \\
 &= \frac{1}{2} [424] - \frac{1}{2} [180]
 \end{aligned}$$

10. A triangular shaped glass with vertices at A(-5,-4), B(1,6) and C(7,-4) has to be painted. If one bucket of paint covers 6 square feet, how many buckets of paint will be required to paint the whole glass, if only one coat of paint is applied.

Solution :

$$\begin{aligned}
 \text{Area of } \Delta ABC &= \frac{1}{2} \begin{vmatrix} -5 & 1 & 7 & -5 \\ -4 & 6 & -4 & -4 \end{vmatrix} \\
 &= \frac{1}{2} [(-30 - 4 - 28) - (-4 + 42 + 20)] \\
 &= \frac{1}{2} [-62 - (58)] \\
 &= \frac{1}{2} [-120] \\
 &= 60 \text{ sq. units (Area can't be } -ve). \\
 \therefore \text{No. of paint cans needed} &= \frac{60}{6} = 10
 \end{aligned}$$

11. In the figure, find the area of (i) triangle AGF (ii) triangle FED (iii) quadrilateral BCEG.

**Solution :**

- i) Area of ΔAGF

$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -5 & -9/2 & -2 & -5 \\ 3 & 1/2 & 3 & 3 \end{vmatrix} \\
 &= \frac{1}{2} [(-2.5 - 13.5 - 6) - (-13.5 - 1 - 15)] \\
 &= \frac{1}{2} [(-22) - (-29.5)] \\
 &= \frac{1}{2} [7.5] \\
 &= 3.75 \text{ sq. units}
 \end{aligned}$$

ii) Area of ΔFED

$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -2 & 3/2 & 1 & -2 \\ 3 & 1 & 3 & 3 \end{vmatrix} \\
 &= \frac{1}{2} [(-2 + 4.5 + 3) - (-4.5 + 1 - 6)] \\
 &= \frac{1}{2} [5.5 - (-0.5)] \\
 &= \frac{1}{2} [6] \\
 &= 3 \text{ sq. units}
 \end{aligned}$$

iii) Area of quadrilateral BCEG

$$\begin{aligned}
 &= \frac{1}{2} \begin{vmatrix} -4 & 2 & 3/2 & -9/2 & -4 \\ -2 & -1 & 1 & 1/2 & -2 \end{vmatrix} \\
 &= \frac{1}{2} [(4 + 2 + 0.75 + 9) - (-4 - 1.5 - 4.5 - 2)] \\
 &= \frac{1}{2} [15.75 + 12] \\
 &= \frac{27.75}{2} \\
 &= 13.875 \\
 &\approx 13.88 \text{ sq. units}
 \end{aligned}$$

II. INCLINATION AND SLOPE OF STRAIGHT LINE

Key Points

- ✓ The inclination of a line or the angle of inclination of a line is the angle which a straight line part of the line above the X axis.
- ✓ The inclination of X axis and every line parallel to X axis is 0° .
- ✓ The inclination of Y axis and every line parallel to Y axis is 90° .
- ✓ If θ is the angle of inclination of a non-vertical straight line, then $\tan \theta$ is called the slope or gradient of the line and is denoted by m .
- ✓ The slope of the straight line is $m = \tan \theta$, $0 \leq 180^\circ$, $\theta \neq 90^\circ$.
- ✓ The slope of the line through (x_1, y_1) and (x_2, y_2) with $x_1 \neq x_2$ is $\frac{y_2 - y_1}{x_2 - x_1}$.
- ✓ If $\theta = 0^\circ$, the line is parallel to the positive direction of X axis.
- ✓ If $0 < \theta < 90^\circ$, the line has positive slope.
- ✓ If $90^\circ < \theta < 180^\circ$, the line has negative slope.
- ✓ If $\theta = 180^\circ$, the line is parallel to the negative direction of X axis.
- ✓ If $\theta = 90^\circ$, the slope is undefined.
- ✓ Non vertical lines are parallel if and only if their slopes are equal.
- ✓ Two non-vertical lines with slopes m_1 and m_2 are perpendicular if and only if $m_1 m_2 = -1$.

Example 5.8

(i) What is the slope of a line whose inclination is 30° ?

(ii) What is the inclination of a line whose slope is $\sqrt{3}$?

Solution :

(i) Here $\theta = 30^\circ$

Slope $m = \tan \theta$

Therefore, slope $m = \tan 30^\circ = \frac{1}{\sqrt{3}}$

(ii) Given $m = \sqrt{3}$, let θ be the inclination of the line

$$\tan \theta = \sqrt{3}$$

We get, $\theta = 60^\circ$

Example 5.9

Find the slope of a line joining the given points

(i) $(-6, 1)$ and $(-3, 2)$ (ii) $\left(-\frac{1}{3}, \frac{1}{2}\right)$
and $\left(\frac{2}{7}, \frac{3}{7}\right)$ (iii) $(14, 10)$ and $(14, -6)$

Solution :

(i) $(-6, 1)$ and $(-3, 2)$

$$\text{The slope} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{2 - 1}{-3 - (-6)} = \frac{1}{3}.$$

(ii) $\left(-\frac{1}{3}, \frac{1}{2}\right)$ and $\left(\frac{2}{7}, \frac{3}{7}\right)$

$$\begin{aligned} \text{The slope} &= \frac{\frac{3}{7} - \frac{1}{2}}{\frac{2}{7} - \left(-\frac{1}{3}\right)} = \frac{\frac{6-7}{14}}{\frac{6+7}{21}} \\ &= -\frac{1}{14} \times \frac{21}{13} = -\frac{3}{26}. \end{aligned}$$

(iii) $(14, 10)$ and $(14, -6)$

$$\text{The slope} = \frac{-6 - 10}{14 - 14} = \frac{-16}{0}.$$

The slope is undefined.

Example 5.10

The line r passes through the points $(-2, 2)$ and $(5, 8)$ and the line s passes through the points $(-8, 7)$ and $(-2, 0)$. Is the line r perpendicular to s ?

Solution :

$$\text{The slope of line } r \text{ is } m_1 = \frac{8 - 2}{5 - (-2)} = \frac{6}{7}$$

$$\text{The slope of line } s \text{ is } m_2 = \frac{0 - 7}{-2 - (-8)} = \frac{-7}{6}$$

$$\text{The product of slopes} = \frac{6}{7} \times \frac{-7}{6} = -1$$

$$\text{That is, } m_1 m_2 = -1$$

Therefore, the line r is perpendicular to line s .

Example 5.11

The line p passes through the points $(3, -2)$, $(12, 2)$ and $(12, 2)$. Is p parallel to q ?

Solution :

$$\text{The slope of line } p \text{ is } m_1 = \frac{4 - (-2)}{12 - 3} = \frac{6}{9} = \frac{2}{3}$$

$$\text{The slope of line } q \text{ is } m_2 = \frac{2 + 2}{12 - 6} = \frac{4}{6} = \frac{2}{3}$$

Thus, slope of line p = slope of line q .

Therefore, line p is parallel to the line q .

Example 5.12


Show that the points $(-2, 5)$, $(6, -1)$ and $(2, 2)$ are collinear.

Solution :

The vertices are A $(-2, 5)$, B $(6, -1)$ and C $(2, 2)$.

$$\text{Slope of AB} = \frac{-1 - 5}{6 - (-2)} = \frac{-6}{8} = \frac{-3}{4}$$

$$\text{Slope of BC} = \frac{2 + 1}{2 - 6} = \frac{3}{-4} = \frac{-3}{4}$$



We get, Slope of AB = Slope of BC.

Therefore, the points A, B, C all lie in a same straight line.

Hence the points A, B and C are collinear.

Example 5.13

Let A (1, -2), B (6, -2), C(5, 1) and D (2, 1) be four points.

(a) AB (b) CD

(ii) Find the slope of the line segments

(a) BC (b) AD

(iii) What can you deduce from your answer?

Solution :

$$(i) \quad (a) \text{ Slope of AB} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-2 + 2}{6 - 1} = 0$$

$$(b) \text{ Slope of CD} = \frac{1 - 1}{2 - 5} = \frac{0}{-3} = 0$$

$$(ii) \quad (a) \text{ Slope of BC} = \frac{1 + 2}{5 - 6} = \frac{3}{-1} = -3$$

$$(b) \text{ Slope of AD} = \frac{1 + 2}{2 - 1} = \frac{3}{1} = 3$$

(iii) The slope of AB and CD are equal so AB, CD are parallel.

Similarly the lines AD and BC are not parallel, since their slopes are not equal.


So, we can deduce that the quadrilateral ABCD is a trapezium.

Example 5.14

Consider the graph representing growth of population (in crores). Find the slope of the line AB and hence estimate the population in the year 2030?

Solution :

The points A(2005, 96) and B(2015, 100) are on the line AB.



$$\text{Slope of AB} = \frac{100 - 96}{2015 - 2005} = \frac{4}{10} = \frac{2}{5}$$

Let the growth of population in 2030 be k crores.

Assuming that the point C(2030, k) is on AB,

we have, slope of AC = slope of AB

$$\frac{k - 96}{2030 - 2005} = \frac{2}{5} \text{ gives } \frac{k - 96}{25} = \frac{2}{5}$$

$$k - 96 = 10$$

$$k = 106$$

Hence the estimated population in 2030 = 106 Crores.

Example 5.15

Without using Pythagoras theorem, show that the points (1, -4), (2, -3) and (4, -7) form a right angled triangle.

Solution :

Let the given points be A(1, -4), B(2, -3) and C(4, -7).

$$\text{The slope of AB} = \frac{-3 + 4}{2 - 1} = \frac{1}{1} = 1$$

$$\text{The slope of BC} = \frac{-7 + 3}{4 - 2} = \frac{-4}{2} = -2$$

$$\text{The slope of AC} = \frac{-7 + 4}{4 - 1} = \frac{-3}{3} = -1$$

$$\text{Slope of AB} \times \text{slope of AC} = (1)(-1) = -1$$

AB is perpendicular to AC. $\angle A = 90^\circ$

Therefore, $\triangle ABC$ is a right angled triangle.


Example 5.16

Prove analytically that the line segment joining the mid-points of two sides of a triangle is parallel to the third side and is equal to half of its length.

Solution :

Let P(a, b), Q(c, d) and R(e, f) be the vertices of a triangle.

Let S be the mid-point of PQ and T be the mid-point of PR.



$$\text{Therefore } S = \left(\frac{a+c}{2}, \frac{b+d}{2} \right)$$

$$\text{and } T = \left(\frac{a+e}{2}, \frac{b+f}{2} \right)$$

$$\text{Now, slope of ST} = \frac{\frac{b+f}{2} - \frac{b+d}{2}}{\frac{a+e}{2} - \frac{a+c}{2}} = \frac{f-d}{e-c}$$

$$\text{And slope of QR} = \frac{f-d}{e-c}$$

Therefore, ST is parallel to QR. (since, their slopes are equal)

Also

$$\begin{aligned} ST &= \sqrt{\left(\frac{a+e}{2} - \frac{a+c}{2} \right)^2 + \left(\frac{b+f}{2} - \frac{b+d}{2} \right)^2} \\ &= \frac{1}{2} \sqrt{(e-c)^2 + (f-d)^2} \end{aligned}$$

$$ST = \frac{1}{2} QR$$

Thus ST is parallel to QR and half of it.

EXERCISE 5.2

- What is the slope of a line whose inclination with positive direction of x-axis is
(i) 90° (ii) 0°

Solution :

$$\text{i) } \theta = 90^\circ \quad m = \tan 90^\circ = \text{undefined.}$$

$$\text{ii) } \theta = 0^\circ \quad m = \tan 0^\circ = 0$$

- What is the inclination of a line whose slope is (i) 0 (ii) 1

Solution :

$$\text{i) } m = 0 \Rightarrow \tan \theta = 0 \quad \therefore \theta = 0^\circ$$

$$\text{ii) } m = 1 \Rightarrow \tan \theta = 1 \quad \therefore \theta = 45^\circ$$

3. Find the slope of a line joining the points**(i) $(5, \sqrt{5})$ with the origin****(ii) $(\sin \theta, -\cos \theta)$ and $(-\sin \theta, \cos \theta)$** **Solution :****i) Slope of the line joining $(5, \sqrt{5})$, $(0, 0)$**

$$\therefore m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - \sqrt{5}}{0 - 5} = \frac{1}{\sqrt{5}}$$

$$\therefore \text{Slope} = \frac{1}{\sqrt{5}}$$

**ii) Slope of line joining $(\sin \theta, -\cos \theta)$
 $(-\sin \theta, \cos \theta)$**

$$\begin{aligned} m &= \frac{y_2 - y_1}{x_2 - x_1} = \frac{\cos \theta + \cos \theta}{-\sin \theta - \sin \theta} \\ &= \frac{2 \cos \theta}{-2 \sin \theta} \\ &= -\cot \theta \end{aligned}$$

4. What is the slope of a line perpendicular to the line joining A $(5, 1)$ and P where P is the mid-point of the segment joining $(4, 2)$ and $(-6, 4)$.**Solution :**Given P is the midpoint of $(4, 2)$, $(-6, 4)$

$$\begin{aligned} \Rightarrow P &= \left(\frac{4 - 6}{2}, \frac{2 + 4}{2} \right) \\ &= (-1, 3) \end{aligned}$$

 \therefore Slope of the line joining A $(5, 1)$, P $(-1, 3)$

$$\begin{aligned} m &= \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 1}{-1 - 5} \\ &= \frac{2}{-6} \\ &= \frac{-1}{3} \end{aligned}$$

\therefore Slope of the line perpendicular to the line joining A and P is $\frac{-1}{m} = 3$

5. Show that the given points are collinear: $(-3, -4)$, $(7, 2)$ and $(12, 5)$.**Solution :**Given points are A $(-3, -4)$, B $(7, 2)$, C $(12, 5)$

$$\begin{aligned} \text{Slope of AB} &= \frac{2 + 4}{7 + 3} \\ &= \frac{6}{10} \\ &= \frac{3}{5} \end{aligned}$$

$$\begin{aligned} \text{Slope of BC} &= \frac{5 - 2}{12 - 7} \\ &= \frac{3}{5} \end{aligned}$$

 \therefore Slope of AB = Slope of BC

But B is the common point.

 \therefore A, B, C are collinear.**6. If the three points $(3, -1)$, $(a, 3)$ and $(1, -3)$ are collinear, find the value of a .****Solution :**Given points A $(3, -1)$, B $(a, 3)$, C $(1, -3)$ are collinear. **\therefore Slope of AB = Slope of BC**

$$\begin{aligned} \Rightarrow \frac{4}{a - 3} &= \frac{-6}{1 - a} \\ \Rightarrow 4 - 4a &= -6a + 18 \\ \Rightarrow 2a &= 14 \\ a &= 7 \end{aligned}$$

7. The line through the points $(-2, a)$ and $(9, 3)$ has slope $-\frac{1}{2}$. Find the value of a .

Solution :

Slope of the line joining $(-2, a), (9, 3) = -\frac{1}{2}$

$$\Rightarrow \frac{3-a}{9+2} = \frac{-1}{2}$$

$$\Rightarrow \frac{3-a}{11} = \frac{-1}{2}$$

$$\Rightarrow 6 - 2a = -11$$

$$\Rightarrow 2a = 17$$

$$\therefore a = \frac{17}{2}$$

8. The line through the points $(-2, 6)$ and $(4, 8)$ is perpendicular to the line through the points $(8, 12)$ and $(x, 24)$. Find the value of x .

Solution :

Slope of line joining $(-2, 6), (4, 8)$

$$m_1 = \frac{8-6}{4+2} = \frac{2}{6} = \frac{1}{3}$$

Slope of line joining $(8, 12), (x, 24)$

$$m_2 = \frac{24-12}{x-8} = \frac{12}{x-8}$$

Since two lines are perpendicular,

$$\Rightarrow \frac{1}{3} \times \frac{12}{x-8} = -1$$

$$\Rightarrow \frac{4}{x-8} = -1$$

$$\Rightarrow -x + 8 = 4$$

$$\Rightarrow x = 4$$

9. Show that the given points form a right angled triangle and check whether they satisfies pythagoras theorem

(i) A $(1, -4)$, B $(2, -3)$ and C $(4, -7)$

(ii) L $(0, 5)$, M $(9, 12)$ and N $(3, 14)$

Solution :

i) Given A $(1, -4)$, B $(2, -3)$, C $(4, -7)$

$$\text{Slope of AB} = \frac{-3+4}{2-1} = \frac{1}{1} = 1$$

$$\text{Slope of BC} = \frac{-7+3}{4-2} = \frac{-4}{2} = -2$$

$$\text{Slope of CA} = \frac{-7+4}{4-1} = \frac{-3}{3} = -1$$

$$\therefore \text{Slope of AB} \times \text{Slope of CA} = 1 \times -1 = -1$$

\therefore AB is perpendicular to AC.

$$\therefore \angle A = 90^\circ$$

$\therefore \triangle ABC$ is a right angled triangle.

ii) Given L $(0, 5)$, M $(9, 12)$, N $(3, 14)$

$$\text{Slope of LM} = \frac{12-5}{9-0} = \frac{7}{9}$$

$$\frac{14-12}{3-9} = \frac{2}{-6} = \frac{-1}{3}$$

$$\text{Slope of LN} = \frac{14-5}{3-0} = \frac{9}{3} = 3$$

\therefore Slope of MN \times Slope of LN

$$= \frac{-1}{3} \times 3$$

$$= -1$$

\therefore MN is perpendicular to LN.

$$\therefore \angle N = 90^\circ$$


$\therefore \triangle LMN$ is a right angled \triangle .

10. Show that the given points form a parallelogram :

A $(2.5, 3.5)$, B $(10, -4)$, C $(2.5, -2.5)$ and D $(-5, 5)$

Solution :

Plot the points and taking in anticlockwise direction.



Let A (2.5, 3.5), B (-5, 5), C (2.5, -2.5),
D (10, -4)

$$\text{Slope of AB} = \frac{5 - 3.5}{-5 - 2.5} = \frac{1.5}{-7.5} = \frac{-1}{5}$$

$$\text{Slope of CD} = \frac{-4 + 2.5}{10 - 2.5} = \frac{-1.5}{7.5} = \frac{-1}{5}$$

\therefore

\therefore AB and CD are parallel.

Also,

$$\text{Slope of AD} = \frac{-4 - 3.5}{10 - 2.5} = \frac{-7.5}{7.5} = -1$$

$$\text{Slope of BC} = \frac{-2.5 - 5}{2.5 + 5} = \frac{-7.5}{7.5} = -1$$

\therefore Slope of AD = Slope of BC.

\therefore AD and BC are parallel.

\therefore ABCD is a parallelogram.

11. If the points A (2, 2), B (-2, -3), C (1, -3) and D (x, y) form a parallelogram then find the value of x and y.

Solution :

Given A (2, 2), B (-2, -3), C (1, -3), D (x, y) form a parallelogram.

\therefore Slope of AB = Slope of CD.

$$\Rightarrow \frac{-3 - 2}{-2 - 2} = \frac{y + 3}{x - 1}$$

$$\Rightarrow \frac{-5}{-4} = \frac{y + 3}{x - 1}$$

$$\Rightarrow \frac{5}{4} = \frac{y + 3}{x - 1}$$

$$\Rightarrow 5x - 5 = 4y + 12$$

$$\Rightarrow 5x - 4y = 17 \quad \text{--- (1)}$$

Also

Slope of AD = Slope of BC

$$\Rightarrow \frac{y - 2}{x - 2} = \frac{-3 + 3}{1 + 2}$$

$$\Rightarrow \frac{y - 2}{x - 2} = 0$$

$$\Rightarrow y - 2 = 0$$

Sub. in (1)

$$5x - 8 = 17$$

$$\Rightarrow 5x = 25$$

$$\therefore x = 5$$


$$\therefore x = 5, y = 2$$

12. Let A (3, -4), B (9, -4), C (5, -7) and D (7, -7). Show that ABCD is a trapezium.

Solution :

Given points are (3, -4), (9, -4), (5, -7), (7, -7)

Plotting the given points in a plane and taking in anti clockwise direction.



Let A (9, -4), B (3, -4), C (5, -7) D (7, -7)

$$\text{Slope of AB} = \frac{-4 + 4}{3 - 9} = 0$$

$$\text{Slope of CD} = \frac{-7 + 7}{7 - 5} = 0$$

\therefore Slope of AB = Slope of CD.

\therefore AB and CD are parallel.

$$\text{Slope of AD} = \frac{-7 + 4}{7 - 9} = \frac{-3}{-2} = \frac{3}{2}$$

$$\text{Slope of BC} = \frac{-7 + 4}{5 - 3} = \frac{-3}{2}$$

\therefore Slope of AD \neq Slope of BC


\therefore One pair of opposite sides is equal.

\therefore ABCD is a trapezium.

- 13. A quadrilateral has vertices at A (-4,-2) B(5, -1) , C(6, 5) and D(-7, 6). Show that the mid-points of its sides form a parallelogram.**

Solution :

Given points are (- 4, - 2), (5, -1), (6, 5), (- 7, 6) which forms a quadrilateral.



$$\text{Mid point of AB} = \left(\frac{-4 + 5}{2}, \frac{-2 - 1}{2} \right) = \left(\frac{1}{2}, \frac{-3}{2} \right) \text{P}$$

$$\text{Mid point of BC} = \left(\frac{5 + 6}{2}, \frac{-1 + 5}{2} \right) = \left(\frac{11}{2}, 2 \right) \text{Q}$$

$$\text{Midpoint of CD} = \left(\frac{-7 + 6}{2}, \frac{6 + 5}{2} \right) = \left(\frac{-1}{2}, \frac{11}{2} \right) \text{R}$$

$$\text{Midpoint of AD} = \left(\frac{-7 - 4}{2}, \frac{6 - 2}{2} \right) = \left(\frac{-11}{2}, 2 \right) \text{S}$$

To prove : PQRS is a parallelogram.

$$\begin{aligned} \text{Slope of PQ} &= \frac{2 + \frac{3}{2}}{\frac{11}{2} - \frac{1}{2}} \\ &= \frac{\frac{7}{2}}{\frac{10}{2}} \\ &= \frac{7}{10} \end{aligned}$$

$$\begin{aligned} \text{Slope of SR} &= \frac{2 - \frac{11}{2}}{\frac{-11}{2} + \frac{1}{2}} \\ &= \frac{\frac{-7}{2}}{\frac{-10}{2}} \\ &= \frac{7}{10} \end{aligned}$$

\therefore Slope of PQ = Slope of SR

\therefore PQ and SR are parallel.

Also,

$$\begin{aligned}\text{Slope of PS} &= \frac{2 + \frac{3}{2}}{\frac{-11}{2} - \frac{1}{2}} \\ &= \frac{\frac{7}{2}}{\frac{-12}{2}} \\ &= \frac{-7}{12}\end{aligned}$$

$$\begin{aligned}\text{Slope of QR} &= \frac{\frac{11}{2} - 2}{\frac{-1}{2} - \frac{11}{2}} \\ &= \frac{\frac{7}{2}}{\frac{-12}{2}} \\ &= \frac{-7}{12}\end{aligned}$$


\therefore Slope of PS = Slope of QR

\therefore PS and QR are parallel.

\therefore PQRS is a parallelogram.

- 14. PQRS is a rhombus. Its diagonals PR and QS intersect at the point M and satisfy $QS = 2PR$. If the coordinates of S and M are (1, 1) and (2, -1) respectively, find the coordinates of P.**

Solution :



Given, In rhombus PQRS, diagonals PR and QS meet at M such that $QS = 2PR$.

Also, given S (1, 1), M (2, -1)

Let Q be (x, y)

Midpoint of SQ = M (\because rhombus)

$$\left(\frac{x+1}{2}, \frac{y+1}{2} \right) = (2, -1)$$

$$\Rightarrow x + 1 = 4 \quad y + 1 = -2$$

$$\Rightarrow x = 3, \quad y = -3$$

\therefore Q is (3, -3)

Since $QS = 2PR$

$$QS^2 = 4 \cdot PR^2$$

$$(3 - 1)^2 + (-3 - 1)^2 = 4 \cdot PR^2$$

$$\therefore PR^2 = 5$$

$$\therefore PR = \sqrt{5}$$

$$\Rightarrow PM = \frac{\sqrt{5}}{2}, \quad M(2, -1)$$

Let P be (l, m)

III. EQUATIONS OF STRAIGHT LINES:

Key Points

- ✓ Equation of OY(Y axis) is $x = 0$.
- ✓ Equation of OX (X axis) is $y = 0$.
- ✓ Equation of a straight line parallel to X axis is $y = b$.
- ✓ If $b > 0$, then the line $y=b$ lies above the X axis, If $b < 0$, then the line $y=b$ lies below the X axis, If $b = 0$, then the line $y=b$ is the X axis itself.
- ✓ Equation of a Straight line parallel to the Y axis is $x = c$.
If $c > 0$, then the line $x=c$ lies right to the side of the Y axis
If $c < 0$, then the line $x=c$ lies left to the side of the Y axis
If $c = 0$, then the line $x=c$ is the Y axis itself.
- ✓ Slope-Intercept Form A line with slope m and y intercept c can be expressed through the equation $y = mx + c$.
- ✓ Point-Slope form $y - y_1 = m(x - x_1)$.
- ✓ $\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$ is the equation of the line in two-point form.
- ✓ Intercept Form $\frac{x}{a} + \frac{y}{b} = 1$.

Example 5.17

Find the equation of a straight line passing through (5,7) and is (i) parallel to X axis (ii) parallel to Y axis.

Solution :

- (i) The equation of any straight line parallel to X axis is $y=b$.

Since it passes through (5,7), $b = 7$.

Therefore, the required equation of the line is $y=7$.

- (ii) The equation of any straight line parallel to Y axis is $x=c$

Since it passes through (5,7), $c = 5$

Therefore, the required equation of the line is $x = 5$.

Example 5.18

Find the equation of a straight line whose (i) Slope is 5 and y intercept is -9 (ii) Inclination is 45° and y intercept is 11

Solution :

- (i) Given, Slope = 5, y intercept, $c = -9$

Therefore, equation of a straight line is

$$y = mx + c$$

$$y = 5x - 9 \text{ gives } 5x - y - 9 = 0$$

- (ii) Given, $\theta = 45^\circ$, y intercept, $c = 11$

$$\text{Slope } m = \tan \theta = \tan 45^\circ = 1$$

Therefore, equation of a straight line is of the form $y = mx + c$

Hence we get, $y = x + 11$ gives $x - y + 11 = 0$.

Example 5.19

Calculate the slope and y intercept of the straight line $8x - 7y + 6 = 0$.

Solution :

Equation of the given straight line is
 $8x - 7y + 6 = 0$

$7y = 8x + 6$ (bringing it to the form

$y = mx + c$)

Slope $m = \frac{8}{7}$ and y intercept $c = \frac{6}{7}$

Example 5.20

The graph relates temperatures y (in Fahrenheit degree) to temperatures x (in Celsius degree)

(a) Find the slope and y intercept (b) Write an equation of the line (c) What is the mean temperature of the earth in Fahrenheit degree if its mean temperature is 25° Celsius?

Solution :


(a) From the figure, slope

$$= \frac{\text{change in } y \text{ coordinate}}{\text{change in } x \text{ coordinate}}$$

$$= \frac{68 - 32}{20 - 0} = \frac{36}{20} = \frac{9}{5} = 1.8$$

The line crosses the Y axis at $(0, 32)$

So the slope is $\frac{9}{5}$ and y intercept is 32.



(b) Use the slope and y intercept to write an equation

The equation is $y = \frac{9}{5}x + 32$.

(c) In Celsius, the mean temperature of the earth is 25° . To find the mean temperature in Fahrenheit, we find the value of y when $x = 25$.

$$y = \frac{9}{5}x + 32$$

$$y = \frac{9}{5}(25) + 32$$

$$y = 77$$

Therefore, the mean temperature of the earth is 77° F .

Example 5.21

point $(3, -4)$ and having slope $-\frac{5}{7}$.

Solution :

Given, $(x, y) = (3, -4)$ and $m = -\frac{5}{7}$

The equation of the point-slope form of the straight line is $y - y_1 = m(x - x_1)$

we write it as $y + 4 = -\frac{5}{7}(x - 3)$

gives us $5x + 7y + 13 = 0$.


Example 5.22

Find the equation of a line passing through the point $A(1, 4)$ and perpendicular to the line joining points $(2, 5)$ and $(4, 7)$.

Solution :

Let the given points be $A(1, 4)$, $B(2, 5)$ and $C(4, 7)$.

$$\text{Slope of line BC} = \frac{7-5}{4-2} = \frac{2}{2} = 1.$$



Let m be the slope of the required line.

Since the required line is perpendicular to BC ,

$$m \times 1 = -1$$

$$m = -1$$

The required line also pass through the point $A(1,4)$.

The equation of the required straight line is

$$y - y_1 = m(x - x_1)$$

$$y - 4 = -1(x - 1)$$

$$y - 4 = -x + 1$$

$$\text{we get, } x + y - 5 = 0.$$

Example 5.23

Find the equation of a straight line passing through $(5, -3)$ and $(7, -4)$.

Solution :

The equation of a straight line passing through the two points (x_1, y_1) and (x_2, y_2) is

$$\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$$

Substituting the points we get,

$$\frac{y + 3}{-4 + 3} = \frac{x - 5}{7 - 5}$$

$$\text{gives } 2y + 6 = -x + 5$$

$$\text{Therefore, } x + 2y + 1 = 0.$$




Example 5.24

Two buildings of different heights are located at opposite sides of each other. If a heavy rod is attached joining the terrace of the buildings from $(6, 10)$ to $(14, 12)$, find the equation of the rod joining the buildings ?

Solution :

Let $A(6,10)$, $B(14,12)$ be the points denoting the terrace of the buildings.



The equation of the rod is the equation of the straight line passing through $A(6,10)$ and $B(14,12)$

$$\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1} \text{ gives } \frac{y - 10}{12 - 10} = \frac{x - 6}{14 - 6}$$

$$\frac{y - 10}{2} = \frac{x - 6}{8}$$

$$\text{Therefore, } x - 4y + 34 = 0.$$

$$\text{Hence, equation of the rod is } x - 4y + 34 = 0.$$

Example 5.25

Find the equation of a line which passes through $(5,7)$ and makes intercepts on the axes equal in magnitude but opposite in sign.

Solution :

Let the x intercept be ' a ' and y intercept be ' $-a$ '.

The equation of the line in intercept form is

$$\frac{x}{a} + \frac{y}{b} = 1$$

gives $\frac{x}{a} + \frac{y}{-a} = 1$ (Here $b = -a$)

Therefore, $x - y = a$ (1)

Since (1) passes through (5,7)

Therefore, $5 - 7 = a$ gives $a = -2$

Thus the required equation of the straight line is $x - y = -2$; or $x - y + 2 = 0$.

Example 5.26

Find the intercepts made by the line $4x - 9y + 36 = 0$ on the coordinate axes.

Solution :

Equation of the given line is $4x - 9y + 36 = 0$.

(bringing it to the normal form)

Dividing by -36 we get, $\frac{x}{-9} + \frac{y}{4} = 1$ (1)

Comparing (1) with intercept form, we get x intercept $a = -9$; y intercept $b = 4$


Example 5.27

A mobile phone is put to use when the battery power is 100%. The percent of battery power 'y' (in decimal) remaining after using the mobile phone for x hours is assumed as $y = -0.25x + 1$

- Draw a graph of the equation.
- Find the number of hours elapsed if the battery power is 40%.
- How much time does it take so that the battery has no power?

Solution :

(i)



- (ii) To find the time when the battery power is 40%, we have to take $y = 0.40$

$$0.40 = -0.25x + 1 \text{ gives } 0.25x = 0.60$$

$$\text{we get, } x = \frac{0.60}{0.25} = 2.4 \text{ hours.}$$

- (iii) If the battery power is 0 then $y = 0$

Therefore, $0 = -0.25x + 1$ gives $-0.25x = 1$ hence $x = 4$ hours.

Thus, after 4 hours, the battery of the mobile phone will have no power.

Example 5.28

A line makes positive intercepts on coordinate axes whose sum is 7 and it passes through $(-3, 8)$. Find its equation.

Solution :

If a and b are the intercepts then $a + b = 7$ or $b = 7 - a$

$$\text{By intercept form } \frac{x}{a} + \frac{y}{b} = 1 \quad \dots(1)$$

$$\text{We have } \frac{x}{a} + \frac{y}{7-a} = 1$$

As this line pass through the point $(-3, 8)$, we have

$$\frac{-3}{a} + \frac{8}{7-a} = 1$$

gives $-3(7-a) + 8a = a(7-a)$.

$$-21 + 3a + 8a = 7a - a^2$$

$$\text{So, } a^2 + 4a - 21 = 0.$$

Solving this equation $(a-3)(a+7) = 0$

$$a = 3 \text{ or } a = -7$$

Solving this equation $(a-3)(a+7) = 0$

$$a = 3 \text{ or } a = -7$$

Since a is positive, we have $a = 3$ and

$$b = 7 - a = 7 - 3 = 4.$$


$$\text{Hence } \frac{x}{3} + \frac{y}{4} = 1$$

Therefore, $4x + 3y - 12 = 0$ is the required equation.

Example 5.29

A circular garden is bounded by East Avenue and Cross Road. Cross Road intersects North Street at D and East Avenue at E. AD is tangential to the circular garden at A(3, 10). Using the figure.

- (a) Find the equation of
- East Avenue.
 - North Street
 - Cross Road
- (b) Where does the Cross Road intersect the
- East Avenue ?
 - North Street ?



Solution :

(a) (i) East Avenue is the straight line joining C(0, 2) and B(7, 2). Thus the equation of East Avenue is obtained by using two-point form which is

$$\frac{y-2}{2-2} = \frac{x-0}{7-0}$$

$$\frac{y-2}{0} = \frac{x}{7} \text{ gives } y = 2$$

(ii) Since the point D lie vertically above C(0, 2). The x coordinate of D is 0.

Since any point on North Street has x coordinate value 0.

The equation of North Street is $x = 0$

(iii) To find equation of Cross Road.

Center of circular garden M is at (7, 7),

We first find slope of MA, which we call m_1

$$\text{Thus } m_1 = \frac{10-7}{3-7} = \frac{-3}{4}$$

Since the Cross Road is perpendicular to MA, if m_2 is the slope of the Cross Road then,

$$m_1 m_2 = -1 \text{ gives } \frac{-3}{4} m_2 = -1 \text{ so } m_2 = \frac{4}{3}$$

Now, the cross road has slope $\frac{4}{3}$ and it passes through the point A(3, 10).

The equation of the Cross Road is

$$y - 10 = \frac{4}{3}(x - 3).$$

$$3y - 30 = 4x - 12$$

$$\text{Hence, } 4x - 3y + 18 = 0$$

(b) (i) If D is (0, k) then D is a point on the Cross Road.

Therefore, substituting $x = 0$, $y = k$ in the equation of Cross Road,

$$\text{we get, } 0 - 3k + 18 = 0$$

$$\text{Value of } k = 6$$

Therefore, D is (0, 6)

(ii) To find E, let E be (q, 2)

Put $y = 2$ in the equation of the Cross Road,

$$\text{we get, } 4q - 6 + 18 = 0$$

$$4q = -12 \text{ gives } q = -3$$

Therefore, The point E is (-3, 2)

Thus the Cross Road meets the North Street at D(0, 6) and East Avenue at E (-3, 2).

EXERCISE 5.3

passing through the mid-point of a line segment joining the points (1, -5), (4, 2) and parallel to (i) X axis (ii) Y axis

Solution :

Mid point of the line joining the points (1, -5), (4, 2) is

$$= \left(\frac{1+4}{2}, \frac{-5+2}{2} \right)$$

$$= \left(\frac{5}{2}, \frac{-3}{2} \right)$$

i) Equation of straight line passing through

$$\left(\frac{5}{2}, \frac{-3}{2} \right) \text{ and}$$

a) Parallel to x-axis is

$$y = -\frac{3}{2} \Rightarrow 2y + 3 = 0$$

b) Parallel to y-axis is

$$x = \frac{5}{2} \Rightarrow 2x - 5 = 0$$

2. The equation of a straight line is $2(x - y) + 5 = 0$. Find its slope, inclination and intercept on the Y axis.

Solution :

Given equation of a straight line is

$$2(x - y) + 5 = 0$$

$$\Rightarrow 2x - 2y + 5 = 0 \quad \text{--- (1)}$$

$$\begin{aligned} \text{i) Slope of the line} &= \frac{-\text{coefficient of } x}{\text{coefficient of } y} \\ &= \frac{-2}{-2} \\ &= 1 \end{aligned}$$

ii) Slope of the line = 1

$$\tan \theta = 1$$

$$\therefore \theta = 45^\circ.$$

iii) Interecept on y-axis

Put $x = 0$ in (1)

$$-2y + 5 = 0$$

$$\Rightarrow -2y = -5$$

$$\Rightarrow y = \frac{5}{2}$$

$$\therefore y\text{-intercept} = \frac{5}{2}$$

3. Find the equation of a line whose inclination is 30° and making an intercept -3 on the Y axis.

Solution :

$$\text{Given } \theta = 30^\circ \Rightarrow m = \tan 30^\circ = \frac{1}{\sqrt{3}} \text{ and}$$

$$y\text{-intercept} = -3$$

The required equation of the line is

$$y = mx + c$$

$$\Rightarrow y = \frac{1}{\sqrt{3}}x - 3$$

$$\Rightarrow \sqrt{3}y = x - 3\sqrt{3}$$

$$\Rightarrow x - \sqrt{3}y - 3\sqrt{3} = 0$$

4. Find the slope and y intercept of $\sqrt{3}x + (1 - \sqrt{3})y = 3$.

Solution :

Given line is $\sqrt{3}x + (1 - \sqrt{3})y - 3 = 0$

$$\Rightarrow (1 - \sqrt{3})y = -\sqrt{3}x + 3$$

$$\Rightarrow y = \frac{-\sqrt{3}}{1 - \sqrt{3}}x + \frac{3}{1 - \sqrt{3}}$$

This is of the form $y = mx + c$.

$$\begin{aligned} m &= \frac{-\sqrt{3}}{1 - \sqrt{3}} & c &= \frac{3}{1 - \sqrt{3}} \\ &= \frac{\sqrt{3}}{\sqrt{3} - 1} \cdot \frac{\sqrt{3} + 1}{\sqrt{3} + 1} & &= \frac{3}{1 - \sqrt{3}} \cdot \frac{1 + \sqrt{3}}{1 + \sqrt{3}} \\ &= \frac{3\sqrt{3} + 3}{2} & &= \frac{3 + 3\sqrt{3}}{-2} \end{aligned}$$

$$\therefore \text{Slope} = \frac{3 + \sqrt{3}}{2}, \text{ y-intercept} = \frac{3 + 3\sqrt{3}}{-2}$$

5. Find the value of 'a', if the line through $(-2, 3)$ and $(8, 5)$ is perpendicular to $y = ax + 2$.

Solution :

Slope of the line joining $(-2, 3)$, $(8, 5)$.

$$\begin{aligned} &= \frac{5 - 3}{8 + 2} \\ &= \frac{2}{10} \\ m_1 &= \frac{1}{5} \end{aligned}$$


Slope of the line $y = ax + 2$ is $m_2 = a$.

Since the 2 lines are perpendicular.

$$\begin{aligned} m_1 m_2 &= -1 \\ \Rightarrow \frac{1}{5} \times a &= -1 \\ \Rightarrow a &= -5 \end{aligned}$$

6. The hill in the form of a right triangle has its foot at $(19, 3)$. The inclination of the hill to the ground is 45° . Find the equation of the hill joining the foot and top.

Solution :



C – Foot of the hill.

$$\therefore \text{Slope of AC} = m = \tan 45^\circ = 1$$

\therefore Equation of AC whose slope 1 and passing through C $(19, 3)$ is

$$\begin{aligned} \Rightarrow y - 3 &= 1(x - 19) \\ \Rightarrow x - y - 16 &= 0. \end{aligned}$$

7. Find the equation of a line through the given pair of points

(i) $\left(2, \frac{2}{3}\right)$ and $\left(\frac{-1}{2}, -2\right)$

(ii) $(2, 3)$ and $(-7, -1)$

Solution :

Given points are

$$\left(2, \frac{2}{3}\right), \left(\frac{-1}{2}, -2\right)$$

The equation of the line passing through 2 points

$$\begin{aligned} \frac{y - \frac{2}{3}}{-2 - \frac{2}{3}} &= \frac{x - 2}{\frac{-1}{2} - 2} \\ \Rightarrow \frac{3y - 2}{-8} &= \frac{x - 2}{\frac{-5}{2}} \end{aligned}$$

$$\Rightarrow \frac{3y-2}{-8} = \frac{2x-4}{-5}$$

$$\Rightarrow 15y - 10 = 16x - 32$$

$$\Rightarrow 16x - 15y - 22 = 0$$

- ii) Given points are (2, 3), (-7, -1)

Equation through 2 points is

$$\frac{y-3}{-1-3} = \frac{x-2}{-7-2}$$


$$\Rightarrow \frac{y-3}{-4} = \frac{x-2}{-9}$$

$$\Rightarrow 9y - 27 = 4x - 8$$

$$\Rightarrow 4x - 9y + 19 = 0$$

8. A cat is located at the point(-6,-4) in xy plane. A bottle of milk is kept at (5,11). The cat wish to consume the milk travelling through shortest possible distance. Find the equation of the path it needs to take its milk.

Solution :



C (-6, -4) is the position of cat.

M (5, 11) is the position of milk.

Equation of the path CM is

$$\frac{y-y_1}{y_2-y_1} = \frac{x-x_1}{x_2-x_1}$$

$$\Rightarrow \frac{y+4}{11+4} = \frac{x+6}{5+6}$$


$$\Rightarrow \frac{y+4}{15} = \frac{x+6}{11}$$

$$\Rightarrow 15x + 90 = 11y + 44$$

$$\Rightarrow 15x - 11y + 46 = 0$$

9. Find the equation of the median and altitude of $\triangle ABC$ through A where the vertices are A(6,2), B(-5,-1) and C(1,9)

Solution :



- i) Equation of the median through A.
mid point of BC = $\left(\frac{-5+1}{2}, \frac{-1+9}{2} \right)$

$$= D(-2, 4)$$

Equation of AD is [A (6, 2), D (-2, 4).

$$\frac{y-y_1}{y_2-y_1} = \frac{x-x_1}{x_2-x_1}$$

$$\Rightarrow \frac{y-2}{4-2} = \frac{x-6}{-2-6}$$

$$\Rightarrow \frac{y-2}{2} = \frac{x-6}{-8}$$

$$= \frac{y-2}{1} = \frac{x-6}{-4}$$

$$\Rightarrow x - 6 = -4y + 8$$

$$\Rightarrow x + 4y - 14 = 0$$

- ii) Equation of altitude through 'A'

$$\text{Slope of BC} = \frac{9+1}{1+5} = \frac{10}{6} = \frac{5}{3}$$

Since $AD \perp BC$, slope of $AD = \frac{-3}{5}$

and A is (6, 2).

\therefore Equation of altitude AD is

$$y - y_1 = m(x - x_1)$$

$$\Rightarrow y - 2 = \frac{-3}{5}(x - 6)$$

$$\Rightarrow 5y - 10 = -3x + 18$$

$$\Rightarrow 3x + 5y - 28 = 0$$

- 10. Find the equation of a straight line which has slope $\frac{-5}{4}$ and passing through the point $(-1, 2)$.**

Solution :

Given slope of the line is $\frac{-5}{4}$ and $(-1, 2)$ is

\therefore its equation is $y - y_1 = m(x - x_1)$

$$\Rightarrow y - 2 = \frac{-5}{4}(x + 1)$$

$$\Rightarrow 4y - 8 = -5x - 5$$

$$\Rightarrow 5x + 4y - 3 = 0$$

- 11. You are downloading a song. The percent y (in decimal form) of mega bytes remaining to get downloaded in x seconds is given by $y = -0.1x + 1$.**

(i) graph the equation.

(ii) find the total MB of the song.

(iii) after how many seconds will 75% of the song gets downloaded?

(iv) after how many seconds the song will be downloaded completely?


Solution :

Given $y = -0.1x + 1$.

where x - time (in sec.)

y - remaining data to be downloaded.

x	0	2	4	6	8	10
y	1	0.8	0.6	0.4	0.2	0



ii) $y = -0.1x + 1$

\therefore Total MB of the song = 1 MB

iii) 75% of MB downloaded.

\therefore 25% of MB to be downloaded

\therefore Put $y = 0.25$ in (1)

$$\Rightarrow 0.25 = -0.1x + 1$$

$$\Rightarrow 0.1x = 0.75$$

$$x = \frac{0.75}{0.10} = \frac{15}{2} = 7.5$$

\therefore Required time = 7.5 sec.

iv) Time when songs completely downloaded

Put $y = 0$ in (1)

$$\text{ie, } 0.1x = 1$$

$$x = 10 \text{ sec.}$$

\therefore Songs will be downloaded completely after 10 sec.

- 12. Find the equation of a line whose intercepts on the x and y axes are given below.**

(i) 4, -6 (ii) -5, $\frac{3}{4}$

Solution :

- i) Given x-intercept = 4 = a

$$y - \text{intercept} = -6 = b$$

Equation of line in intercept form is

$$\frac{x}{a} + \frac{y}{b} = 1$$

$$\Rightarrow \frac{x}{4} + \frac{y}{-6} = 1$$

$$\Rightarrow \frac{x}{4} - \frac{y}{6} = 1$$

$$\Rightarrow \frac{3x - 2y}{12} = 1$$

$$\Rightarrow 3x - 2y - 12 = 0$$

- ii) Given x-intercept = -5 = a

$$y - \text{intercept} = \frac{3}{4} = b$$

Equation of line in intercept form is

$$\frac{x}{a} + \frac{y}{b} = 1$$

$$\Rightarrow \frac{x}{-5} + \frac{y}{\frac{3}{4}} = 1$$

$$\Rightarrow \frac{x}{-5} - \frac{4y}{3} = 1$$

$$\Rightarrow \frac{-3x + 20y}{15} = 1$$

$$\Rightarrow -3x + 20y - 15 = 0$$

$$\Rightarrow 3x - 20y + 15 = 0$$

- 13. Find the intercepts made by the following lines on the coordinate axes.**

(i) $3x - 2y - 6 = 0$ (ii) $4x + 3y + 12 = 0$

Solution :

Method - 1 :

$$3x - 2y = 6$$

$$\Rightarrow \frac{3x}{6} - \frac{2y}{6} = 1$$

$$\Rightarrow \frac{x}{2} + \frac{y}{-3} = 1$$

$$\therefore x\text{-int} = 2, y\text{-int} = -3$$

Method - 2 :

$$3x - 2y = 6$$

$$\text{Put } y = 0 \Rightarrow 3x = 6$$

$$\Rightarrow x = 2 \quad (x\text{-int})$$

$$\text{Put } x = 0 \Rightarrow -2y = 6$$

$$\Rightarrow y = -3 \quad (y\text{-int})$$

- ii) Given line is $4x + 3y + 12 = 0$

$$\Rightarrow 4x + 3y = -12$$

$$\Rightarrow \frac{4x}{-12} + \frac{3y}{-12} = 1$$

$$\Rightarrow \frac{x}{-3} + \frac{y}{-4} = 1$$

$$\therefore x\text{-int} = -3, y\text{-int} = -4.$$

- 14. Find the equation of a straight line**

(i) passing through (1,-4) and has intercepts which are in the ratio 2:5

(ii) passing through (-8, 4) and making equal intercepts on the coordinate axes

Solution :

- i) Required line is passing through (1, -4) and has intercepts in the ratio 2 : 5.

Equation of line in intercept form is

$$\begin{aligned} \frac{x}{a} + \frac{y}{b} &= 1 & \text{where } a : b &= 2 : 5 \\ \Rightarrow \frac{x}{a} + \frac{y}{\frac{5a}{2}} &= 1 & \Rightarrow \frac{a}{b} &= \frac{2}{5} \\ \Rightarrow \frac{x}{a} + \frac{2y}{5a} &= 1 & \Rightarrow b &= \frac{5a}{2} \\ \Rightarrow \frac{x}{a} + \frac{2y}{5a} &= 1 \dots(1) \end{aligned}$$

Since (1) passes through $(1, -4)$

$$\begin{aligned} \frac{1}{a} - \frac{8}{5a} &= 1 \\ \Rightarrow \frac{5-8}{5a} &= 1 \\ \Rightarrow 5a &= -3 \\ a &= \frac{-3}{5} \end{aligned}$$

$$\therefore (1) \Rightarrow \frac{-3}{5} + \frac{2y}{-3} = 1$$

$$\begin{aligned} \Rightarrow \frac{5x}{-3} + \frac{2y}{-3} &= 1 \\ \Rightarrow 5x + 2y &= -3 \\ \Rightarrow 5x + 2y + 3 &= 0 \end{aligned}$$

ii) Required line is passing through $(-8, 4)$ and making equal intercepts on the axes.

Equation of line in intercept form is

$$\begin{aligned} \frac{x}{a} + \frac{y}{b} &= 1 \quad \text{where } a = b \\ \Rightarrow \frac{x}{a} + \frac{y}{a} &= 1 \\ \Rightarrow x + y &= a \dots(1) \end{aligned}$$

Since (1) passes through $(-8, 4)$

$$-8 + 4 = a$$

$$a = -4$$

$$\therefore (1) \Rightarrow x + y = -4 \Rightarrow x + y + 4 = 0$$

IV. GENERAL FORM OF A STRAIGHT LINE

Key Points

- ✓ The equation of all lines parallel to the line $ax + by + c = 0$ can be put in the form $ax + by + k = 0$ for different values of k .
- ✓ The equation of all lines perpendicular to the line $ax + by + c = 0$ can be written as $bx - ay + k = 0$ for different values of k .
- ✓ Slope of a straight line $m = \frac{-\text{coefficient of } x}{\text{coefficient of } y}$
- ✓ y intercept $= \frac{-\text{constant term}}{\text{coefficient of } y}$

Example 5.30

Find the slope of the straight line $6x + 8y + 7 = 0$.

Solution :

$$\text{Given } 6x + 8y + 7 = 0$$

$$\text{slope } m = \frac{-\text{coefficient of } x}{\text{coefficient of } y} = -\frac{6}{8} = -\frac{3}{4}$$

Therefore, the slope of the straight line is $-\frac{3}{4}$.

Example 5.31

Find the slope of the line which is

- (i) parallel to $3x - 7y = 11$
 (ii) perpendicular to $2x - 3y + 8 = 0$.

Solution :

- (i) Given straight line is $3x - 7y = 11$

$$\text{gives } 3x - 7y - 11 = 0$$

$$\text{Slope } m = \frac{-3}{-7} = \frac{3}{7}$$

Since parallel lines have same slopes, slope of any line parallel to

$$3x - 7y = 11 \text{ is } \frac{3}{7}.$$

- (ii) Given straight line is $2x - 3y + 8 = 0$

$$\text{Slope } m = \frac{-2}{-3} = \frac{2}{3}$$

Since product of slopes is -1 for perpendicular lines, slope of any line perpendicular to

$$2x - 3y + 8 = 0 \text{ is } \frac{-1}{\frac{2}{3}} = \frac{-3}{2}$$

Example 5.32

Show that the straight lines $2x + 3y - 8 = 0$ and $4x + 6y + 18 = 0$ are parallel.

Solution :

Slope of the straight line $2x + 3y - 8 = 0$ is

$$m_1 = \frac{-\text{coefficient of } x}{\text{coefficient of } y}$$

$$m_1 = \frac{-2}{3}$$

Slope of the straight line $4x + 6y + 18 = 0$ is

$$m_2 = \frac{-4}{6} = \frac{-2}{3}$$

$$\text{Here, } m_1 = m_2$$

That is, slopes are equal. Hence, the two straight lines are parallel.

Example 5.33

Show that the straight lines $x - 2y + 3 = 0$ and $6x + 3y + 8 = 0$ are perpendicular.

Solution :

Slope of the straight line $x - 2y + 3 = 0$ is

$$m_1 = \frac{-1}{-2} = \frac{1}{2}$$

Slope of the straight line $6x + 3y + 8 = 0$ is

$$m_2 = \frac{-6}{3} = -2$$

$$\text{Now, } m_1 \times m_2 = \frac{1}{2} \times (-2) = -1$$

Hence, the two straight lines are perpendicular.

Example 5.34

Find the equation of the straight line parallel to the line $3x - 7y = 12$ and passing through the point $(6, 4)$.

Solution :

Equation of the straight line, parallel to $3x - 7y - 12 = 0$ is $3x - 7y + k = 0$

Since it passes through the point $(6, 4)$

$$3(6) - 7(4) + k = 0$$

$$k = 28 - 18 = 10$$

Therefore, equation of the required straight line is $3x - 7y + 10 = 0$.

Example 5.35

Find the equation of a straight line perpendicular to the line $y = \frac{4}{3}x - 7$ and passing through the point $(7, -1)$.

Solution :

The equation $y = \frac{4}{3}x - 7$ can be written as $4x - 3y - 21 = 0$.

Equation of a straight line perpendicular to $4x - 3y - 21 = 0$ is $3x + 4y + k = 0$

Since it passes through the point $(7, -1)$,

$$21 - 4 + k = 0 \text{ we get, } k = -17$$

Therefore, equation of the required straight line is $3x + 4y - 17 = 0$.

Example 5.36

Find the equation of a straight line parallel to Y axis and passing through the point of intersection of the lines $4x + 5y = 13$ and $x - 8y + 9 = 0$.

Solution :

Given lines $4x + 5y - 13 = 0 \dots(1)$

$$x - 8y + 9 = 0 \dots(2)$$

$$\begin{array}{rcccl} & x & & y & 1 \\ 5 & & -13 & & 4 \\ -8 & & 9 & & 1 \end{array} \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} \searrow \\ \nearrow \end{array} \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} 5 \\ -8 \end{array}$$

$$\frac{x}{45 - 104} = \frac{y}{-13 - 36} = \frac{1}{-32 - 5}$$

$$\frac{x}{-59} = \frac{y}{-49} = \frac{1}{-37}$$

$$x = \frac{59}{37}, y = \frac{49}{37}$$

Therefore, the point of intersection

$$(x, y) = \left(\frac{59}{37}, \frac{49}{37} \right)$$

The equation of line parallel to Y axis is $x = c$.

It passes through $(x, y) = \left(\frac{59}{37}, \frac{49}{37} \right)$.

$$\text{Therefore, } c = \frac{59}{37}$$

The equation of the line is $x = \frac{59}{37}$ gives $37x - 59 = 0$.


Example 5.37

The line joining the points $A(0, 5)$ and $B(4, 1)$ is a tangent to a circle whose centre C is at the point $(4, 4)$ find

- the equation of the line AB.
- the equation of the line through C which is perpendicular to the line AB.
- the coordinates of the point of contact of tangent line AB with the circle.

Solution :

- Equation of line AB, $A(0, 5)$ and $B(4, 1)$



$$\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$$

$$\frac{y - 5}{1 - 5} = \frac{x - 0}{4 - 0}$$

$$4(y - 5) = -4x \text{ gives } y - 5 = -x$$

$$x + y - 5 = 0$$

- The equation of a line which is perpendicular to the line AB : $x + y - 5 = 0$ is $x - y + k = 0$

Since it is passing through the point $(4, 4)$, we have

$$4 - 4 + k = 0 \text{ gives } k = 0$$

The equation of a line which is perpendicular to AB and through C is

$$x - y = 0 \quad \dots(2)$$

(iii) The coordinate of the point of contact P of the tangent line AB with the circle is

$$x + y - 5 = 0 \text{ and } x - y = 0$$

$$\text{Solving, we get } x = \frac{5}{2} \text{ and } y = \frac{5}{2}$$

Therefore, the coordinate of the point of contact is $P\left(\frac{5}{2}, \frac{5}{2}\right)$.

EXERCISE 5.4

1. Find the slope of the following straight lines

$$(i) 5y - 3 = 0 \quad (ii) 7x - \frac{3}{17} = 0$$

Solution :

$$\begin{aligned} \text{Slope} &= \frac{\text{Co. eff. of } x}{\text{Co. eff. of } y} \\ &= \frac{-0}{5} \\ &= 0 \end{aligned}$$

$$(ii) \text{ Given line is } 7x - \frac{3}{17} = 0$$

$$\begin{aligned} \text{Slope} &= \frac{\text{Co. eff. of } x}{\text{Co. eff. of } y} \\ &= \frac{-7}{0} \\ &= \text{undefined} \end{aligned}$$

2. Find the slope of the line which is

(i) parallel to $y = 0.7x - 11$

(ii) perpendicular to the line $x = -11$.

Solution :

$$\begin{aligned} (i) \text{ Given line is } y &= 0.7x - 11 \\ \text{whose slope is } &0.7 \end{aligned}$$

\therefore Slope of the line parallel to

$$y = 0.7x - 11 \text{ is also } 0.7$$

(ii) Given line is $x = -11$,

whose slope is 0

\therefore Slope of the line perpendicular to

$$x = -11 \text{ is } \frac{-1}{0}$$

which is undefined.

3. Check whether the given lines are parallel or perpendicular

$$(i) \frac{x}{3} + \frac{y}{4} + \frac{1}{7} = 0 \text{ and } \frac{2x}{3} + \frac{y}{2} + \frac{1}{10} = 0$$

$$(ii) 5x + 23y + 14 = 0 \text{ and } 23x - 5y + 9 = 0$$

Solution :

(i) Given pair of lines

$$\frac{x}{3} + \frac{y}{4} + \frac{1}{7} = 0 \quad \frac{2x}{3} + \frac{y}{2} + \frac{1}{10} = 0$$

Their slope,

$$\begin{aligned} m_1 &= \frac{-1/3}{1/4} \quad \& \quad m_2 = \frac{-2/3}{1/2} \\ &= -4/3 \quad \quad \quad = -4/3 \end{aligned}$$

$$\therefore m_1 = m_2$$

\Rightarrow the 2 lines are parallel

(ii) Given lines are

$$5x + 23y + 14 = 0, \quad 23x - 5y + 9 = 0$$

Their slope,

$$m_1 = \frac{-5}{23} \quad m_2 = \frac{23}{5}$$

$$\therefore m_1 \times m_2 = -1$$

\therefore the 2 lines are perpendicular.

4. If the straight lines $12y = -(p + 3)x + 12$,

$12x - 7y = 16$ are perpendicular then find 'p'.

Solution :

Given lines

$$12y = -(p+3)x + 12,$$

$12x - 7y = 16$ are perpendicular

$$\Rightarrow (p+3)x + 12y = 12$$

$$m_1 = \frac{-(p+3)}{12} \quad m_2 = \frac{12}{7}$$

Since 2 lines are perpendicular,

$$m_1 \times m_2 = -1$$

$$\Rightarrow \frac{-(p+3)}{12} \times \frac{12}{7} = -1$$

$$\Rightarrow -(p+3) = -7$$

$$\Rightarrow p+3 = 7$$

$$\Rightarrow p = 4$$

5. Find the equation of a straight line passing through the point P(-5,2) and parallel to the line joining the points Q(3,-2) and R(-5, 4) .

Solution :

The required line is passing through P(-5, 2) and parallel to the line joining the points Q(3, -2), R(-5, 4)

$$\text{Slope of QR} = \frac{4+2}{-5-3} = \frac{6}{-8} = \frac{-3}{4}$$

\therefore Equn. of the line is

$$y - y_1 = m(x - x_1)$$

$$\Rightarrow y - 2 = \frac{-3}{4}(x + 5)$$

$$\Rightarrow 4y - 8 = -3x - 15$$

$$\Rightarrow 3x + 4y + 7 = 0$$

6. Find the equation of a line passing through (6,-2) and perpendicular to the line joining the points (6,7) and (2,-3).

Solution :

The required line is passing through (6, -2) and perpendicular to the line joining (6, 7), (2,-3)

\therefore Slope of the line joining (6, 7), (2, -3)

$$= \frac{-3-7}{2-6} = \frac{-10}{-4} = \frac{5}{2}$$

\therefore Slope of the line perpendicular to it is $\frac{-2}{5}$

\therefore Equation of the required line is

$$y + 2 = \frac{-2}{5}(x - 4)$$

$$\Rightarrow 5y + 10 = -2x + 12$$

$$\Rightarrow 2x + 5y - 2 = 0$$

7. A(-3, 0) B(10,-2) and C(12, 3) are the vertices of $\triangle ABC$. Find the equation of the altitude through A and B.

Solution :

Given vertices of Δ are

A(-3, 0), B(10, -2), C(12, 3)

Equation of altitude AD :

$$\text{Slope of BC} = \frac{3+2}{12-10} = \frac{5}{2}$$


\therefore Slope of AD is $= \frac{-2}{5}$ ($AD \perp BC$)

\therefore Equation of AD is

$$y - 0 = \frac{-2}{5}(x + 3)$$

$$5y = -2x - 6$$

$$\Rightarrow 2x + 5y + 6 = 0$$



∴ Equation of altitude BE is

$$\text{Slope of AC} = \frac{3-0}{12+3} = \frac{3}{15} = \frac{1}{5}$$

∴ Slope of BE = -5 ($\because BE \perp AC$)

∴ Equation of BE is

$$y + 2 = -5(x - 10)$$


$$\Rightarrow y + 2 = -5x + 50$$

$$\Rightarrow 5x + y - 48 = 0$$

8. Find the equation of the perpendicular bisector of the line joining the points A(-4,2) and B(6,-4).

Solution :

Given AB and CD are perpendicular & D is the midpoint of AB



$$\therefore D = \left(\frac{-4+6}{2}, \frac{2-4}{2} \right) = (1, -1)$$

$$\text{Slope of AB} = \frac{-4-2}{6+4} = \frac{-6}{10} = \frac{-3}{5}$$

$$\therefore \text{Slope of CD} = \frac{5}{3} \quad (\because CD \perp AB)$$

∴ Equation of perpendicular bisector

CD is

$$y + 1 = \frac{5}{3}(x - 1)$$

$$\Rightarrow 3y + 3 = 5x - 5$$

$$\Rightarrow 5x - 3y - 8 = 0$$

9. Find the equation of a straight line through the intersection of lines $7x + 3y = 10$, $5x - 4y = 1$ and parallel to the line $13x + 5y + 12 = 0$.

Solution :

The required line is passing through the intersection of the lines

$$7x + 3y = 10 \quad \dots\dots\dots (1)$$

$$5x - 4y = 1 \quad \dots\dots\dots (2)$$

and parallel to the line $13x + 5y + 12 = 0$

Solving (1) & (2)

$$(1) \times 4 \Rightarrow 28x + 12y = 40$$

$$(2) \times 3 \Rightarrow \frac{15x - 12y = 3}{43x = 43}$$

$$x = 1$$

Sub $x = 1$ in (1)

$$7(1) + 3y = 10$$

$$\Rightarrow 3y = 3$$

$$y = 1$$

∴ The required line is

Since it passes through (1, 1)

$$13 + 5 + k = 0$$

$$k = -18$$

$$\therefore 13x + 5y - 18 = 0$$

10. Find the equation of a straight line through the intersection of lines $5x - 6y = 2$, $3x + 2y = 10$ and perpendicular to the line $4x - 7y + 13 = 0$.

Solution :

Given lines are

$$5x - 6y = 2 \quad \dots\dots\dots (1)$$

$$3x + 2y = 10 \quad \dots\dots\dots (2)$$

$$(1) \Rightarrow 5x - 6y = 2$$

$$(2) \times 3 \Rightarrow \frac{9x + 6y = 30}{14x = 32}$$

$$x = \frac{16}{7}$$

Sub in (2)

$$\begin{aligned} 48/7 + 2y &= 10 \Rightarrow 2y = 10 - 48/7 \\ &\Rightarrow 2y = 22/7 \\ &y = 11/7 \end{aligned}$$

The required line is perpendicular to

$$4x - 7y + 13 = 0$$

Equation of the required line is

$$7x + 4y + k = 0$$

\therefore Since it passes through

$$\left(\frac{16}{7}, \frac{11}{7}\right)$$

$$\Rightarrow 7\left(\frac{16}{7}\right) + 4\left(\frac{11}{7}\right) + k = 0$$

$$\Rightarrow 16 + \frac{44}{7} + k = 0$$

$$\Rightarrow k = -16 - \frac{44}{7}$$

$$\Rightarrow k = \frac{-156}{7}$$

$$\therefore 7x + 4y - \frac{156}{7} = 0$$

$$\Rightarrow 49x + 28y - 156 = 0$$

- 11. Find the equation of a straight line joining the point of intersection of $3x + y + 2 = 0$ and $x - 2y - 4 = 0$ to the point of intersection of $7x - 3y = -12$ and $2y = x + 3$.**

Solution :

$$(1) \times 2 \Rightarrow 6x + 2y = -4$$

$$(2) \Rightarrow \frac{x - 2y = 4}{7x = 0}$$

$$x = 0$$

$$(1) \Rightarrow y = -2$$

\therefore The point of int. of (1) & (2) is $(0, -2)$

Now, to find the point of int. of the lines

$$7x - 3y = -12 \quad \dots\dots (3)$$

$$x - 2y + 3 = 0 \quad \dots\dots (4)$$

$$(3) \Rightarrow 7x - 3y = -12$$

$$(4) \times 7 \Rightarrow \frac{7x - 14y = -21}{11y = 9}$$

$$y = \frac{9}{11}$$

Sub in (4)

$$x - \frac{18}{11} + 3 = 0$$

$$x = \frac{18}{11} - 3 = \frac{-15}{11}$$

$$\left(\frac{-15}{11}, \frac{9}{11}\right)$$

The required equation of the line joining

$$(0, -2), \left(\frac{-15}{11}, \frac{9}{11}\right)$$

$$\frac{y + 2}{\frac{9}{11} + 2} = \frac{x - 0}{\frac{-15}{11}}$$

$$\Rightarrow \frac{y + 2}{31} = \frac{x}{-15}$$

$$\Rightarrow 31x = -15y - 30$$

$$\Rightarrow 31x + 15y + 30 = 0$$

- 12. Find the equation of a straight line through the point of intersection of the lines $8x + 3y = 18$, $4x + 5y = 9$ and bisecting the line segment joining the points $(5, -4)$ and $(-7, 6)$.**

Solution :

To find : The point of int. of

$$8x + 3y = 18 \quad \dots\dots (1)$$

$$4x + 5y = 9 \quad \dots\dots (2)$$

$$(1) \Rightarrow 8x + 3y = 18$$

$$(2) \times 2 \Rightarrow \begin{array}{r} 8x + 10y = 18 \\ -7y = 0 \\ y = 0 \end{array}$$

$$(2) \Rightarrow 4x = 9$$

$$\therefore x = \frac{9}{4}$$

 \therefore The point of int. of (1) & (2) in

$$\left(\frac{9}{4}, 0\right)$$

Mid point of the line joining

$$(5, -4), (-7, 6)$$

$$= \left(\frac{5-7}{2}, \frac{-4+6}{2}\right)$$

$$= (-1, 1)$$

Equation of the required line joining

$$\left(\frac{9}{5}, 0\right), (-1, 1)$$

$$\frac{y-0}{1} = \frac{x-\frac{9}{5}}{-1-\frac{9}{5}}$$

$$y = \frac{4x-9}{-13}$$

$$\Rightarrow 4x + 13y - 9 = 0$$

EXERCISE 5.5**Multiple choice questions :**

1. The area of triangle formed by the points $(-5,0)$, $(0,-5)$ and $(5,0)$ is


- (1) 0 sq.units (2) 25 sq.units
(3) 5 sq.units (4) none of these

Hint :**Ans : (2)**

$$\text{Area of ABC} = \frac{1}{2} \times b \times h$$

$$= \frac{1}{2} \times 10 \times 5$$


$$= 25 \text{ sq.units}$$



2. A man walks near a wall, such that the distance between him and the wall is 10

The path travelled by the man is

- (1) $x = 10$ (2) $y = 10$
(3) $x = 0$ (4) $y = 0$

Hint : **Ans : (1)**Equation of path travelled by the man is $x = 10$

3. The straight line given by the equation $x = 11$ is

- (1) parallel to X axis
(2) parallel to Y axis
(3) passing through the origin
(4) passing through the point $(0, 11)$

Hint :**Ans : (2)**Equation $x = C$ is a line parallel to y - axis

4. If $(5,7)$, $(3,p)$ and $(6,6)$ are collinear, then the value of p is

- (1) 3 (2) 6 (3) 9 (4) 12

Hint :**Ans : (3)**

A (5, 7), B (3, p), C (6, 6) are collinear

 \therefore Slope of AB = Slope of BC

$$\frac{p-7}{-2} = \frac{6-p}{3}$$

$$\Rightarrow 3p - 21 = -12 + 2p$$

$$\Rightarrow p = 9$$

5. The point of intersection of $3x - y = 4$ and $x + y = 8$ is

(1) (5,3) (2) (2,4) (3) (3,5) (4) (4,4)

Hint :**Ans : (3)**

Substitute and check the point to satisfy the given lines.

6. The slope of the line joining (12, 3), (4, a)

$$\frac{1}{8}$$

(1) 1 (2) 4 (3) -5 (4) 2

Hint :**Ans : (4)**

$$\text{Slope of } (12, 3), (4, a) = \frac{1}{8}$$

$$\Rightarrow \frac{a-3}{-8} = \frac{1}{8}$$

$$\Rightarrow a - 3 = -1$$

$$\Rightarrow a = 2$$

7. The slope of the line which is perpendicular to a line joining the points (0,0) and (-8,8) is

(1) -1 (2) 1 (3) $\frac{1}{3}$ (4) -8**Hint :****Ans : (2)**

Slope of the line joining (0, 0), (-8, 8)

$$= \frac{8-0}{-8-0}$$

$$= -1$$

 \therefore Slope of the line perpendicular to it = 1.

8. If slope of the line PQ is $\frac{1}{\sqrt{3}}$ then slope of the perpendicular bisector of PQ is

(1) $\sqrt{3}$ (2) $-\sqrt{3}$ (3) $\frac{1}{\sqrt{3}}$ (4) 0**Hint :****Ans : (2)**

$$\text{Slope of PQ} = \frac{+1}{\sqrt{3}}$$


$$\text{Slope of its perpendicular bisector} = -\sqrt{3}$$

9. If A is a point on the Y axis whose ordinate is 8 and B is a point on the X axis whose abscissae is 5 then the equation of the line AB is

(1) $8x + 5y = 40$ (2) $8x - 5y = 40$ (3) $x = 8$ (4) $y = 5$ Here $a = 5$, $b = 8$

$$\therefore \text{Equn. of the line is } \frac{x}{5} - \frac{y}{8} = 1$$

$$\Rightarrow 8x + 5y - 40 = 0$$



10. The equation of a line passing through the origin and perpendicular to the line $7x - 3y + 4 = 0$ is

(1) $7x - 3y + 4 = 0$ (2) $3x - 7y + 4 = 0$ (3) $3x + 7y = 0$ (4) $7x - 3y = 0$ **Hint :****Ans : (3)**

Equation of the line perpendicular to

$$7x - 3y + 4 = 0 \text{ is}$$

$$3x + 7y + k = 0$$

Since it passes through (0, 0), $k = 0$

$$\therefore 3x + 7y = 0$$

11. Consider four straight lines

(i) $l_2 : 3y = 4x + 5$ (ii) $l_2 : 4y = 3x - 1$

(iii) $l_3 : 4y + 3x + 7$ (iv) $l_4 : 4x + 3y = 2$

Which of the following statement is true?

(1) l_1 and l_2 are perpendicular(2) l_1 and l_4 are parallel(3) l_2 and l_4 are perpendicular(4) l_2 and l_3 are parallel*Hint :***Ans : (3)**

i) Slope of $l_1 = \frac{4}{3}$

ii) Slope of $l_2 = \frac{3}{4}$

iii) Slope of $l_3 = -\frac{3}{4}$

iv) Slope of $l_4 = \frac{1}{3}$

Here l_1 and l_3 are perpendicular l_2 and l_4 are perpendicular

But 3rd option is a contradiction

12. A straight line has equation $8y = 4x + 21$. Which of the following is true

(1) The slope is 0.5 and the y intercept is 2.6

(2) The slope is 5 and the y intercept is 1.6

(3) The slope is 0.5 and the y intercept is 1.6

(4) The slope is 5 and the y intercept is 2.6

*Hint :***Ans :**

(1)

Given equation is $8y = 4x + 21$

$$\Rightarrow y = \frac{1}{2}x + \frac{21}{8}$$

$$\Rightarrow y = 0.5x + 2.6$$

 \therefore Slope = 0.5, y - int = 2.6**13. When proving that a quadrilateral is a trapezium, it is necessary to show**

(1) Two sides are parallel.

(2) Two parallel and two non-parallel sides.

(3) Opposite sides are parallel.

(4) All sides are of equal length.

*Hint :***Ans : (2)**

A quadrilateral is trapezoid if one pair of opposite sides are parallel and another pair is non parallel.

14. When proving that a quadrilateral is a parallelogram by using slopes you must find

(1) The slopes of two sides

(2) The slopes of two pair of opposite sides

(3) The lengths of all sides

(4) Both the lengths and slopes of two side

*Hint :***Ans : (1)**

We should find the slopes of all the sides when proving a quadrilateral is a parallelogram.

15. (2, 1) is the point of intersection of two lines.

(1) $x - y - 3 = 0$; $3x - y - 7 = 0$

(2) $x + y = 3$; $3x + y = 7$

(3) $3x + y = 3$; $x + y = 7$

(4) $x + 3y - 3 = 0$; $x - y - 7 = 0$

*Hint :***Ans : (2)**


Substitute (2, 3) & check in all pair of lines.

UNIT EXERCISE - 5

1. PQRS is a rectangle formed by joining the points P(-1,-1), Q(-1, 4), R(5, 4) and S(5,-1). A, B, C and D are the mid-points of PQ, QR, RS and SP respectively. Is the quadrilateral ABCD a square, a rectangle or a rhombus? Justify your answer.

Solution :

Given P = (-1, -1) Q (-1, 4), R (5, 4), S (5, -1)



$$A = \text{Mid point of } PQ = \left(\frac{-1-1}{2}, \frac{-1+4}{2} \right) = \left(-1, \frac{3}{2} \right)$$

$$B = \text{Mid point of } QR = \left(\frac{-1+5}{2}, \frac{4+4}{2} \right) = (2, 4)$$

$$C = \text{Mid point of } RS = \left(\frac{5+5}{2}, \frac{4-1}{2} \right) = \left(5, \frac{3}{2} \right)$$

$$D = \text{Mid point of } PS = \left(\frac{-1+5}{2}, \frac{-1-1}{2} \right) = (2, -1)$$

$$\text{Slope of } AB = \frac{4 - \frac{3}{2}}{2 - (-1)} = \frac{5}{3}$$

$$\text{Slope of } BC = \frac{\frac{3}{2} - 4}{5 - 2} = \frac{-\frac{5}{2}}{3} = -\frac{5}{6}$$

$$\text{Slope of } CD = \frac{-1 - \frac{3}{2}}{2 - 5} = \frac{-\frac{5}{2}}{-3} = \frac{5}{6}$$

$$\text{Slope of } AD = \frac{-1 - \frac{3}{2}}{2 + 1} = -\frac{5}{3}$$

$\therefore AB \parallel CD, BC \parallel AD$

$$\begin{aligned} \text{Mid Point of } AC &= \left(\frac{-1+5}{2}, \frac{\frac{3}{2} + \frac{3}{2}}{2} \right) \\ &= \left(2, \frac{3}{2} \right) \end{aligned}$$

$$\begin{aligned} \text{Mid Point of } BD &= \left(\frac{2+2}{2}, \frac{4-1}{2} \right) \\ &= \left(2, \frac{3}{2} \right) \end{aligned}$$

\therefore diagonals bisect each other & opposite sides are parallel.

ABCD is a rhombus.

2. The area of a triangle is 5 sq.units. Two of its vertices are (2,1) and (3, -2). The third vertex is (x, y) where $y = x + 3$. Find the coordinates of the third vertex.

Solution :

and A (2, 1), B (3, -2), C (x, y) where $y = x + 3$

\therefore Area of Δ

$$= \frac{1}{2} \begin{vmatrix} 2 & 3 & x & 2 \\ 1 & -2 & y & 1 \end{vmatrix} = 5$$

$$\Rightarrow (-4 + 3y + x) - (3 - 2x + 2y) = 10$$

$$\Rightarrow x + 3y - 4 - 3 + 2x - 2y = 10$$

$$\Rightarrow 3x + y = 17 \quad \dots\dots (1)$$

$$\text{Also given, } x - y = -3 \quad \dots\dots (2)$$

$$\text{Adding, } 4x = 14$$

$$x = \frac{7}{2}$$

$$\text{Sub, } x = \frac{7}{2} \text{ in (2)}$$

$$\frac{7}{2} - y = -3$$

$$y = \frac{7}{2} + 3 = \frac{13}{2}$$

\therefore Third vertex is $\left(\frac{7}{2}, \frac{13}{2} \right)$

3. Find the area of a triangle formed by the lines $3x + y - 2 = 0$, $5x + 2y - 3 = 0$ and $2x - y - 3 = 0$.

Solution :

Given lines are

$$3x + y - 2 = 0 \quad \dots\dots (1)$$

$$5x + 2y - 3 = 0 \quad \dots\dots (2)$$

$$2x - y - 3 = 0 \quad \dots\dots (3)$$

Solving (1) & (2)

$$(1) \times 2 \Rightarrow 6x + 2y = 4$$

$$(2) \Rightarrow \frac{5x + 2y = 3}{x = 1}$$

Sub. in (1)

$$3 + y - 2 = 0$$

$$y = -1$$

$$\therefore A(1, -1)$$

Solving (1) & (2)

$$3x + y = 2$$

$$\frac{2x - y = 3}{5x = 5}$$

$$5x = 5$$

$$x = 1$$

$$\therefore y = -1$$

$$\therefore B \text{ is } (1, -1)$$

Solving (2) & (3)

$$(2) \Rightarrow 5x + 2y = 3$$

$$(3) \times 2 \Rightarrow \frac{4x - 2y = 6}{9x = 9}$$

$$x = 1$$

$$\therefore (3) \Rightarrow 2 - y - 3 = 0$$

$$\Rightarrow -y = 1$$

$$\therefore y = -1$$

$$\therefore C \text{ is } (1, -1)$$

$$\therefore A(1, -1), B(1, -1), C(1, -1)$$

\therefore All point line on the same line

\therefore Area of $\Delta = 0$ sq. units

4. If vertices of a quadrilateral are at $A(-5,7)$, $B(-4,k)$, $C(-1,-6)$ and $D(4,5)$ and its area is 72 sq.units. Find the value of k .

Solution :

Given vertices of quadrilateral are

$A(-5, 7)$, $B(-4, k)$, $C(-1, -6)$, $D(4, 5)$ & its area = 72 sq.units

$$\Rightarrow \frac{1}{2} \begin{bmatrix} -5 & -4 & -1 & 4 & -5 \\ 7 & k & -6 & 5 & 7 \end{bmatrix} = 72$$

$$(-5k + 24 - 5 + 28) - (-28 - k - 24 - 25) = 144$$

$$(-5k + 47) - (-k - 77) = 144$$

$$-4k + 124 = 144$$

$$-4k = 20$$

$$k = -5$$

5. Without using distance formula, show

$(-3, 2)$ are vertices of a parallelogram.

Solution :

Given vertices are

$$(-2, -1), (4, 0), (3, 3), (-3, 2)$$

Let $A(-2, -1)$, $B(4, 0)$, $C(3, 3)$, $D(-3, 2)$

$$\text{Slope of } AB = \frac{0 + 1}{4 + 2} = \frac{1}{6}$$

$$\text{Slope of } CD = \frac{3 - 2}{3 + 3} = \frac{1}{6}$$

$\therefore AB$ & CD are parallel

$$\text{Slope of } AD = \frac{2 + 1}{-3 + 2} = \frac{3}{-1} = -3$$

$$\text{Slope of } BC = \frac{0 - 3}{4 - 3} = \frac{-3}{1} = -3$$

$\therefore AD$ & BC are parallel

$\therefore ABCD$ is a parallelogram

6. Find the equations of the lines, whose sum and product of intercepts are 1 and -6 respectively.

Solution :

Given, sum of intercepts = 1

$$\Rightarrow a + b = 1$$

$$\therefore b = 1 - a$$

Given, product of intercepts = -6

$$\Rightarrow ab = -6$$

$$\therefore ab = -6 \Rightarrow a(1 - a) = -6$$

$$\Rightarrow a - a^2 = -6$$

$$\Rightarrow a^2 - a - 6 = 0$$

$$\Rightarrow (a - 3)(a + 2) = 0$$

$$\therefore a = 3, -2$$

If $a = 3, b = -2$

If $a = -2, b = 3$

$$a = 3, b = -2 \Rightarrow \frac{x}{3} + \frac{y}{-2} = 1$$

$$\Rightarrow \frac{x}{3} - \frac{y}{2} = 1$$

$$\Rightarrow 2x - 3y - 6 = 0$$

$$a = -2, b = 3 \Rightarrow \frac{x}{-2} + \frac{y}{3} = 1$$

$$\Rightarrow \frac{-3x + 2y}{6} = 1$$

$$\Rightarrow -3x + 2y = 6$$

$$\Rightarrow 3x - 2y + 6 = 0$$

7. The owner of a milk store finds that, he can sell 980 litres of milk each week at ₹14/litre and 1220 litres of milk each week at ₹16/litre. Assuming a linear relationship between selling price and demand, how many litres could he sell weekly at ₹17/litre?

Solution :

By data given,

the linear relationship between selling price per litre and demand is the equation of the line passing through the points

(14, 980) and (16, 1220) is

$$\therefore \frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$$

$$\Rightarrow \frac{y - 980}{1220 - 980} = \frac{x - 14}{16 - 14}$$

$$\Rightarrow \frac{y - 980}{240} = \frac{x - 14}{2}$$

$$\Rightarrow y - 980 = 120(x - 14)$$

$$\Rightarrow y = 120(x - 14) + 980 \quad \dots\dots\dots (1)$$

When $x = \text{Rs.}17 / \text{litre}$

$$y = 120(3) + 980$$

$$= 360 + 980$$


$$= 1340$$

\therefore He can sell weekly 1340 litres at Rs.17/litre

8. Find the image of the point (3,8) with respect to the line $x + 3y = 7$ assuming the line to be a plane mirror.

Solution :

To find the image of (3, 8) w.r.to the line $x + 3y = 7$



Let Q (h, k) be the image of P (3, 8) about the line $x + 3y = 7$ (1)

Since the line is assumed as a plane mirror P & Q are equidistant from R (x, y)

\therefore R is the midpoint and PQ is a perpendicular bisector of (1)

$$\therefore (x, y) = \left(\frac{h+3}{2}, \frac{k+8}{2} \right)$$

$$\therefore x = \frac{h+3}{2}, y = \frac{k+8}{2}$$

Since R (x, y) is a point on (1)

$$\left(\frac{h+3}{2} \right) + 3 \left(\frac{k+8}{2} \right) = 7$$

$$\Rightarrow h + 3 + 3k + 24 = 14$$

$$\Rightarrow h + 3k = -13 \quad \text{..... (2)}$$

Also, slope of PQ \times Slope of (1) = -1

$$\Rightarrow \frac{k-8}{h-3} \times \frac{-1}{3} = -1$$

$$\Rightarrow \frac{k-8}{h-3} = 3$$

$$\Rightarrow k - 8 = 3h - 9$$

$$\Rightarrow 3h - k = 1 \quad \text{..... (3)}$$

Solving (2) & (3)

$$(2) \Rightarrow h + 3k = -13$$

$$(3) \times 3 \Rightarrow \frac{9h - 3k = 3}{10h = -10}$$

$$h = -1$$

Sub in (2)

$$-1 + 3k = -13$$

$$3k = -12$$

$$k = -4$$

\therefore Q is (-1, -4), which is the image of P (3, 8)

9. Find the equation of a line passing through the point of intersection of the lines $4x + 7y - 3 = 0$ and $2x - 3y + 1 = 0$ that has equal intercepts on the axes.

Solution :

First, we find the point of intersection of the lines

$$4x + 7y = 3 \quad \text{..... (1)}$$

$$2x - 3y = -1 \quad \text{..... (2)}$$

$$(1) \Rightarrow 4x + 7y = +3$$

$$(2) \times 2 \Rightarrow 4x - 6y = -2$$

$$\frac{13y = 5}{y = \frac{5}{13}}$$

Sub in (1)

$$4x + \frac{35}{13} = 3$$

$$\Rightarrow x = -\frac{35}{13}$$

$$\Rightarrow 4x = \frac{4}{13}$$

$$\Rightarrow x = \frac{1}{13}$$

$$\therefore \text{The point is } \left(\frac{1}{13}, \frac{5}{13} \right)$$

Equation of plane in intercept form is

$$\frac{x}{a} + \frac{y}{b} = 1, \text{ where } a = b$$

$$\Rightarrow \frac{x}{a} + \frac{y}{b} = 1$$

$$\Rightarrow x + y = a \quad \text{..... (1)}$$

Since (1) passes through $\left(\frac{1}{13}, \frac{5}{13} \right)$


$$a = \frac{1}{13} + \frac{5}{13} = \frac{6}{13}$$

$$\therefore x + y = \frac{6}{13}$$

$$\Rightarrow \boxed{13x + 13y - 6 = 0}$$

10. A person standing at a junction (crossing) of two straight paths represented by the equations $2x - 3y + 4 = 0$ and $3x + 4y - 5 = 0$ seek to reach the path whose equation is $6x - 7y + 8 = 0$ in the least time. Find the equation of the path that he should follow.

Solution :



Given straight paths are

$$2x - 3y + 4 = 0 \quad \dots\dots (1) \quad (AB)$$

To reach the path

$$6x - 7y + 8 = 0 \quad \dots\dots (3) \quad (BC)$$

in the least time

To find : Equation of the path (AP)

A → Position of the person

Solving (1) & (2)

$$\begin{aligned} (1) \times 3 &\Rightarrow 6x - 9y = -12 \\ (2) \times 2 &\Rightarrow 6x + 8y = 10 \\ \hline &-17y = -22 \end{aligned}$$

$$\therefore y = \frac{22}{17}$$

Sub in (1)

$$\begin{aligned} 2x - \frac{66}{17} &= -4 \\ \Rightarrow 2x &= \frac{66}{17} - 4 \\ \Rightarrow 2x &= \frac{66 - 68}{17} \\ \Rightarrow 2x &= \frac{-2}{17} \\ \Rightarrow x &= \frac{-1}{17} \\ \therefore A \text{ is } &\left(\frac{-1}{17}, \frac{22}{17}\right) \end{aligned}$$

Also AP is perpendicular to BC, whose slope is $\frac{6}{7}$

$$\therefore \text{Slope of } AP = -\frac{7}{6}$$

\therefore Equation of the required path AP is

$$\begin{aligned} y - y_1 &= m(x - x_1) \\ \Rightarrow y - \frac{22}{17} &= -\frac{7}{6}\left(x + \frac{1}{17}\right) \\ \Rightarrow \frac{17y - 22}{17} &= -\frac{7}{6}\left(\frac{7x + 1}{17}\right) \\ \Rightarrow \frac{17y - 22}{17} &= \frac{-(7x + 1)}{6} \end{aligned}$$

$$\Rightarrow 6(17y - 22) = -17(7x + 1)$$

$$\Rightarrow 102y - 132 = -119x - 17$$

$$\Rightarrow 119x + 102y - 125 = 0$$

is the required path.

PROBLEMS FOR PRACTICE

- If $P\left(\frac{a}{2}, 4\right)$ is the mid point of the line joining the points A $(-6, 5)$, B $(-2, 3)$, then find 'a'. **(Ans : a = -8)**
- Find the area of ΔABC whose vertices are
 - A $(3, 8)$, B $(-4, 2)$, C $(5, -1)$ **(Ans : 37.5)**
 - A $(1, 2)$, B $(-3, 4)$, C $(-5, -6)$ **(Ans : 22)**
 - A $(0, 1)$, B $(2, 3)$, C $(3, 4)$ **(Ans : 0)**
- If the area of Δ is 12 sq. units with vertices $(a, -3)$, $(3, a)$, $(-1, 5)$ find 'a'. **(Ans : 1, 3)**
- If the area of Δ formed by (x, y) , $(1, 2)$, $(2, 1)$ is 6 sq. units, prove that $x + y = 15$.
- For what values of k, are the points $(8, 1)$, $(3, -2k)$, and $(k, -5)$ are collinear. **(Ans : k = 2, $\frac{11}{2}$)**

6. Find the value of 'p' if the area of Δ formed by $(p + 1, 2p - 2)$, $(p - 1, p)$ and $(p - 3, 2p - 6)$ is 0. **(Ans : p = 4)**
7. Find the area of quadrilateral whose vertices are,
 i) A (3, -1), B (9, -5), C (14, 0), D (9, 19) **(Ans : 132)**
 ii) P (-5, -3), Q (-4, -6), R (2, -3), S (1, 2) **(Ans : 28)**
 iii) E (-3, 2), F (5, 4), G (7, -6), H (-5, -4) **(Ans : 80)**
 iv) A (-4, 5), B (0, 7), C (5, -5), D (-4, -2) **(Ans : 60.5)**
8. If (3, 3), (6, y), (x, 7) and (5, 6) are the vertices of a parallelogram taken in order, find x and y. **(Ans : x = 8, y = 4)**
9. If the points (p, q), (m, n), $(p - m, q - n)$ are collinear, show that $pn = qm$.
10. Three vertices of a parallelogram ABCD are (1, 2), (4, 3), (6, 6). Find the 4th vertex D. **(Ans : 3, 5)**
11. The line joining A (0, 5) and B (4, 2) is perpendicular to the line joining C (-1, -2), D (5, b) find 'b'. **(Ans : b = 6)**
12. Find the equation of the line passing through (9, -1) having its x-intercept thrice as its y-intercept. **(Ans : x + 3y - 6 = 0)**
13. Find the slope and y-intercept of the line $10x + 15y + 6 = 0$. **(Ans : $m = -\frac{2}{5}$, $c = -\frac{2}{5}$)**
14. Find whether the lines drawn through the two pair of points are parallel (or) perpendicular.
 i) (5, 2), (0, 5) and (0, 0), (-5, 3) **(Ans : parallel)**
 ii) (4, 5), (0, -2) and (-5, 1), (2, -3) **(Ans : perpendicular)**
15. A line passing through the points (2, 7) and (3, 6) is parallel to the line joining (9, a) and (11, 2) find 'a'. **(Ans : -2 (or) 4)**
16. Find the equation of a straight line whose slope is $\frac{2}{3}$ and passing through the point (5, -4) **(Ans : 2x - 3y - 22 = 0)**
17. Without using distance formula, show that the points P (3, 2), Q (0, -3), R (-3, -2) and S (0, 1) are the vertices of a parallelogram.
18. A triangle has vertices at (3, 4), (1, 2), (-5, -6). Find the slopes of the medians. **(Ans : $\frac{6}{5}$, $\frac{3}{2}$, $\frac{9}{7}$)**
19. Find the equation of altitude from A of a Δ ABC whose vertices are (1, -3), (-2, 5), (-3, 4). **(Ans : x + y + 2 = 0)**
20. Find the values of 'p' of the straight lines are perpendicular to each other. **(Ans : p = 1, 2)**
21. Find the equation of the straight line passing through (1, 4) and having intercepts in the ratio 3 : 5 **(Ans : 5x + 3y = 17)**
22. Find the area of the triangle formed by sides $x + 4y - 9 = 0$, $9x + 10y + 23 = 0$, $7x + 2y - 11 = 0$ **(Ans : 26 sq.units)**
23. Find the equation of the line through the point of intersection of the lines $2x + y - 5 = 0$, $x + y - 3 = 0$ and bisecting the line segment joining the points (3, -2) (-5, 6) **(Ans : x + 3y - 5 = 0)**
24. Find the image of the point (-2, 3) w.r.to the line $x + 2y - 9 = 0$ **(Ans : 0, 7)**
25. The equation of the diagonals of a rectangle are $4x - 7y = 0$, $8x - y = 26$ and one of its sides is $2x + 3y = 0$, find the equation of the other sides. **(Ans : 2x+3y-26=0, 3x-2y-13=0, 3x-2y = 0)**

OBJECTIVE TYPE QUESTIONS

1. The point whose the line $3x - y + 6 = 0$ meets the x - axis is

a) (0, 6) b) (-2, 0)
c) (-1, 3) d) (2, 0) **Ans : (b)**

2. The point of intersection of the lines $2x + y - 3 = 0$, $5x + y - 6 = 0$ lies in the quadrant

a) I b) II c) IV d) III

Ans : (a)

3. The value of 'k' if the lines $3x + 6y + 7 = 0$ and $2x + ky - 5 = 0$ are perpendicular is

a) 1 b) -1 c) 2 d) $\frac{1}{2}$

Ans : (b)

4. The slope of the line which is parallel to the line joining the points (0, 0) and (-5, 5) is

a) 1 b) -1 c) 2 d) -2

Ans : (a)

5. The area of triangle formed by (0, 4) (4, 0) and origin is

a) 8 b) 16 c) 2 d) 4 **Ans : (a)**

6. The equation of a straight line which has the y-intercept 5 and slope 2 is

a) $2x + y + 5 = 0$ b) $2x - y + 5 = 0$
c) $2x - y - 5 = 0$ d) $2x + y - 5 = 0$

Ans : (b)

7. If the point (a, a) lies on the line $3x + 4y - 14 = 0$ then 'a' is

a) 2 b) -2 c) 1 d) 0 **Ans : (a)**

8. Equation of the line parallel to y-axis and passing through (-2, 3) is

a) $x = 3$ b) $y = -2$
c) $x = -2$ d) $y = 3$

Ans (c)

9. The x-intercept of the line $3x - 2y + 12 = 0$ is

a) 6 b) -6 c) 4 d) -4

Ans : (d)

10. AB is parallel to CD. If A and B are (2, 3) and (6, 9) the slope of CD is

a) $\frac{4}{9}$ b) $\frac{3}{2}$ c) $\frac{2}{3}$ d) $\frac{9}{4}$

Ans : (b)

11. If (1, 2), (4, 6), (x, 6) and (3, 2) are the vertices of a parallelogram taken in order, then x is

a) 6 b) 2 c) 1 d) 3

Ans : (a)

12. If the slope of a line is $-\sqrt{3}$, then the angle of inclination is

a) 60 b) 30 c) 120 d) 150

Ans : (c)

13. The value of a for which (-a, a) is collinear with the points (2, 0), (0, 1) is

a) 1 b) 2 c) -2 d) -1

Ans : (b)

14. The x - coordinates of the point of intersection of the lines $x - 7y + 5 = 0$, $3x + y = 0$ is

a) $\frac{15}{22}$ b) $\frac{5}{22}$ c) $\frac{-5}{22}$ d) $\frac{-10}{22}$

Ans : (c)

15. The equations of the 4 sides of a rectangle are $x = 1$, $y = 2$, $x = 4$, $y = 5$. One vertex of the rectangle is at

a) (2, 4) b) (5, 1)
c) (2, 5) d) (4, 2)


Ans : (d)

CHAPTER 6

TRIGONOMETRY

TRIGONOMETRIC RATIOS

Key Points



$$\sin \theta = \frac{\text{Opposite side}}{\text{Hypotenuse}} = \frac{MP}{OP}$$

$$\cos \theta = \frac{\text{Adjacent side}}{\text{Hypotenuse}} = \frac{OM}{OP}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}; \cot \theta = \frac{\cos \theta}{\sin \theta};$$

$$\operatorname{cosec} \theta = \frac{1}{\sin \theta}; \sec \theta = \frac{1}{\cos \theta}$$

$$\sin (90^\circ - \theta) = \cos \theta \qquad \cos (90^\circ - \theta) = \sin \theta \qquad \tan (90^\circ - \theta) = \cot \theta$$

$$\operatorname{cosec} (90^\circ - \theta) = \sec \theta \qquad \sec (90^\circ - \theta) = \operatorname{cosec} \theta \qquad \cot (90^\circ - \theta) = \tan \theta$$

Trigonometric Ratio \ θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	undefined
$\operatorname{cosec} \theta$	undefined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
$\sec \theta$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	undefined
$\cot \theta$	undefined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

I. TRIGONOMETRIC IDENTITIES**Key Points**

✓ $\sin^2\theta + \cos^2\theta = 1 \Rightarrow$	$\sin^2\theta = 1 - \cos^2\theta$ (or) $\cos^2\theta = 1 - \sin^2\theta$
✓ $1 + \tan^2\theta = \sec^2\theta \Rightarrow$	$\tan^2\theta = \sec^2\theta - 1$ (or) $\sec^2\theta - \tan^2\theta = 1$
✓ $1 + \cot^2\theta = \operatorname{cosec}^2\theta \Rightarrow$	$\cot^2\theta = \operatorname{cosec}^2\theta - 1$ (or) $\operatorname{cosec}^2\theta - \cot^2\theta = 1$

Example 6.1

Prove that $\tan^2\theta - \sin^2\theta = \tan^2\theta \sin^2\theta$

Solution :

$$\begin{aligned}\tan^2\theta - \sin^2\theta &= \tan^2\theta - \frac{\sin^2\theta}{\cos^2\theta} \cdot \cos^2\theta \\ &= \tan^2\theta (1 - \cos^2\theta) = \tan^2\theta \sin^2\theta\end{aligned}$$

Example 6.2

Prove that $\frac{\sin A}{1 + \cos A} = \frac{1 - \cos A}{1 - \cos A}$

Solution :

$$\frac{\sin A}{1 + \cos A} = \frac{\sin A}{1 + \cos A} \times \frac{1 - \cos A}{1 - \cos A}$$

[multiply numerator and denominator by the conjugate of $1 + \cos A$]

$$\begin{aligned}&= \frac{\sin A (1 - \cos A)}{(1 + \cos A)(1 - \cos A)} = \frac{\sin A (1 - \cos A)}{1 - \cos^2 A} \\ &= \frac{\sin A (1 - \cos A)}{\sin^2 A} = \frac{1 - \cos A}{\sin A}\end{aligned}$$

Example 6.3

Prove that $1 + \frac{\cot^2\theta}{1 + \operatorname{cosec}\theta} = \operatorname{cosec}\theta$

Solution :

$$\begin{aligned}&= 1 + \frac{\cot^2\theta}{1 + \operatorname{cosec}\theta} \\ &= 1 + \frac{\operatorname{cosec}^2\theta - 1}{\operatorname{cosec}\theta + 1} \quad \left[\text{since } \operatorname{cosec}^2\theta - 1 = \cot^2\theta \right] \\ &= 1 + \frac{(\operatorname{cosec}\theta + 1)(\operatorname{cosec}\theta - 1)}{\operatorname{cosec}\theta + 1} \\ &= 1 + (\operatorname{cosec}\theta - 1) = \operatorname{cosec}\theta\end{aligned}$$

Example 6.4

Prove that $\sec\theta - \cos\theta = \tan\theta \sin\theta$

Solution :

$$\begin{aligned}\sec\theta - \cos\theta &= \frac{1}{\cos\theta} - \cos\theta = \frac{1 - \cos^2\theta}{\cos\theta} \\ &= \frac{\sin^2\theta}{\cos\theta} \quad \left[\text{since } 1 - \cos^2\theta = \sin^2\theta \right] \\ &= \frac{\sin\theta}{\cos\theta} \times \sin\theta = \tan\theta \sin\theta\end{aligned}$$

Example 6.5

Prove that $\sqrt{\frac{1 + \cos\theta}{1 - \cos\theta}} = \operatorname{cosec}\theta + \cot\theta$

Solution :

$$\sqrt{\frac{1 + \cos\theta}{1 - \cos\theta}} = \sqrt{\frac{1 + \cos\theta}{1 - \cos\theta}} \times \frac{1 + \cos\theta}{1 + \cos\theta}$$

[multiply numerator and denominator by the conjugate of $1 - \cos\theta$]

$$\begin{aligned}&= \sqrt{\frac{(1 + \cos\theta)^2}{1 - \cos^2\theta}} = \frac{1 + \cos\theta}{\sqrt{\sin^2\theta}} \\ &\quad \left[\text{since } \sin^2\theta + \cos^2\theta = 1 \right] \\ &= \frac{1 + \cos\theta}{\sin\theta} = \operatorname{cosec}\theta + \cot\theta\end{aligned}$$

Example 6.6

Prove that $\frac{\sec\theta}{\sin\theta} - \frac{\sin\theta}{\cos\theta} = \cot\theta$

Solution :

$$\begin{aligned}
 &= \frac{\sec \theta}{\sin \theta} - \frac{\sin \theta}{\cos \theta} \\
 &= \frac{1}{\cos \theta} - \frac{\sin \theta}{\cos \theta} \\
 &= \frac{1}{\sin \theta \cos \theta} - \frac{\sin \theta}{\cos \theta} \\
 &= \frac{1 - \sin^2 \theta}{\sin \theta \cos \theta} = \frac{\cos^2 \theta}{\sin \theta \cos \theta} \\
 &= \cot \theta
 \end{aligned}$$

Example 6.7

Prove that $\sin^2 A \cos^2 B + \cos^2 A \sin^2 B + \cos^2 A \cos^2 B + \sin^2 A \sin^2 B = 1$

Solution :

$$\begin{aligned}
 &\sin^2 A \cos^2 B + \sin^2 A \sin^2 B + \cos^2 A \sin^2 B + \cos^2 A \cos^2 B \\
 &= \sin^2 A (\cos^2 B + \sin^2 B) + \cos^2 A (\sin^2 B + \cos^2 B) \\
 &= \sin^2 A (1) + \cos^2 A (1) \\
 &\quad (\text{since } \sin^2 B + \cos^2 B = 1) \\
 &= \sin^2 A + \cos^2 A = 1
 \end{aligned}$$

Example 6.8

If $\cos \theta + \sin \theta = \sqrt{2} \cos \theta$, then prove that $\cos \theta - \sin \theta = \sqrt{2} \sin \theta$

Solution :

$$\begin{aligned}
 &\text{Now, } \cos \theta + \sin \theta = \sqrt{2} \cos \theta \\
 &\text{Squaring both sides,} \\
 &(\cos \theta + \sin \theta)^2 = (\sqrt{2} \cos \theta)^2 \\
 &\cos^2 \theta + \sin^2 \theta + 2 \sin \theta \cos \theta = 2 \cos^2 \theta
 \end{aligned}$$

$$2 \cos^2 \theta - \cos^2 \theta - \sin^2 \theta = 2 \sin \theta \cos \theta$$

$$\cos^2 \theta - \sin^2 \theta = 2 \sin \theta \cos \theta$$

$$\begin{aligned}
 &(\cos \theta + \sin \theta) (\cos \theta - \sin \theta) \\
 &\quad = 2 \sin \theta \cos \theta
 \end{aligned}$$

$$\begin{aligned}
 \cos \theta - \sin \theta &= \frac{2 \sin \theta \cos \theta}{\cos \theta + \sin \theta} = \frac{2 \sin \theta \cos \theta}{\sqrt{2} \cos \theta} \\
 &= \sqrt{2} \sin \theta
 \end{aligned}$$

$$[\text{since } \cos \theta + \sin \theta = \sqrt{2} \cos \theta]$$

$$\text{Therefore, } \cos \theta - \sin \theta = \sqrt{2} \sin \theta$$

Example 6.9

Prove that $(\operatorname{cosec} \theta - \sin \theta) (\sec \theta - \cos \theta) (\tan \theta + \cot \theta) = 1$

Solution :

$$\begin{aligned}
 &= \left(\frac{1}{\sin \theta} - \sin \theta \right) \left(\frac{1}{\cos \theta} - \cos \theta \right) \left(\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} \right) \\
 &= \frac{1 - \sin^2 \theta}{\sin \theta} \times \frac{1 - \cos^2 \theta}{\cos \theta} \times \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta} \\
 &= \frac{\cos^2 \theta \sin^2 \theta \times 1}{\sin^2 \theta \cos^2 \theta} = 1
 \end{aligned}$$

Example 6.10

Prove that $\frac{\sin A}{1 + \cos A} + \frac{\sin A}{1 - \cos A} = 2 \operatorname{cosec} A$

Solution :

$$\begin{aligned}
 &= \frac{\sin A}{1 + \cos A} + \frac{\sin A}{1 - \cos A} \\
 &= \frac{\sin A (1 - \cos A) + \sin A (1 + \cos A)}{(1 + \cos A) (1 - \cos A)} \\
 &= \frac{\sin A - \sin A \cos A + \sin A + \sin A \cos A}{1 - \cos^2 A} \\
 &= \frac{2 \sin A}{1 - \cos^2 A} = \frac{2 \sin A}{\sin^2 A} = 2 \operatorname{cosec} A
 \end{aligned}$$

Example 6.11

If $\operatorname{cosec} \theta + \cot \theta = P$, then prove that

$$\cos \theta = \frac{P^2 - 1}{P^2 + 1}$$

Solution :

$$\text{Given } \operatorname{cosec} \theta + \cot \theta = P \quad \dots (1)$$

$$\operatorname{cosec}^2 \theta - \cot^2 \theta = 1 \text{ (identity)}$$

$$\operatorname{cosec} \theta - \cot \theta = \frac{1}{\operatorname{cosec} \theta + \cot \theta}$$

$$\operatorname{cosec} \theta - \cot \theta = \frac{1}{P} \quad \dots (2)$$

Adding (1) and (2) we get,

$$2 \operatorname{cosec} \theta = P + \frac{1}{P}$$

$$2 \operatorname{cosec} \theta = \frac{P^2 + 1}{P} \quad \dots (3)$$

Subtracting (2) from (1), we get,

$$2 \cot \theta = P - \frac{1}{P}$$

$$2 \cot \theta = \frac{P^2 - 1}{P} \quad \dots (4)$$

Dividing (4) by (3) we get,

$$\frac{2 \cot \theta}{2 \operatorname{cosec} \theta} = \frac{P^2 - 1}{P} \times \frac{P}{P^2 + 1}$$

$$\text{gives, } \cos \theta = \frac{P^2 - 1}{P^2 + 1}$$

Example 6.12

Prove that $\tan^2 A - \tan^2 B = \frac{\sin^2 A - \sin^2 B}{\cos^2 A \cos^2 B}$

Solution :

$$\tan^2 A - \tan^2 B$$

$$= \frac{\sin^2 A}{\cos^2 A} - \frac{\sin^2 B}{\cos^2 B}$$

$$= \frac{\sin^2 A \cos^2 B - \sin^2 B \cos^2 A}{\cos^2 A \cos^2 B}$$

$$= \frac{\sin^2 A (1 - \sin^2 B) - \sin^2 B (1 - \sin^2 A)}{\cos^2 A \cos^2 B}$$

$$= \frac{\sin^2 A - \sin^2 A \sin^2 B - \sin^2 B + \sin^2 A \sin^2 B}{\cos^2 A \cos^2 B}$$

$$= \frac{\sin^2 A - \sin^2 B}{\cos^2 A \cos^2 B}$$

Example 6.13

Prove that

$$\left| \frac{\cos^3 A - \sin^3 A}{\cos A - \sin A} \right| - \left| \frac{\cos^3 A + \sin^3 A}{\cos A + \sin A} \right| = 2 \sin A \cos A$$

Solution :

$$= \left(\frac{\cos^3 A - \sin^3 A}{\cos A - \sin A} \right) - \left(\frac{\cos^3 A + \sin^3 A}{\cos A + \sin A} \right)$$

$$= \left(\frac{(\cos A - \sin A)(\cos^2 A + \sin^2 A + \cos A \sin A)}{\cos A - \sin A} \right) -$$

$$= \left(\frac{(\cos A + \sin A)(\cos^2 A + \sin^2 A - \cos A \sin A)}{\cos A + \sin A} \right)$$

$$\left[\begin{array}{l} \text{since } a^3 - b^3 = (a - b)(a^2 + b^2 + ab) \\ a^3 + b^3 = (a + b)(a^2 + b^2 - ab) \end{array} \right]$$

$$= (1 + \cos A \sin A) - (1 - \cos A \sin A)$$

$$= 2 \cos A \sin A$$

Example 6.14

Prove that

$$\frac{\sin A}{\sec A + \tan A - 1} + \frac{\cos A}{\operatorname{cosec} A + \cot A - 1} = 1$$

Solution :

$$\begin{aligned}
&= \frac{\sin A}{\sec A + \tan A - 1} + \frac{\cos A}{\operatorname{cosec} A + \cot A - 1} \\
&= \frac{\sin A (\operatorname{cosec} A + \cot A - 1) + \cos A (\sec A + \tan A - 1)}{(\sec A + \tan A - 1)(\operatorname{cosec} A + \cot A - 1)} \\
&= \frac{\sin A \operatorname{cosec} A + \sin A \cot A - \sin A + \cos A \sec A + \cos A \tan A - \cos A}{(\sec A + \tan A - 1)(\operatorname{cosec} A + \cot A - 1)} \\
&= \frac{1 + \cos A - \sin A + 1 + \sin A - \cos A}{\left(\frac{1}{\cos A} + \frac{\sin A}{\cos A} - 1\right)\left(\frac{1}{\sin A} + \frac{\cos A}{\sin A} - 1\right)} \\
&= \frac{2}{\left(\frac{1 + \sin A - \cos A}{\cos A}\right)\left(\frac{1 + \cos A - \sin A}{\sin A}\right)} \\
&= \frac{2 \sin A \cos A}{(1 + \sin A - \cos A)(1 + \cos A - \sin A)} \\
&= \frac{2 \sin A \cos A}{[(1 + \sin A - \cos A)][(1 - \sin A - \cos A)]} \\
&= \frac{2 \sin A \cos A}{1 - (1 - 2 \sin A \cos A)} \\
&= \frac{2 \sin A \cos A}{1 - 1 + 2 \sin A \cos A} = \frac{2 \sin A \cos A}{2 \sin A \cos A} = 1
\end{aligned}$$

Example 6.15

Show that $\left(\frac{1 + \tan^2 A}{1 + \cot^2 A}\right) = \left(\frac{1 - \tan A}{1 - \cot A}\right)^2$

Solution :

LHS

$$\begin{aligned}
\left(\frac{1 + \tan^2 A}{1 + \cot^2 A}\right) &= \frac{1 + \tan^2 A}{1 + \frac{1}{\tan^2 A}} \\
&= \frac{1 + \tan^2 A}{\frac{\tan^2 A + 1}{\tan^2 A}} = \tan^2 A \quad \dots\dots (1)
\end{aligned}$$

RHS

$$\begin{aligned}
\left(\frac{1 - \tan A}{1 - \cot A}\right)^2 &= \left(\frac{1 - \tan A}{1 - \frac{1}{\tan A}}\right)^2 \\
&= \left(\frac{1 - \tan A}{\frac{\tan A - 1}{\tan A}}\right)^2 = (-\tan A)^2 = \tan^2 A
\end{aligned}$$

From (1) and (2),

$$\left(\frac{1 + \tan^2 A}{1 + \cot^2 A}\right) = \left(\frac{1 - \tan A}{1 - \cot A}\right)^2$$

Example 6.16

Prove that

$$\frac{(1 + \cot A + \tan A)(\sin A - \cos A)}{\sec^3 A - \operatorname{cosec}^3 A} = \sin^2 A \cos^2 A$$

Solution :

$$\begin{aligned}
&\frac{(1 + \cot A + \tan A)(\sin A - \cos A)}{\sec^3 A - \operatorname{cosec}^3 A} \\
&= \frac{\left(1 + \frac{\cos A}{\sin A} + \frac{\sin A}{\cos A}\right)(\sin A - \cos A)}{(\sec A - \operatorname{cosec} A)(\sec^2 A + \sec A \operatorname{cosec} A + \operatorname{cosec}^2 A)} \\
&= \frac{(\sin A \cos A + \cos^2 A + \sin^2 A)(\sin A - \cos A)}{\sin A \cos A} \\
&= \frac{(\sec A - \operatorname{cosec} A)\left(\frac{1}{\cos^2 A} + \frac{1}{\cos A \sin A} + \frac{1}{\sin^2 A}\right)}{(\sin A \cos A + 1)\left(\frac{\sin A}{\sin A \cos A} - \frac{\cos A}{\sin A \cos A}\right)} \\
&= \frac{(\sec A - \operatorname{cosec} A)\left(\frac{\sin^2 A + \sin A \cos A + \cos^2 A}{\sin^2 A \cos^2 A}\right)}{(\sec A - \operatorname{cosec} A)(1 + \sin A \cos A)} \\
&= \frac{(\sin A \cos A + 1)(\sec A - \operatorname{cosec} A)}{(\sec A - \operatorname{cosec} A)(1 + \sin A \cos A)} \times \sin^2 A \cos^2 A \\
&= \sin^2 A \cos^2 A
\end{aligned}$$

Example 6.17

If $\frac{\cos^2 \theta}{\sin \theta} = p$ and $\frac{\sin^2 \theta}{\cos \theta} = q$, then prove that

$$p^2 q^2 (p^2 + q^2 + 3) = 1$$

Solution :

We have $\frac{\cos^2 \theta}{\sin \theta} = p$ (1) and $\frac{\sin^2 \theta}{\cos \theta} = q$ (2)

$$p^2 q^2 (p^2 + q^2 + 3) =$$

$$\left(\frac{\cos^2 \theta}{\sin \theta}\right)^2 \left(\frac{\sin^2 \theta}{\cos \theta}\right)^2 \times \left[\left(\frac{\cos^2 \theta}{\sin \theta}\right)^2 + \left(\frac{\sin^2 \theta}{\cos \theta}\right)^2 + 3\right]$$

[from (1) and (2)]

$$= \left(\frac{\cos^4 \theta}{\sin^2 \theta}\right) \left(\frac{\sin^4 \theta}{\cos^2 \theta}\right) \times \left[\left(\frac{\cos^4 \theta}{\sin^2 \theta}\right) + \left(\frac{\sin^4 \theta}{\cos^2 \theta}\right) + 3\right]$$

$$= (\cos^2 \theta \times \sin^2 \theta) \times$$

$$\left[\left(\frac{\cos^6 \theta + \sin^6 \theta + 3 \sin^2 \theta \cos^2 \theta}{\sin^2 \theta \cos^2 \theta}\right)\right]$$

$$= \cos^6 \theta + \sin^6 \theta + 3 \sin^2 \theta \cos^2 \theta$$

$$= (\cos^2 \theta)^3 \times (\sin^2 \theta)^3 + 3 \sin^2 \theta \cos^2 \theta$$

$$= [(\cos^2 \theta \sin^2 \theta)^3 - 3 \cos^2 \theta \sin^2 \theta (\cos^2 \theta \sin^2 \theta)]$$

$$+ 3 \sin^2 \theta \cos^2 \theta$$

$$= 1 - 3 \cos^2 \theta \sin^2 \theta (1) + 3 \cos^2 \theta \sin^2 \theta = 1$$

EXERCISE 6.1

1. Prove the following identities.

i) $\cot \theta + \tan \theta = \sec \theta \operatorname{cosec} \theta$

ii) $\tan^4 \theta + \tan^2 \theta = \sec^4 \theta - \sec^2 \theta$

Solution:

i) LHS

$$= \cot \theta + \tan \theta$$

$$= \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{\cos \theta}$$

$$= \frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta \cdot \cos \theta}$$

$$= \frac{1}{\sin \theta \cdot \cos \theta}$$

$$= \frac{1}{\cos \theta} \times \frac{1}{\sin \theta}$$

$$= \sec \theta \cdot \operatorname{cosec} \theta$$

$$= \text{RHS}$$

ii) RHS

$$= \tan^4 \theta + \tan^2 \theta$$

$$= \tan^2 \theta (\tan^2 \theta + 1)$$

$$= (\sec^2 \theta - 1) \cdot (\sec^2 \theta)$$

$$= \sec^4 \theta - \sec^2 \theta$$

$$= \text{RHS}$$

2. Prove the following identities.

i) $\frac{1 - \tan^2 \theta}{\cot^2 \theta - 1} = \tan^2 \theta$

ii) $\frac{\cos \theta}{1 + \sin \theta} = \sec \theta - \tan \theta$

Solution :

i) LHS

$$= \frac{1 - \tan^2 \theta}{\cot^2 \theta - 1}$$

$$= \frac{1 - \tan^2 \theta}{\frac{1}{\tan^2 \theta} - 1}$$

$$= \frac{1 - \tan^2 \theta}{\frac{1 - \tan^2 \theta}{\tan^2 \theta}}$$

$$= \tan^2 \theta$$

$$= \text{RHS}$$

ii) LHS

$$= \frac{\cos \theta}{1 + \sin \theta}$$

$$= \frac{\cos \theta}{1 + \sin \theta} \times \frac{1 - \sin \theta}{1 - \sin \theta}$$

$$= \frac{\cos \theta (1 - \sin \theta)}{\cos^2 \theta}$$

$$= \frac{1 - \sin \theta}{\cos \theta}$$

$$= \frac{1}{\cos \theta} - \frac{\sin \theta}{\cos \theta}$$

$$= \sec \theta - \tan \theta$$

$$= \text{RHS}$$

3. Prove the following identities.

$$i) \sqrt{\frac{1+\sin\theta}{1-\sin\theta}} = \sec\theta + \tan\theta$$

$$ii) \sqrt{\frac{1+\sin\theta}{1-\sin\theta}} + \sqrt{\frac{1-\sin\theta}{1+\sin\theta}} = 2\sec\theta$$

Solution :

i) LHS

$$\begin{aligned} &= \sqrt{\frac{1+\sin\theta}{1-\sin\theta}} \\ &= \sqrt{\frac{1+\sin\theta}{1-\sin\theta} \times \frac{1+\sin\theta}{1+\sin\theta}} \\ &= \sqrt{\frac{(1+\sin\theta)^2}{\cos^2\theta}} \\ &= \frac{1+\sin\theta}{\cos\theta} \\ &= \frac{1}{\cos\theta} + \frac{\sin\theta}{\cos\theta} \\ &= \sec\theta + \tan\theta \\ &= \text{RHS} \end{aligned}$$

ii) LHS

$$\begin{aligned} &= \sqrt{\frac{1+\sin\theta}{1-\sin\theta}} + \sqrt{\frac{1-\sin\theta}{1+\sin\theta}} \\ &= (\sec\theta + \tan\theta) + \frac{1}{\sec\theta + \tan\theta} \\ &= (\sec\theta + \tan\theta) + (\sec\theta - \tan\theta) \\ &= 2\sec\theta \\ &= \text{RHS} \end{aligned}$$

4. Prove the following identities.

$$i) \sec^6\theta = \tan^6\theta + 3\tan^2\theta \sec^2\theta + 1$$

$$ii) (\sin\theta + \sec\theta)^2 + (\cos\theta + \operatorname{cosec}\theta)^2 = 1 + (\sec\theta + \operatorname{cosec}\theta)^2$$

Solution :

i) LHS

$$= \sec^6\theta$$

$$= (\sec^2\theta)^3$$

$$= (1 + \tan^2\theta)^3$$

$$(\because a+b)^3 = a^3 + b^3 + 3ab(a+b)$$

$$= 1 + \tan^6\theta + 3(1)\tan^2\theta(1 + \tan^2\theta)$$

$$= 1 + \tan^6\theta + 3\tan^2\theta \cdot \sec^2\theta$$

$$= \text{RHS}$$

ii) LHS

$$= (\sin\theta + \sec\theta)^2 + (\cos\theta + \operatorname{cosec}\theta)^2$$

$$= \sin^2\theta + \sec^2\theta + 2\sin\theta \cdot \sec\theta + \cos^2\theta + \operatorname{cosec}^2\theta + 2\cos\theta \cdot \operatorname{cosec}\theta$$

$$= (\sin^2\theta + \cos^2\theta) + \sec^2\theta +$$

$$\frac{2\sin\theta}{\cos\theta} + \operatorname{cosec}^2\theta + \frac{2\cos\theta}{\sin\theta}$$

$$= 1 + \sec\theta + \operatorname{cosec}\theta + 2\left(\frac{\sin\theta}{\cos\theta} + \frac{\cos\theta}{\sin\theta}\right)$$

$$= 1 + \sec^2\theta + \operatorname{cosec}^2\theta + 2\left(\frac{\sin^2\theta + \cos^2\theta}{\cos\theta\sin\theta}\right)$$

$$= 1 + \sec^2\theta + \operatorname{cosec}^2\theta + 2\sec\theta \operatorname{cosec}\theta$$

$$= 1 + (\sec\theta \operatorname{cosec}\theta)^2$$

$$= \text{RHS}$$

5. Prove the following identities.

$$i) \sec^4\theta(1 - \sin^4\theta) - 2\tan^2\theta = 1$$

$$ii) \frac{\cot\theta - \cos\theta}{\cot\theta + \cos\theta} = \frac{\operatorname{cosec}\theta - 1}{\operatorname{cosec}\theta + 1}$$

Solution :

i) LHS

$$= \sec^4\theta(1 - \sin^4\theta) - 2\tan^2\theta$$

$$\begin{aligned}
 &= \frac{1}{\cos^4 \theta} (1 + \sin^2 \theta) \cdot (1 - \sin^2 \theta) - \frac{2 \sin^2 \theta}{\cos^2 \theta} \\
 &= \frac{(1 + \sin^2 \theta) \cos^2 \theta}{\cos^4 \theta} - \frac{2 \sin^2 \theta}{\cos^2 \theta} \\
 &= \frac{1 + \sin^2 \theta}{\cos^2 \theta} - \frac{2 \sin^2 \theta}{\cos^2 \theta} \\
 &= \frac{1 - \sin^2 \theta}{\cos^2 \theta} \\
 &= \frac{\cos^2 \theta}{\cos^2 \theta} \\
 &= 1 \\
 &= \text{RHS}
 \end{aligned}$$

ii) LHS

$$\begin{aligned}
 &= \frac{\cot \theta - \cos \theta}{\cot \theta + \cos \theta} \\
 &= \frac{\frac{\cos \theta}{\sin \theta} - \cos \theta}{\frac{\cos \theta}{\sin \theta} + 1} \\
 &= \frac{\cos \theta \left(\frac{1}{\sin \theta} - 1 \right)}{\cos \theta \left(\frac{1}{\sin \theta} + 1 \right)} \\
 &= \frac{\operatorname{cosec} \theta - 1}{\operatorname{cosec} \theta + 1} \\
 &= \text{RHS}
 \end{aligned}$$

6. Prove the following identities.

$$\begin{aligned}
 \text{i)} \quad & \frac{\sin A - \sin B}{\cos A + \cos B} + \frac{\cos A - \cos B}{\sin A + \sin B} = 0 \\
 \text{ii)} \quad & \frac{\sin^3 A + \cos^3 A}{\sin A + \cos A} + \frac{\sin^3 A - \cos^3 A}{\sin A - \cos A} = 2
 \end{aligned}$$

Solution :

i) LHS

$$\begin{aligned}
 &= \frac{\sin A - \sin B}{\cos A + \cos B} + \frac{\cos A - \cos B}{\sin A + \sin B} \\
 &= \frac{(\sin^2 A - \sin^2 B) + (\cos^2 A - \cos^2 B)}{(\cos A + \cos B) \cdot (\sin A + \sin B)} \\
 &= \frac{(\sin^2 A + \cos^2 A) - (\sin^2 B + \cos^2 B)}{(\cos A + \cos B) \cdot (\sin A + \sin B)} \\
 &= \frac{1 - 1}{(\cos A + \cos B) \cdot (\sin A + \sin B)} \\
 &= 0 \\
 &= \text{RHS}
 \end{aligned}$$

ii) LHS

$$\begin{aligned}
 &= \frac{\sin^3 A - \cos^3 A}{\sin A + \cos A} + \frac{\sin^3 A + \cos^3 A}{\sin A - \cos A} \\
 &= \frac{(\sin A + \cos A) \cdot (\sin^2 A - \sin A \cos A + \cos^2 A)}{(\sin A + \cos A) \cdot (\sin A - \cos A)} \\
 &\quad + \frac{(\sin A - \cos A) \cdot (\sin^2 A + \sin A \cos A + \cos^2 A)}{(\sin A - \cos A)} \\
 &= (1 - \sin A \cos A) + (1 + \sin A \cos A) \\
 &= 2 \\
 &= \text{RHS}
 \end{aligned}$$

7. i) If $\sin \theta + \cos \theta = \sqrt{3}$, then prove that $\tan \theta + \cot \theta = 1$

ii) If $\sqrt{3} \sin \theta - \cos \theta = 0$, then show that $\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}$

Solution :

$$\begin{aligned}
 \text{i) Given } \sin \theta + \cos \theta &= \sqrt{3} \\
 \Rightarrow (\sin \theta + \cos \theta)^2 &= 3 \\
 \Rightarrow \sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta &= 3 \\
 \Rightarrow 1 + 2 \sin \theta \cos \theta &= 3 \\
 \Rightarrow \sin \theta \cos \theta &= 1 \quad \dots\dots (1)
 \end{aligned}$$

$$\text{TP : } \tan \theta + \cot \theta = 1$$

$$\text{LHS : } \tan \theta + \cot \theta$$

$$\begin{aligned}
 &= \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} \\
 &= \frac{\sin^2 \theta + \cos^2 \theta}{\cos \theta \sin \theta} \\
 &= \frac{1}{1} \quad (\text{from (1)}) \\
 &= 1 \\
 &= \text{RHS}
 \end{aligned}$$

Hence proved.

ii) Given $\sqrt{3} \sin \theta - \cos \theta = 0$

$$\begin{aligned}
 &\Rightarrow \sqrt{3} \sin \theta = \cos \theta \\
 &\Rightarrow \frac{\sin \theta}{\cos \theta} = \frac{1}{\sqrt{3}} \\
 &\Rightarrow \tan \theta = \frac{1}{\sqrt{3}} \Rightarrow \theta = 30^\circ
 \end{aligned}$$

$$T.P : \tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}$$

LHS

$$\begin{aligned}
 \tan 3\theta &= \tan 3(30^\circ) \\
 &= \tan 90^\circ \\
 &= \text{undefined}
 \end{aligned}$$

RHS

$$\begin{aligned}
 &= \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \\
 &= \frac{3 \left(\frac{1}{\sqrt{3}} \right) - \left(\frac{1}{\sqrt{3}} \right)^3}{1 - 3 \left(\frac{1}{\sqrt{3}} \right)^2} \\
 &= \frac{\sqrt{3} - \frac{1}{3\sqrt{3}}}{1 - 1}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\sqrt{3} - \frac{1}{3\sqrt{3}}}{0}
 \end{aligned}$$

= undefined

= LHS = RHS

= Hence proved.

8. i) If $\frac{\cos \alpha}{\sin \beta} = m$ and $\frac{\cos \alpha}{\sin \beta} = n$, then prove

that $(m^2 + n^2) \cos^2 \beta = n^2$

ii) If $\cot \theta + \tan \theta = x$ and $\sec \theta - \cos \theta = y$,

then prove that $(x^2 y)^{\frac{2}{3}} - (xy^2)^{\frac{2}{3}} = 1$

Solution :

i) Given $\frac{\cos \alpha}{\sin \beta} = m$, $\frac{\cos \alpha}{\sin \beta} = n$

$$\frac{\cos^2 \alpha}{\cos^2 \beta} \quad \frac{\cos^2 \alpha}{\sin^2 \beta}$$

To Prove : $(m^2 + n^2) \cos^2 \beta = n^2$

LHS :

$$\begin{aligned}
 &(m^2 + n^2) \cos^2 \beta \\
 &= \left(\frac{\cos^2 \alpha}{\cos^2 \beta} + \frac{\cos^2 \alpha}{\sin^2 \beta} \right) \cos^2 \beta \\
 &= \cos^2 \alpha \left(\frac{1}{\cos^2 \beta} + \frac{1}{\sin^2 \beta} \right) \cdot \cos^2 \beta \\
 &= \cos^2 \alpha \left(\frac{\sin^2 \beta + \cos^2 \beta}{\cos^2 \beta \cdot \sin^2 \beta} \right) \cdot \cos^2 \beta \\
 &= \frac{\cos^2 \alpha}{\sin^2 \beta} \\
 &= n^2 \\
 &= \text{RHS}
 \end{aligned}$$

ii) Given

$$\begin{aligned} x &= \cot \theta + \tan \theta & y &= \sec \theta - \cos \theta \\ &= \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{\cos \theta} & &= \frac{1}{\cos \theta} - \cos \theta \\ &= \frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta \cos \theta} & &= \frac{1 - \cos^2 \theta}{\cos \theta} \\ &= \frac{1}{\sin \theta \cdot \cos \theta} & &= \frac{\sin^2 \theta}{\cos \theta} \end{aligned}$$

To Prove :

$$(x^2 y)^{\frac{2}{3}} - (xy^2)^{\frac{2}{3}} = 1$$

LHS

$$\begin{aligned} &= \left(\frac{1}{\sin^2 \theta \cos^2 \theta} \times \frac{\sin^2 \theta}{\cos \theta} \right)^{\frac{2}{3}} \\ &\quad - \left(\frac{1}{\sin \theta \cdot \cos \theta} \times \frac{\sin^4 \theta}{\cos^2 \theta} \right)^{\frac{2}{3}} \\ &= \left(\frac{1}{\cos^3 \theta} \right)^{\frac{2}{3}} - \left(\frac{\sin^3 \theta}{\cos^3 \theta} \right)^{\frac{2}{3}} \\ &= (\sec^3 \theta)^{\frac{2}{3}} - (\tan^3 \theta)^{\frac{2}{3}} \\ &= \sec^2 \theta - \tan^2 \theta \\ &= 1 \\ &= 1 \\ &= \text{RHS} \end{aligned}$$

9. i) If $\sin \theta + \cos \theta = p$ and $\sec \theta + \operatorname{cosec} \theta = q$, then prove that $q(p^2 - 1) = 2p$
 ii) If $\sin \theta (1 + \sin^2 \theta) = \cos^2 \theta$, then prove that $\cos^6 \theta - 4 \cos^4 \theta + 8 \cos^2 \theta = 4$

Solution :

i) Given $p = \sin \theta + \cos \theta$,
 $q = \sec \theta + \operatorname{cosec} \theta$

To Prove : $q(p^2 - 1) = 2p$

LHS :

$$\begin{aligned} &q(p^2 - 1) \\ &= (\sec \theta + \operatorname{cosec} \theta) [(\sin \theta + \cos \theta)^2 - 1] \\ &= \left(\frac{1}{\cos \theta} + \frac{1}{\sin \theta} \right) [\sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta - 1] \\ &= \frac{\sin \theta + \cos \theta}{\cos \theta \cdot \sin \theta} [1 + 2 \sin \theta \cos \theta - 1] \\ &= \frac{\sin \theta + \cos \theta}{\cos \theta \cdot \sin \theta} \times 2 \sin \theta \cos \theta \\ &= 2(\sin \theta \cos \theta) \\ &= 2p \\ &= \text{RHS} \end{aligned}$$

ii) Given $\sin \theta (1 + \sin^2 \theta) = \cos^2 \theta$

To Prove :

$$\cos^6 \theta - 4 \cos^4 \theta + 8 \cos^2 \theta = 4$$

10. If $\frac{\cos \theta}{1 + \sin \theta} = \frac{1}{a}$, then prove that

$$\frac{a^2 - 1}{a^2 + 1} = \sin \theta$$

Solution :

Given

$$\frac{\cos \theta}{1 + \sin \theta} = \frac{1}{a}$$

To Prove : $\frac{a^2 - 1}{a^2 + 1} = \sin \theta$

$$\begin{aligned} \therefore a &= \frac{1 + \sin \theta}{\cos \theta} \\ &= (1 + \sin \theta) \sec \theta \\ &= \sec \theta + \tan \theta \end{aligned}$$


∴ LHS :

$$\begin{aligned}
 &= \frac{a^2 - 1}{a^2 + 1} \\
 &= \frac{(\sec \theta + \tan \theta)^2 - 1}{(\sec \theta + \tan \theta)^2 + 1} \\
 &= \frac{\sec^2 \theta + \tan^2 \theta + 2 \sec \theta \tan \theta - 1}{\sec^2 \theta + \tan^2 \theta + 2 \sec \theta \tan \theta + 1} \\
 &= \frac{2 \tan^2 \theta + 2 \sec \theta \tan \theta}{2 \sec^2 \theta + 2 \sec \theta \tan \theta} \\
 &= \frac{\cancel{2} \tan \theta (\cancel{\tan \theta + \sec \theta})}{\cancel{2} \sec \theta (\cancel{\sec \theta + \tan \theta})} \\
 &= \frac{\sin \theta}{\cos \theta} \\
 &= \frac{1}{\cos \theta} \\
 &= \sin \theta \\
 &= \text{RHS}
 \end{aligned}$$

II. PROBLEMS INVOLVING ANGLE OF ELEVATION

Example 6.18

Calculate the size of $\angle BAC$ in the given triangles.



Solution :

(i) In right triangle ABC [see Fig.]

$$\tan \theta = \frac{\text{opposite side}}{\text{adjacent side}} = \frac{4}{5}$$

$$\theta = \tan^{-1} \left(\frac{4}{5} \right) = \tan^{-1} (0.8)$$

$$\theta = 38.7^\circ \text{ (since } \tan 38.7^\circ = 0.8011 \text{)}$$

$$\angle BAC = 38.7^\circ$$

(ii) In right triangle ABC [see Fig.6.12(b)]

$$\tan \theta = \frac{8}{3}$$


$$\theta = \tan^{-1} \left(\frac{8}{3} \right) = \tan^{-1} (2.66)$$

$$\theta = 69.4^\circ \text{ (since } \tan 69.4^\circ = 2.6604 \text{)}$$

$$\angle BAC = 69.4^\circ$$

Example 6.19

A tower stands vertically on the ground. From a point on the ground, which is 48 m away from the foot of the tower, the angle of elevation of the top of the tower is 30° . Find the height of the tower.



Solution :

Let PQ be the height of the tower.

Take $PQ = h$ and QR is the distance between the tower and the point R. In right triangle PQR, $\angle PRQ = 30^\circ$

$$\tan \theta = \frac{PQ}{QR}$$


$$\tan 30^\circ = \frac{h}{48}$$

$$\text{gives, } \frac{1}{\sqrt{3}} = \frac{h}{48} \text{ so, } h = 16\sqrt{3}$$

Therefore the height of the tower is $16\sqrt{3}$ m

Example 6.20

A kite is flying at a height of 75 m above the ground. The string attached to the kite is temporarily tied to a point on the ground. The inclination of the string with the ground is 60° . Find the length of the string, assuming that there is no slack in the string.



Solution :

Let AB be the height of the kite above the ground. Then, $AB = 75$.

Let AC be the length of the string.

In right triangle ABC, $\angle ACB = 60^\circ$

$$\sin \theta = \frac{AB}{AC}$$

$$\sin 60^\circ = \frac{75}{AC}$$


$$\text{gives, } \frac{\sqrt{3}}{2} = \frac{75}{AC} \text{ so, } AC = \frac{150}{\sqrt{3}} = 50\sqrt{3}$$

Hence, the length of the string is $50\sqrt{3}$ m

Example 6.21

Two ships are sailing in the sea on either sides of a lighthouse. The angle of elevation of the top of the lighthouse as observed from the ships are 30° and 45° respectively. If the lighthouse is 200 m high, find the distance between the two ships. ($\sqrt{3} = 1.732$)

Solution :



Let AB be the lighthouse. Let C and D be

Then, $AB = 200$ m.

$$\angle ACB = 30^\circ, \angle ADB = 45^\circ$$

In right triangle BAC,

$$\tan 30^\circ = \frac{AB}{AC}$$

$$\frac{1}{\sqrt{3}} = \frac{200}{AC}$$

$$\text{gives, } AC = 200\sqrt{3} \quad \dots\dots (1)$$

In right triangle BAD,

$$\tan 45^\circ = \frac{AB}{AD}$$

$$1 = \frac{200}{AD}$$

$$\text{gives, } AD = 200 \quad \dots\dots (2)$$

Now, $CD = AC + AD$

$$= 200\sqrt{3} + 200 [\text{by (1) and (2)}]$$

$$CD = 200 (\sqrt{3} + 1)$$

$$= 200 \times 2.732 = 546.4$$

Distance between two ships is 546.4 m.

Example 6.22

From a point on the ground, the angles of elevation of the bottom and top of a tower fixed at the top of a 30 m high building are 45° and 60° respectively. Find the height of the tower. ($\sqrt{3} = 1.732$)

Solution :



Fig. 6.16

Let AC be the height of the tower.

Let AB be the height of the building.

Then, AC = h metres, AB = 30 m

In right triangle CBP,

$$\angle CPB = 60^\circ$$

$$\tan \theta = \frac{BC}{BP}$$

$$\tan 60^\circ = \frac{AB + AC}{BP}$$

$$\text{so, } \sqrt{3} = \frac{30 + h}{BP} \quad \dots\dots (1)$$

In right triangle ABP,

$$\angle APB = 45^\circ$$

$$\tan \theta = \frac{AB}{BP}$$

$$\tan 45^\circ = \frac{30}{BP}$$

$$\text{gives, } BP = 30 \quad \dots\dots (2)$$

Substituting (2) in (1), we get

$$\sqrt{3} = \frac{30 + h}{30}$$

$$h = 30(\sqrt{3} - 1)$$

$$= 30(1.732 - 1)$$

$$= 30(0.732) = 21.96$$

Hence, the height of the tower is 21.96 m.

Example 6.23

A TV tower stands vertically on a bank of a ca-

nal. From a point D on the other bank directly opposite to it, the angle of elevation of the top of the tower is 58° . From another point 20 m away from this point on the line joining this point to the foot of the tower, the angle of elevation of the top of the tower is 30° . Find the height of the tower and the width of the canal. ($\tan 58^\circ = 1.6003$)

Solution :




Fig. 6.17

Let AB be the height of the TV tower.

CD = 20 m.

Let BC be the width of the canal.

In right triangle ABC,

$$\tan 58^\circ = \frac{AB}{BC}$$

$$1.6003 = \frac{AB}{BC} \quad \dots\dots\dots (1)$$

In right triangle ABD,

$$\tan 30^\circ = \frac{AB}{BD} = \frac{AB}{BC + CD}$$

$$\frac{1}{\sqrt{3}} = \frac{AB}{BC + 20} \quad \dots\dots\dots (2)$$

Dividing (1) by (2) we get,

$$\frac{1.6003}{\frac{1}{\sqrt{3}}} = \frac{BC + 20}{BC}$$

$$BC = \frac{20}{1.7791} = 11.24m \quad \dots\dots\dots (3)$$

$$1.6003 = \frac{AB}{11.24} \quad [\text{from (1) and (3)}]$$

$$AB = 17.99$$

Hence, the height of the tower is 17.99 m and the width of the canal is 11.24 m.


Example 6.24

An aeroplane sets off from G on a bearing of 24° towards H, a point 250 km away. At H it changes course and heads towards J on a bearing of 55° and a distance of 180 km away.

- How far is H to the North of G?
- How far is H to the East of G?
- How far is J to the North of H?
- How far is J to the East of H?

$$\left(\begin{array}{ll} \sin 24^\circ = 0.4067 & \sin 11^\circ = 0.1908 \\ \cos 24^\circ = 0.9135 & \cos 11^\circ = 0.9816 \end{array} \right)$$

Solution :



(i) In right triangle GOH,


$$\cos 24^\circ = \frac{OG}{GH}$$

$$0.9135 = \frac{OG}{250}; OG = 228.38 \text{ km}$$

Distance of H to the North of

$$G = 228.38 \text{ km}$$

(ii) In right triangle GOH,



$$\sin 24^\circ = \frac{OH}{GH}$$

$$0.4067 = \frac{OH}{250}; OH = 101.68$$

Distance of H to the East of

$$G = 101.68 \text{ km}$$

(iii) In right triangle HIJ,

$$\sin 11^\circ = \frac{IJ}{HJ}$$

$$0.1908 = \frac{IJ}{180}; IJ = 34.34 \text{ km}$$

Distance of J to the North of

$$H = 34.34 \text{ km}$$

(iv) In right triangle HIJ,

$$\cos 11^\circ = \frac{HI}{HJ}$$

$$0.9816 = \frac{HI}{180}; HI = 176.69 \text{ km}$$

Distance of J to the East of

$$H = 176.69 \text{ km}$$

Example 6.25

Two trees are standing on flat ground. The angle of elevation of the top of both the trees from a point X on the ground is 40° . If the horizontal distance between X and the smaller tree is 8 m and the distance of the top of the two trees is 20 m, calculate

- the distance between the point X and the top of the smaller tree.
- the horizontal distance between the two trees. ($\cos 40^\circ = 0.7660$)

Solution :

Let AB be the height of the bigger tree and CD be the height of the smaller tree and X is the point on the ground.

(i) In right triangle XCD,

$$\cos 40^\circ = \frac{CX}{XD}$$

$$XD = \frac{8}{0.7660} = 10.44 \text{ km}$$

Therefore the distance between X and top of the smaller tree = $XD = 10.44 \text{ m}$

(ii) In right triangle XAB,

$$\cos 40^\circ = \frac{AX}{BX}$$

$$= \frac{AC + CX}{BD + DX}$$

$$= 0.7660 = \frac{AC + 8}{20 + 10.44}$$

$$\text{gives } AC = 23.32 - 8 = 15.32 \text{ m}$$


Therefore the horizontal distance between two trees = $AC = 15.32 \text{ m}$

EXERCISE 6.2

- Find the angle of elevation of the top of a tower from a point on the ground, which is 30 m away from the foot of a tower of


$\sqrt{\quad}$

Solution :



$$\begin{aligned} \text{From the fig, } \tan \theta &= \frac{10\sqrt{3}}{30} \\ &= \frac{1}{\sqrt{3}} \\ \therefore \theta &= 30^\circ \end{aligned}$$

- A road is flanked on either side by continuous rows of houses of height $4\sqrt{3} \text{ m}$ with no space in between them. A pedestrian is standing on the median of the road facing a row house. The angle of elevation from the pedestrian to the top of the house is 30° . Find the width of the road.

Solution :Let $AB =$ Width of the road $P =$ Midpoint of AB $BC =$ height of the row houses

$$= 4\sqrt{3} \text{ m}$$

Let $PB = PA = x \text{ m}$

$$\text{In } \triangle PBC, \tan 30^\circ = \frac{BC}{PB}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{4\sqrt{3}}{x}$$


$$\Rightarrow x = 12 \text{ m}$$

$$= 2x$$

$$= 2(12)$$

$$= 24 \text{ m}$$

3. To a man standing outside his house, the angles of elevation of the top and bottom of a window are 60° and 45° respectively. If the height of the man is 180 cm and if he is 5 m away from the wall, what is the height of the window? ($\sqrt{3} = 1.732$)

Solution :

Let $MN = 180 \text{ cm} = 1.8 \text{ m}$
 $=$ height of the man

$DN = 5 \text{ m} =$ Dist. between
 Man & Wall

$T, B \rightarrow$ Top & Bottom of Window

$BC = x, TB = y$ (Height of window)

$$\text{In } \triangle CMB, \tan 45^\circ = \frac{x}{5}$$

$$\Rightarrow 1 = \frac{x}{5}$$

$$\Rightarrow x = 5$$

$$\text{In } \triangle CMT, \tan 60^\circ = \frac{x+y}{5}$$

$$\Rightarrow \sqrt{3} = \frac{5+y}{5}$$

$$\Rightarrow 5+y = 5\sqrt{3}$$

$$\Rightarrow y = 5\sqrt{3} - 5$$


$$= 5(\sqrt{3} - 1)$$

$$= 5(0.732)$$

$$= 3.66 \text{ m}$$

\therefore Height of window = 3.66 m

4. A statue 1.6 m tall stands on the top of a pedestal. From a point on the ground, the angle of elevation of the top of the statue is 60° and from the same point the angle of elevation of the top of the pedestal is 40° . Find the height of the pedestal. ($\tan 40^\circ = 0.8391, (\sqrt{3} = 1.732)$)

Solution :

Let D → Point of observation on the ground

BC = x m = height of Pedestal

AC = 1.6m = height of statue

$\angle BDC = 40^\circ$, $\angle BDA = 60^\circ$, BD = y m

$$\text{In } \triangle BCD, \tan 40^\circ = \frac{x}{y}$$

$$\begin{aligned} \Rightarrow y &= \frac{x}{\tan 40^\circ} \\ &= \frac{x}{0.8391} \end{aligned} \quad \dots\dots\dots (1)$$

$$\text{In } \triangle BAD, \tan 60^\circ = \frac{AB}{BD}$$

$$\begin{aligned} \Rightarrow \sqrt{3} &= \frac{x+1.6}{y} \\ \Rightarrow y &= \frac{x+1.6}{\sqrt{3}} \end{aligned} \quad \dots\dots\dots (2)$$


∴ From (1) & (2)

$$\begin{aligned} \frac{x}{0.8391} &= \frac{x+1.6}{1.732} \\ \Rightarrow 1.732x &= 0.8391x + 1.6(0.8391) \\ \Rightarrow 0.8929x &= 1.343 \\ \Rightarrow x &= \frac{1.343}{0.8929} \\ &= 1.5 \text{ m} \end{aligned}$$


∴ Height of statue = 1.5 m

5. A flag pole 'h' metres is on the top of the hemispherical dome of radius 'r' metres. A man is standing 7 m away from the dome. Seeing the top of the pole at an angle 45° and moving 5 m away from the dome and seeing the bottom of the pole at an angle 30° . Find (i) the height of the pole (ii) radius of the dome.

($\sqrt{3} = 1.732$)



Solution :



$$\text{In } \triangle ACD, \tan 45^\circ = \frac{AC}{CD}$$

$$1 = \frac{h+r}{r+7}$$

$$\Rightarrow r+7 = h+r$$

$$\Rightarrow h = 7$$

∴ Height of the pole = 7m

$$\text{In } \triangle BCE, \tan 30^\circ = \frac{BC}{CE}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{r}{r+7+5}$$

$$\Rightarrow \sqrt{3}r = r+12$$

$$\Rightarrow \sqrt{3}r - r = 12$$

$$\Rightarrow r(\sqrt{3} - 1) = 12$$

$$\Rightarrow r = \frac{12}{\sqrt{3} - 1} \times \frac{\sqrt{3} + 1}{\sqrt{3} + 1}$$

$$= \frac{12(\sqrt{3} + 1)}{2}$$


$$= 6(2.732)$$

$$= 16.392 \text{ m}$$

∴ Radius of dome = 16.39 m

6. The top of a 15 m high tower makes an angle of elevation of 60° with the bottom of an electronic pole and angle of elevation of 30° with the top of the pole. What is the height of the electric pole?

Solution :



Let $AB = 15 \text{ m} = \text{Height of the tower}$

$CD = x \text{ m} = \text{Height of the pole} = BE$

Let $BD = EC = y$

$$\text{In } \triangle ADB, \tan 60^\circ = \frac{AB}{BD}$$

$$\sqrt{3} = \frac{15}{y}$$

$$\Rightarrow y = \frac{15}{\sqrt{3}} = 5\sqrt{3} \quad \dots\dots\dots (1)$$

$$\text{In } \triangle ACE, \tan 30^\circ = \frac{AE}{EC}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{15-x}{y}$$

$$\Rightarrow 5\sqrt{3} = (15-x)\sqrt{3} \quad (\text{From (1)})$$


$$\Rightarrow 5 = 15 - x$$

$$\Rightarrow x = 10$$

$\therefore \text{Height of the pole} = 10\text{m}$

7. A vertical pole fixed to the ground is divided in the ratio 1 : 9 by a mark on it with lower part shorter than the upper part. If the two parts subtend equal angles at a place on the ground, 25 m away from the base of the pole, what is the height of the pole?

Solution :



Let $AC = \text{height of the pole}$

B is a point on AC which divides it in the ratio 1 : 9 such that $AB = 9x$, $BC = x$

(\because lower part is shorter than upper part)

$CD = 25 \text{ m} = \text{Dist. between pole and point of observation}$

$$\text{In } \triangle BCD, \tan \theta = \frac{BC}{CD} = \frac{x}{25}$$

$$\text{In } \triangle ACD, \tan 2\theta = \frac{AC}{CD} = \frac{10x}{25}$$

$$\Rightarrow \tan 2\theta = \frac{10x}{25}$$

$$\Rightarrow \frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{10x}{25}$$

$$\Rightarrow \frac{2 \left(\frac{x}{25} \right)}{1 - \frac{x^2}{25}} = \frac{2x}{5}$$


$$\Rightarrow \frac{\frac{2x}{25}}{\frac{625 - x^2}{625}} = \frac{2x}{5}$$

$$\begin{aligned} \Rightarrow \frac{2x}{25} \times \frac{625}{625 - x^2} &= \frac{2x}{5} \\ \Rightarrow \frac{25}{625 - x^2} &= \frac{1}{5} \\ \Rightarrow 625 - x^2 &= 125 \\ \Rightarrow x^2 &= 500 \\ \Rightarrow x &= 10\sqrt{5} \end{aligned}$$

$$\begin{aligned} \therefore \text{Height of the pole} &= 10x \\ &= 10(10\sqrt{5}) \\ &= 100\sqrt{5} \text{ m} \end{aligned}$$

8. A traveler approaches a mountain on highway. He measures the angle of elevation to the peak at each milestone. At two consecutive milestones the angles measured are 4° and 8° . What is the height of the peak if the distance between consecutive milestones is 1 mile. ($\tan 4^\circ = 0.0699$, $\tan 8^\circ = 0.1405$)

Solution :



Let $PK = h =$ height of mountain

$M_1, M_2 =$ Mile stones, $M_1, M_2 = 1$ mile

Let $KM_1 = y$

In $\triangle PKM_1$,

$$\begin{aligned} \tan 8^\circ &= \frac{h}{y} \\ \Rightarrow y &= \frac{h}{\tan 8^\circ} \quad \dots\dots\dots (1) \end{aligned}$$

In $\triangle PKM_2$,

$$\begin{aligned} \tan 4^\circ &= \frac{h}{y+1} \\ \Rightarrow y+1 &= \frac{h}{\tan 4^\circ} \\ \Rightarrow y &= \frac{h}{\tan 4^\circ} - 1 \quad \dots\dots\dots (2) \end{aligned}$$

\therefore From (1) & (2)


$$\begin{aligned} \frac{h}{\tan 8^\circ} &= \frac{h}{\tan 4^\circ} - 1 \\ \Rightarrow \frac{h}{\tan 4^\circ} - \frac{h}{\tan 8^\circ} &= 1 \\ \Rightarrow h \left[\frac{1}{\tan 4^\circ} - \frac{1}{\tan 8^\circ} \right] &= 1 \\ \Rightarrow h \left[\frac{\tan 8^\circ - \tan 4^\circ}{\tan 8^\circ \tan 4^\circ} \right] &= 1 \\ \Rightarrow h &= \frac{\tan 8^\circ - \tan 4^\circ}{\tan 8^\circ \tan 4^\circ} \\ &= \frac{0.14 - 0.07}{0.14 \times 0.07} \\ &= \frac{0.0098}{0.07} \\ &= 0.14 \text{ miles} \end{aligned}$$

III. PROBLEMS INVOLVING ANGLE OF DEPRESSION

Example 6.26

A player sitting on the top of a tower of height 20 m observes the angle of depression of a ball lying on the ground as 60° . Find the distance between the foot of the tower and the ball. ($\sqrt{3} = 1.732$)

Solution :



Let BC be the height of the tower and A be the position of the ball lying on the ground. Then,

$$BC = 20 \text{ m and } \angle XCA = 60^\circ = \angle CAB$$

Let AB = x metres.

In right triangle ABC,

$$\tan 60^\circ = \frac{BC}{AB}$$

$$\sqrt{3} = \frac{20}{x}$$

$$x = \frac{20 \times \sqrt{3}}{\sqrt{3} \times \sqrt{3}} = \frac{20 \times 1.732}{3}$$

$$= 11.54 \text{ m}$$

Hence, the distance between the foot of the tower and the ball is 11.54 m.

The horizontal distance between two buildings is 140 m. The angle of depression of the top of the first building when seen from the top of the second building is 30° . If the height of the first building is 60 m, find the height of the second building. ($\sqrt{3} = 1.732$)

Solution :




Fig. 6.22

The height of the first building AB = 60 m. Now, AB = MD = 60 m
Let the height of the second building CD = h. Distance BD = 140 m
Now, AM = BD = 140 m
From the diagram,

$$\angle XCA = 30^\circ = \angle CAM$$

In right triangle AMC,

$$\tan 30^\circ = \frac{CM}{AM}$$

$$\frac{1}{\sqrt{3}} = \frac{CM}{140}$$

$$CM = \frac{140}{\sqrt{3}} = \frac{140\sqrt{3}}{3} \\ = \frac{140 \times 1.732}{3}$$

$$CM = 80.78$$

$$\begin{aligned} \text{Now, } h &= CD \\ &= CM + MD \\ &= 80.78 + 60 = 140.78 \end{aligned}$$

Therefore the height of the second building is 140.78 m

Example 6.28

From the top of a tower 50 m high, the angles of depression of the top and bottom of a tree are observed to be 30° and 45° respectively. Find the height of the tree. ($\sqrt{3} = 1.732$)

Solution :




Fig. 6.23

The height of the tower AB = 50 m

Let the height of the tree

$$CD = y \text{ and } BD = x$$

From the diagram,

$$\angle XAC = 30^\circ = \angle ACM \text{ and}$$

$$\angle XAD = 45^\circ = \angle ADB$$

In right triangle ABD,

$$\tan 45^\circ = \frac{AB}{BD}$$

$$1 = \frac{50}{x} \text{ gives } x = 50 \text{ m}$$

In right triangle AMC,

$$\tan 30^\circ = \frac{AM}{CM}$$

$$\frac{1}{\sqrt{3}} = \frac{AM}{50} \text{ [since } DB = CM]$$

$$\begin{aligned} AM &= \frac{50}{\sqrt{3}} = \frac{50\sqrt{3}}{3} \\ &= \frac{50 \times 1.732}{3} = 28.85 \text{ m} \end{aligned}$$

Therefore, height of the tree =

$$CD = MB = AB - AM$$

$$= 50 - 28.85 = 21.15 \text{ m}$$

As observed from the top of a 60 m high light house from the sea level, the angles of depression of two ships are 28° and 45° . If one ship is exactly behind the other on the same side of the lighthouse, find the distance between the two ships. ($\tan 28^\circ = 0.5317$)

Solution :




Fig. 6.24

Let the observer on the lighthouse CD be at D.

Height of the lighthouse CD = 60 m

From the diagram,

$$\angle XDA = 28^\circ = \angle DAC \text{ and}$$

$$\angle XDB = 45^\circ = \angle DBC$$

In right triangle DCB,

$$\tan 45^\circ = \frac{DC}{BC}$$

$$1 = \frac{60}{BC} \text{ gives } BC = 60 \text{ m}$$

In right triangle DCA,

$$\tan 28^\circ = \frac{DC}{AC}$$

$$0.5317 = \frac{60}{AC}$$

$$\begin{aligned} \text{gives } AC &= \frac{60}{0.5317} \\ &= 112.85 \end{aligned}$$

Distance between the two ships

$$AB = AC - BC = 52.85 \text{ m}$$

Example 6.30

A man is watching a boat speeding away from

depression of 60° with the man's eye when at a distance of 200 m from the tower. After 10 seconds, the angle of depression becomes 45° . What is the approximate speed of the boat (in km / hr), assuming that it is sailing in still water? ($\sqrt{3} = 1.732$)

Solution :




Fig. 6.25

Let AB be the tower.

Let C and D be the positions of the boat.

From the diagram,

$$\angle XAC = 60^\circ = \angle ACB \text{ and}$$

$$\angle XAD = 45^\circ = \angle ADB, BC = 200 \text{ m}$$

$$\text{In right triangle ABC, } \tan 60^\circ = \frac{AB}{BC}$$

$$\text{gives } \sqrt{3} = \frac{AB}{200}$$

$$\text{we get } AB = 200\sqrt{3} \quad \dots\dots\dots (1)$$

$$\text{In right triangle ABD, } \tan 45^\circ = \frac{AB}{BD}$$

$$\text{gives } 1 = \frac{200\sqrt{3}}{BD} \text{ [by (1)]}$$

$$\text{we get } BD = 200\sqrt{3}$$

$$\text{Now, } CD = BD - BC$$

$$\begin{aligned} CD &= 200\sqrt{3} - 200 \\ &= 200(\sqrt{3} - 1) = 146.4 \end{aligned}$$

10 seconds.

That is, the distance of 146.4 m is covered in 10 seconds.


$$\begin{aligned} \text{Therefore, speed of the boat} &= \frac{\text{distance}}{\text{time}} \\ &= \frac{146.4}{10} = 14.64 \text{ m/s} \end{aligned}$$

$$\begin{aligned} \text{gives } 14.64 \times \frac{3600}{1000} \text{ km/hr} \\ = 52.704 \text{ km/hr} \end{aligned}$$

EXERCISE 6.3

- From the top of a rock $50\sqrt{3}$ m high, the angle of depression of a car on the ground is observed to be 30° . Find the distance of the car from the rock.

Solution:



$$RK = 50\sqrt{3} \text{ m} = \text{height of the rock}$$

C = Position of the car

$$KC = x \text{ m}$$

$$\angle TRC = \angle RCK = 30^\circ$$

$$\tan 30^\circ = \frac{RK}{KC}$$


$$\frac{1}{\sqrt{3}} = \frac{50\sqrt{3}}{x}$$

$$\Rightarrow x = 150 \text{ m}$$

\therefore Dist. of the car from the rock = 150 m

- The horizontal distance between two buildings is 70 m. The angle of depression of the top of the first building when seen from the top of the second building is 45° . If the height of the second building is 120 m, find the height of the first building.

Solution:



$$\text{Height of the 1st building} = CD = x \text{ m}$$

$$\text{Height of the 2nd building} = AB = 120 \text{ m}$$

$$\text{Distance between 2 buildings} =$$

$$BD = EC = 70 \text{ m}$$

In $\triangle AEC$,

$$\tan 45^\circ = \frac{AE}{EC}$$

$$1 = \frac{120 - x}{70}$$

$$\Rightarrow 70 = 120 - x$$


$$\Rightarrow x = 120 - 70$$

$$\Rightarrow x = 50$$

\therefore Height of 1st building = 50 m

3. From the top of the tower 60 m high the angles of depression of the top and bottom of a vertical lamp post are observed to be 38° and 60° respectively. Find the height of the lamp post. ($\tan 38^\circ = 0.7813$, $\sqrt{3} = 1.732$)

Solution:



TR = Height of the tower = 60 m

LP = Height of the lamp post = x m = WR

$$\therefore TW = 60 - x$$

Let RP = WL = y

In $\triangle TRP$,

$$\tan 60^\circ = \frac{TR}{RP}$$

$$\Rightarrow \sqrt{3} = \frac{60}{y}$$

$$\Rightarrow y = \frac{60}{\sqrt{3}}$$

$$\Rightarrow y = 20\sqrt{3} \quad \dots\dots\dots (1)$$

In $\triangle TWL$,

$$\tan 30^\circ = \frac{TW}{WL}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{60 - x}{y}$$

$$\Rightarrow y = \sqrt{3} (60 - x) \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\Rightarrow 20\sqrt{3} = \sqrt{3} (60 - x)$$


$$\Rightarrow 20 = 60 - x$$

$$\Rightarrow x = 60 - 20 \Rightarrow x = 40$$

\therefore Height of the lamp post = 40m

4. An aeroplane at an altitude of 1800 m finds that two boats are sailing towards it in the same direction. The angles of depression of the boats as observed from the aeroplane are 60° and 30° respectively. Find the distance between the two boats. ($\sqrt{3} = 1.732$)

Solution:



AP = 1800 m

= height of the plane from the ground

B_1, B_2 = Positions of 2 boats

Let $B_1, B_2 = x$ m ; $PB_1 = y$

In $\triangle APB_1$,

$$\tan 60^\circ = \frac{1800}{y}$$

$$\Rightarrow \sqrt{3} = \frac{1800}{y}$$


$$\Rightarrow y = \frac{1800}{\sqrt{3}} = 600\sqrt{3} \quad \dots\dots\dots (1)$$

In $\triangle APB_2$,

$$\begin{aligned}\tan 30^\circ &= \frac{1800}{y+x} \\ \Rightarrow \frac{1}{\sqrt{3}} &= \frac{1800}{y+x} \\ \Rightarrow y+x &= 1800\sqrt{3} \\ \Rightarrow 600\sqrt{3} + x &= 1800\sqrt{3} \\ (\text{From (1)}) \\ \Rightarrow x &= 1200\sqrt{3} \\ x &= 1200 (1.732) \\ &= 2078.4 \text{ m}\end{aligned}$$

5. From the top of a lighthouse, the angle of depression of two ships on the opposite sides of it are observed to be 30° and 60° . If the height of the lighthouse is h m through the foot of the lighthouse, show that the distance between the ships is $\frac{4h}{\sqrt{3}}$ m.

Solution:



LH = h m = height of the light house
 S_1, S_2 = Positions of 2 ships
 $S_1H = x$ m, $S_2H = y$ m
 To find : $x + y$
 In $\triangle LS_1H$,

$$\tan 30^\circ = \frac{h}{x}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{x}$$

$$\Rightarrow x = \sqrt{3}h \quad \dots\dots\dots (1)$$

In $\triangle LS_2H$,

$$\tan 60^\circ = \frac{h}{y}$$

$$\Rightarrow \sqrt{3} = \frac{h}{y}$$

$$\Rightarrow y = \frac{h}{\sqrt{3}} \quad \dots\dots\dots (2)$$

\therefore Adding (1) & (2)


$$\begin{aligned}x + y &= \sqrt{3}h + \frac{h}{\sqrt{3}} \\ &= \frac{3h + h}{\sqrt{3}} = \frac{4h}{\sqrt{3}}\end{aligned}$$

\therefore Distance between 2 ships

$$= \frac{4h}{\sqrt{3}} \text{ m}$$

6. A lift in a building of height 90 feet with transparent glass walls is descending from the top of the building. At the top of the building, the angle of depression to a fountain in the garden is 60° . Two minutes later, the angle of depression reduces to 30° . If the fountain is $30\sqrt{3}$ feet from the entrance of the lift, find the speed of the lift which is descending.

Solution:



LT = height of the lift = 90 ft

F = Position of the fountain

FT = Distance between fountain & lift

$$= 30\sqrt{3} \text{ ft.}$$

$$LP = x \text{ ft} \Rightarrow PT = 90^\circ - x$$

Time taken from L to P = 2 min.

In ΔPFT ,

$$\tan 30^\circ = \frac{90 - x}{30\sqrt{3}}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{90 - x}{30\sqrt{3}}$$

$$\Rightarrow 30 = 90 - x$$

$$\Rightarrow x = 60 \text{ ft}$$

$$\therefore \text{Speed of the lift} = \frac{\text{Dist.}}{\text{Time}} = \frac{60}{2}$$


$$= 30 \text{ ft/min.}$$

PROBLEMS INVOLVING ANGLE OF ELEVATION AND DEPRESSION

Example 6.31

From the top of a 12 m high building, the angle of elevation of the top of a cable tower is 60° and the angle of depression of its foot is 30° . Determine the height of the tower.

Solution :



As shown in Fig., OA is the building, O is the point of observation on the top of the building OA. Then, OA = 12 m.

PP' is the cable tower with P as the top and P' as the bottom.

Then the angle of elevation of P, $\angle MOP = 60^\circ$.

And the angle of depression of P', $\angle MOP' = 30^\circ$

Suppose, height of the cable tower PP' = h metres.

Through O, draw $OM \perp PP'$

$$MP = PP' - MP' = h - OA = h - 12$$

In right triangle OMP, $\frac{MP}{OM} = \tan 60^\circ$

$$\text{gives } \frac{h - 12}{OM} = \sqrt{3}$$

$$\text{so, } OM = \frac{h - 12}{\sqrt{3}} \quad \dots\dots\dots (1)$$

In right triangle OMP', $\frac{MP'}{OM} = \tan 30^\circ$

$$\text{gives } \frac{12}{OM} = \frac{1}{\sqrt{3}}$$

$$\text{so, } OM = 12\sqrt{3} \quad \dots\dots\dots (2)$$

From (1) and (2) we have, $\frac{h - 12}{\sqrt{3}} = 12\sqrt{3}$

$$\text{gives, } h - 12 = 12\sqrt{3} \times \sqrt{3} \text{ we get, } h = 48$$

Hence, the required height of the cable tower is 48 m.

Example 6.32

A pole 5 m high is fixed on the top of a tower. The angle of elevation of the top of the pole observed from a point 'A' on the ground is 60° and the angle of depression to the point 'A' from the top of the tower is 45° . Find the height of the tower. ($\sqrt{3} = 1.732$)

Solution :




Fig. 6.28

Let BC be the height of the tower and CD be the height of the pole.

Let 'A' be the point of observation.

Let BC = x and AB = y.

From the diagram,

$\angle BAD = 60^\circ$ and $\angle XCA = 45^\circ = \angle BAC$

In right triangle ABC, $\tan 45^\circ = \frac{BC}{AB}$

$$\text{gives, } 1 = \frac{x}{y} \text{ so, } x = y \quad \dots\dots\dots (1)$$

In right triangle ABD, $\tan 60^\circ$

$$= \frac{BD}{AB} = \frac{BC + CD}{AB}$$

$$\text{gives, } \sqrt{3} = \frac{x+5}{y} \text{ so, } \sqrt{3}y = x+5$$

$$\text{we get, } \sqrt{3}x = x+5 \quad [\text{From (1)}]$$

$$\begin{aligned} \text{so, } x &= \frac{5}{\sqrt{3}-1} = \frac{5}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} \\ &= \frac{5(1.732+1)}{2} = 6.83 \end{aligned}$$


Hence, height of the tower is 6.83 m.

Example 6.33

From a window (h metres high above the ground) of a house in a street, the angles of elevation and depression of the top and the foot of another

house on the opposite side of the street are θ_1 and θ_2 respectively. Show that the height of the opposite house is $h \left(1 + \frac{\cot \theta_2}{\cot \theta_1} \right)$

Solution :



Let W be the point on the window where the angles of elevation and depression are measured.

Let PQ be the house on the opposite side.

Then WA is the width of the street.

$$= AQ \quad (WR = AQ)$$

Let PA = x metres.

$$\text{In right triangle } PAW, \tan \theta_1 = \frac{AP}{AW}$$

$$\text{gives } \tan \theta_1 = \frac{x}{AW}$$

$$\text{so, } AW = \frac{x}{\tan \theta_1}$$

$$\text{we get, } AW = x \cot \theta_1 \quad \dots\dots (1)$$

$$\text{In right triangle } QAW, \tan \theta_2 = \frac{AQ}{AW}$$

$$\text{gives } \tan \theta_2 = \frac{h}{AW}$$

$$\text{we get, } AW = h \cot \theta_2 \quad \dots\dots (2)$$

From (1) and (2) we get, $x \cot \theta_1 = h \cot \theta_2$

$$\text{gives, } x = h \frac{\cot \theta_2}{\cot \theta_1}$$

Therefore, height of the opposite house =


$$PA + AQ = x + h = h \frac{\cot \theta_2}{\cot \theta_1} + h = h \left(1 + \frac{\cot \theta_2}{\cot \theta_1} \right)$$

Hence Proved.

EXERCISE 6.4

1. From the top of a tree of height 13 m the angle of elevation and depression of the top and bottom of another tree are 45° and 30° respectively. Find the height of the second tree. ($\sqrt{3} = 1.732$)

Solution:



$CD = 13\text{m} = \text{height of tree 1}$

$AB = x + 13 = \text{height of tree 2}$

$BD = EC = y \text{ m} = \text{dist. between 2 trees}$

In $\triangle BCD$,

$$\tan 30^\circ = \frac{CD}{BD}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{13}{y}$$

$$\Rightarrow y = 13\sqrt{3} \text{ m}$$

In $\triangle ACE$,

$$\tan 45^\circ = \frac{AE}{EC}$$

$$\Rightarrow 1 = \frac{x}{y}$$

$$\Rightarrow y = x$$

$$\Rightarrow x = 13\sqrt{3} \text{ m}$$

\therefore Height of 2nd tree = $x + 13$

$$= 13\sqrt{3} + 13$$

$$= 13(\sqrt{3} + 1)$$


$$= 13 \times 2.732$$

$$= 35.516$$

$$\approx 35.52 \text{ m}$$

2. A man is standing on the deck of a ship, which is 40 m above water level. He observes the angle of elevation of the top of a hill as 60° and the angle of depression of the base of the hill as 30° . Calculate the distance of the hill from the ship and the height of the hill. ($\sqrt{3} = 1.732$)

Solution :



$D \rightarrow$ Deck of a ship

$DS = 40\text{m} = \text{Deck of ship from water level}$

$HL = \text{Height of the hill} = x + 40$

$SL = DB = \text{Dist. between ship \& hill}$

In $\triangle DLS$,

$$\tan 30^\circ = \frac{DS}{SL}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{40}{y}$$

$$\Rightarrow y = 40\sqrt{3} \quad \dots\dots\dots (1)$$

In $\triangle DHB$,

$$\tan 60^\circ = \frac{HB}{DB}$$

$$\Rightarrow \sqrt{3} = \frac{x}{y}$$

$$\Rightarrow y = \frac{x}{\sqrt{3}} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\frac{x}{\sqrt{3}} = 40\sqrt{3}$$

$$\Rightarrow x = 40\sqrt{3} \times \sqrt{3}$$

$$= 120\text{m}$$

\therefore Height of the hill = $x + 40$

$$= 120 + 40$$

$$= 160 \text{ m and}$$

$$y = 40\sqrt{3}$$


$$= 40 (1.732)$$

$$= 69.28 \text{ m}$$

3. If the angle of elevation of a cloud from a point 'h' metres above a lake is θ_1 and the angle of depression of its reflection in the lake is θ_2 . Prove that the height that the cloud is located from the ground is

$$\frac{h(\tan\theta_1 + \tan\theta_2)}{\tan\theta_2 + \tan\theta_1}$$

Solution :



LE \rightarrow Surface of the lake

P \rightarrow Point of observation 'h' mrs. from lake

A, A' \rightarrow Positions of cloud & its reflection

AE = A'E, PL = CE = h m

To find :

$$EA = \frac{h(\tan\theta_1 + \tan\theta_2)}{\tan\theta_2 - \tan\theta_1}$$

In $\triangle APC$,

$$\tan\theta_1 = \frac{x}{y}$$

$$\Rightarrow y = \frac{x}{\tan\theta_1} \quad \dots\dots\dots (1)$$

In $\triangle PA'C$,

$$\tan\theta_2 = \frac{CA'}{PC}$$

$$= \frac{x + 2h}{y}$$

$$\Rightarrow y = \frac{(x + 2h)}{\tan\theta_2} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\frac{x}{\tan\theta_1} = \frac{x + 2h}{\tan\theta_2}$$

$$\Rightarrow x \tan\theta_2 = x \tan\theta_1 + 2h \tan\theta_1$$

$$\Rightarrow x (\tan\theta_2 - \tan\theta_1) = 2h \tan\theta_1$$

$$\Rightarrow x = \frac{2h \tan\theta_1}{\tan\theta_2 - \tan\theta_1}$$

$$\therefore AE = h + x$$

$$= h + \frac{2h \tan\theta_1}{\tan\theta_2 - \tan\theta_1}$$


$$= h \left[1 + \frac{2 \tan\theta_1}{\tan\theta_2 - \tan\theta_1} \right]$$

$$= h \left[\frac{\tan\theta_2 + \tan\theta_1}{\tan\theta_2 - \tan\theta_1} \right]$$

Hence proved.

4. The angle of elevation of the top of a cell phone tower from the foot of a high apartment is 60° and the angle of depression of the foot of the tower from the top of the apartment is 30° . If the height of the apartment is 50 m, find the height of the cell phone tower. According to radiation control norms, the minimum height of a cell phone tower should be 120 m. State if the height of the above mentioned cell phone tower meets the radiation norms.

Solution :



$CD = 50\text{ m} = \text{height of the apartment} = EB$

$AB = (x + 50)\text{ m} = \text{height of the cellphone tower}$

$BD = EC = y\text{ m} = \text{dist. between tower \& apartment}$

In $\triangle CDB$,

$$\tan 30^\circ = \frac{CD}{BD}$$

$$\frac{1}{\sqrt{3}} = \frac{50}{y}$$

$$\Rightarrow y = 50\sqrt{3}\text{ m} \quad \dots\dots\dots (1)$$

In $\triangle ADB$,

$$\tan 60^\circ = \frac{AB}{BD}$$

$$\sqrt{3} = \frac{x+50}{y}$$

$$\Rightarrow y = \frac{x+50}{\sqrt{3}}$$

$$\Rightarrow 50\sqrt{3} = \frac{x+50}{\sqrt{3}} \quad (\text{from (1)})$$

$$\Rightarrow 150 = x+50$$

$$\therefore x = 100\text{ m}$$

\therefore Height of cell phone tower

$$= x + 50$$

$$= 150\text{ m} > 120\text{ m}$$

\therefore The tower does not meet the radiation norms.


5. The angles of elevation and depression of the top and bottom of a lamp post from the top of a 66 m high apartment are 60° and 30° respectively. Find

(i) The height of the lamp post.

(ii) The difference between height of the lamp post and the apartment.

(iii) The distance between the lamp post and the apartment. ($\sqrt{3} = 1.732$)

Solution :



$AT = \text{height of apartment} = 66\text{ m} = EP$

$LP = \text{height of the lamp post} = x + 66$

$PT = EA = y\text{ m} = \text{dist. between post and apartment}$

In $\triangle APT$,

$$\tan 30^\circ = \frac{AT}{y}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{66}{y}$$

$$\Rightarrow y = 66\sqrt{3} \quad \dots\dots\dots (1)$$

In $\triangle ALE$,

$$\tan 60^\circ = \frac{LE}{EA}$$

$$\Rightarrow \sqrt{3} = \frac{x}{y}$$

$$\Rightarrow y = \frac{x}{\sqrt{3}} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\frac{x}{\sqrt{3}} = 66\sqrt{3}$$

$$\Rightarrow x = 66 \times 3$$

$$= 198$$

i) Height of the lamp post = $x + 66$

$$= 198 + 66$$

$$= 264 \text{ mrs}$$

ii) Difference between height of lamp post & apartment = $264 - 66 = 198 \text{ m}$

iii) Dist. between lamp post & the apartment

$$\Rightarrow y = 66\sqrt{3} \text{ m}$$


$$= 66(1.732)$$

$$= 114.312 \text{ m}$$


6. Three villagers A, B and C can see each other across a valley. The horizontal distance between A and B is 8 km and the horizontal distance between B and C is 12 km. The angle of depression of B from A is 20° and the angle of elevation of C from B is 30° . Calculate :

(i) the vertical height between A and B.

(ii) the vertical height between B and C.
($\tan 20^\circ = 0.3640$, ($\sqrt{3} = 1.732$))



Solution :



A, B, C \rightarrow Positions of 3 villagers

To find i) AD ii) CE

In $\triangle ABD$,

$$\tan 20^\circ = \frac{x}{8}$$

$$\Rightarrow x = 8 \cdot \tan 20^\circ$$

$$= 8(0.3640)$$

$$= 2.912$$

$$\therefore AD = 2.912 \text{ km}$$

In $\triangle CBE$,

$$\tan 30^\circ = \frac{y}{12}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{y}{12} \Rightarrow y = \frac{12}{\sqrt{3}}$$

$$= 4\sqrt{3}$$

$$= 4(1.732)$$

$$= 6.928$$

$$\approx 6.93$$

$$\therefore CE = 6.93 \text{ km}$$

EXERCISE 6.5**Multiple choice questions**

1. The value of $\sin^2 \theta + \frac{1}{1 + \tan^2 \theta}$ is equal to

- (1) $\tan^2 \theta$ (2) 1 (3) $\cot^2 \theta$
 (4) 0

Hint :

Ans : (2)

$$\begin{aligned} &= \sin^2 \theta + \frac{1}{1 + \tan^2 \theta} \\ &= \sin^2 \theta + \frac{1}{\sec^2 \theta} \\ &= \sin^2 \theta + \cos^2 \theta \\ &= 1 \end{aligned}$$

2. $\tan \theta \operatorname{cosec}^2 \theta - \tan \theta$ is equal to

- (3) $\sin \theta$ (4) $\cot \theta$

Hint :

Ans : (4)

$$\begin{aligned} &= \tan \theta \cdot \operatorname{cosec}^2 \theta - \tan \theta \\ &= \tan \theta (\operatorname{cosec}^2 \theta - 1) \\ &= \tan \theta \cdot \cot^2 \theta \\ &= \frac{1}{\cot \theta} \times \cot^2 \theta \\ &= \cot \theta \end{aligned}$$

3. If $(\sin \alpha + \operatorname{cosec} \alpha)^2 + (\cos \alpha + \sec \alpha)^2 = k + \tan^2 \alpha + \cot^2 \alpha$, then the value of k is equal to

- (1) 9 (2) 7 (3) 5 (4) 3

Hint :

Ans : (2)

$$\begin{aligned} &= (\sin \alpha + \operatorname{cosec} \alpha)^2 + (\cos \alpha + \sec \alpha)^2 \\ &= k + \tan^2 \alpha + \cot^2 \alpha \\ \Rightarrow &\sin^2 \alpha + \operatorname{cosec}^2 \alpha + 2 \sin \alpha \cdot \operatorname{cosec} \alpha \\ &\quad + \cos^2 \alpha + \sec^2 \alpha + 2 \cos \alpha \sec \alpha \end{aligned}$$

$$= k + \tan^2 \alpha + \cot^2 \alpha$$

$$\Rightarrow 1 + 2 + 2 + \operatorname{cosec}^2 \alpha + \sec^2 \alpha$$

$$= k + \tan^2 \alpha + \cot^2 \alpha$$

$$\Rightarrow 5 + 1 + \cot^2 \alpha + 1 + \tan^2 \alpha$$

$$= k + \tan^2 \alpha + \cot^2 \alpha$$

$$\Rightarrow 7 + \cot^2 \alpha + \tan^2 \alpha = k + \tan^2 \alpha + \cot^2 \alpha$$

$$\therefore k = 7$$

4. If $\sin \theta + \cos \theta = a$ and $\sec \theta + \operatorname{cosec} \theta = b$, then the value of b ($a^2 - 1$) is equal to

- (1) 2a (2) 3a (3) 0
 (4) 2ab

Hint :

Ans : (1)

$$\begin{aligned} b(a^2 - 1) &= (\sec \theta + \operatorname{cosec} \theta) [(\sin \theta + \cos \theta)^2 - 1] \\ &= \left(\frac{1}{\cos \theta} + \frac{1}{\sin \theta} \right) [2 \sin \theta \cos \theta] \\ &= 2 \sin \theta + 2 \cos \theta \\ &= 2 (\sin \theta + \cos \theta) \\ &= 2a \end{aligned}$$

5. If $5x = \sec \theta$ and $\frac{5}{x} = \tan \theta$, then $x^2 - \frac{1}{x^2}$ is equal to

- (1) 25 (2) $\frac{1}{25}$ (3) 5 (4) 1

Hint :

Ans : (2)

$$\begin{aligned} 5x &= \sec \theta, \quad \frac{5}{x} = \tan \theta \\ \Rightarrow \sec^2 \theta - \tan^2 \theta &= 1 \\ \Rightarrow 25x^2 - \frac{25}{x^2} &= 1 \\ \Rightarrow 25 \left(x^2 - \frac{1}{x^2} \right) &= 1 \\ \Rightarrow x^2 - \frac{1}{x^2} &= \frac{1}{25} \end{aligned}$$

6. If $\sin \theta = \cos \theta$, then $2 \tan^2 \theta + \sin^2 \theta - 1$ is equal to

(1) $\frac{-3}{2}$ (2) $\frac{3}{2}$ (3) $\frac{2}{3}$ (4) $\frac{-2}{3}$

Hint :

Ans : (2)

$$\text{Given } \sin \theta = \cos \theta \Rightarrow \theta = 45^\circ$$

$$\therefore 2 \tan^2 \theta + \sin^2 \theta - 1$$

$$= 2 \tan^2 45^\circ + \sin^2 45^\circ - 1$$

$$= 2(1) + \left(\frac{1}{\sqrt{2}}\right)^2 - 1$$

$$= 2 + \frac{1}{2} - 1$$

$$= \frac{3}{2}$$

7. If $x = a \tan \theta$ and $y = b \sec \theta$ then

(1) $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$ (2) $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

(3) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ (4) $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 0$

Hint :

Ans : (1)

$$x = a \tan \theta, \quad y = b \sec \theta$$

$$\therefore \tan \theta = \frac{x}{a}, \quad \sec \theta = \frac{y}{b}$$

$$\sec^2 \theta - \tan^2 \theta = 1$$

$$\Rightarrow \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$$

8. $(1 + \tan \theta + \sec \theta)(1 + \cot \theta - \operatorname{cosec} \theta)$ is equal to

(1) 0 (2) 1 (3) 2 (4) -1

Hint :

Ans : (3)

$$\begin{aligned} &= (1 + \tan \theta + \sec \theta)(1 + \cot \theta - \operatorname{cosec} \theta) \\ &= \left(1 + \frac{\sin \theta}{\cos \theta} + \frac{1}{\cos \theta}\right) \cdot \left(1 + \frac{\cos \theta}{\sin \theta} - \frac{1}{\sin \theta}\right) \\ &= \left(\frac{(\cos \theta + \sin \theta) + 1}{\cos \theta}\right) \left(\frac{(\sin \theta + \cos \theta) - 1}{\sin \theta}\right) \\ &= \left(\frac{(\cos \theta + \sin \theta)^2 - 1}{\cos \theta \cdot \sin \theta}\right) = \frac{2 \sin \theta \cos \theta}{\cos \theta \sin \theta} \\ &= 2 \end{aligned}$$

9. $a \cot \theta + b \operatorname{cosec} \theta = p$ and $b \cot \theta + a \operatorname{cosec} \theta = q$ then $p^2 - q^2$ is equal to

(1) $a^2 - b^2$ (2) $b^2 - a^2$
(3) $a^2 + b^2$ (4) $b - a$

Hint :

Ans : (2)


$$\begin{aligned} p^2 - q^2 &= (a \cot \theta + b \operatorname{cosec} \theta)^2 - (b \cot \theta + a \operatorname{cosec} \theta)^2 \\ &= a^2 \cot^2 \theta + b^2 \operatorname{cosec}^2 \theta + 2ab \cot \theta \operatorname{cosec} \theta \\ &\quad - [b^2 \cot^2 \theta + a^2 \operatorname{cosec}^2 \theta + 2ab \cot \theta \operatorname{cosec} \theta] \\ &= a^2 (\cot^2 \theta - \operatorname{cosec}^2 \theta) + b^2 (\operatorname{cosec}^2 \theta - \cot^2 \theta) \\ &= a^2 (-1) + b^2 (1) \\ &= b^2 - a^2 \end{aligned}$$

10. If the ratio of the height of a tower and the length of its shadow is $\sqrt{3} : 1$, then the angle of elevation of the sun has measure

(1) 45° (2) 30° (3) 90°
(4) 60°

Hint :

Ans : (4)




$$\tan \theta = \frac{h}{s} = \frac{\sqrt{3}}{1} = \sqrt{3} \Rightarrow \theta = 60^\circ$$

11. The electric pole subtends an angle of 30° at a point on the same level as its foot. At a second point 'b' metres above the first, the depression of the foot of the tower is 60° . The height of the tower (in metres) is equal to

(1) $\sqrt{3}b$ (2) $\frac{b}{a}$ (3) $\frac{b}{2}$ (4) $\frac{b}{\sqrt{3}}$

Hint :

Ans : (4)



$$\begin{aligned} \tan 30^\circ &= \frac{b}{y} & \tan 60^\circ &= \frac{x+b}{y} \\ \frac{1}{\sqrt{3}} &= \frac{b}{y} & \sqrt{3} &= \frac{x+b}{y} \\ \Rightarrow y &= \sqrt{3}b & \Rightarrow y &= \frac{x+b}{\sqrt{3}} \end{aligned}$$

$$\therefore \sqrt{3}(x+b) = \frac{b}{\sqrt{3}}$$

$$\Rightarrow 3(x+b) = b$$

$$\Rightarrow b+x = \frac{b}{3}$$


$$\Rightarrow \text{height of tower} = \frac{b}{3} \text{ mts}$$

12. A tower is 60 m height. Its shadow is x metres shorter when the sun's altitude is 45° than when it has been 30° , then x is equal to

(1) 41.92 m (2) 43.92 m
(3) 43 m (4) 45.6 m

Hint :

Ans : (3)



In $\triangle ABC$,

$$\tan 45^\circ = \frac{AB}{BC} = \frac{60}{y}$$

$$\Rightarrow 1 = \frac{60}{y}$$

$$\Rightarrow y = 60$$

In $\triangle ABD$,

$$\tan 30^\circ = \frac{AB}{BD}$$

$$\frac{1}{\sqrt{3}} = \frac{60}{x+y}$$

$$\Rightarrow x+y = 60\sqrt{3}$$

$$\Rightarrow x+60 = 60\sqrt{3}$$

$$\Rightarrow x = 60\sqrt{3} - 60$$

$$= 60(\sqrt{3} - 1)$$

$$= 60 \times 0.732$$


$$= 43.92 \text{ m}$$

13. The angle of depression of the top and bottom of 20 m tall building from the top of a multistoried building are 30° and 60° respectively. The height of the multistoried building and the distance between two buildings (in metres) is

(1) 20, $10\sqrt{3}$ (2) 30, $5\sqrt{3}$
(3) 20, 10 (4) 30, $10\sqrt{3}$

Hint :

Ans : (4)



In $\triangle ACE$,

$$\tan 30^\circ = \frac{AE}{EC}$$

$$\frac{1}{\sqrt{3}} = \frac{x}{y}$$

$$\Rightarrow y = \sqrt{3}x \quad \dots\dots\dots (1)$$

In $\triangle ADB$,

$$\frac{AB}{BD}$$

$$\Rightarrow \sqrt{3} = \frac{x+20}{y}$$

$$\Rightarrow y = \frac{x+20}{\sqrt{3}} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\sqrt{3}x = \frac{x+20}{\sqrt{3}}$$

$$\Rightarrow 3x = x+20$$

$$\Rightarrow 2x = 20$$

$$\therefore x = 10$$

\therefore Height of multistoried building

$$= x + 20$$

$$= 10 + 20$$

$$= 30 \text{ m}$$

\therefore Distance between 2 buildings


$$\begin{aligned} y &= \frac{x+20}{\sqrt{3}} = \frac{30}{\sqrt{3}} = 10\sqrt{3} \\ &= 10(1.732) \\ &= 17.32 \text{ m} \end{aligned}$$

14. Two persons are standing 'x' metres apart from each other and the height of the first person is double that of the other. If from the middle point of the line joining their feet an observer finds the angular elevations of their tops to be complementary, then the height of the shorter person (in metres) is

- (1) $\sqrt{2}x$ (2) $\frac{x}{2\sqrt{2}}$ (3) $\frac{x}{\sqrt{2}}$ (4) $2x$

Hint :

Ans : ()



$CD = h =$ height of shorter person

$AB = 2h =$ height of taller person

$BD = x$ mtrs. $\Rightarrow BO = OD = \frac{x}{2}$

In $\triangle AOB$,

$$\tan \theta = \frac{2h}{x/2} = \frac{4h}{x} \quad \dots\dots\dots (1)$$

In $\triangle COD$,

$$\tan(90^\circ - \theta) = \frac{h}{x/2}$$

$$\Rightarrow \cot \theta = \frac{h}{x/2} = \frac{2h}{x}$$

$$\Rightarrow \tan \theta = \frac{x}{2h} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\frac{4h}{x} = \frac{x}{2h}$$

$$\Rightarrow 8h^2 = x^2$$

$$\Rightarrow h^2 = \frac{x^2}{8}$$

$$\Rightarrow h = \frac{x}{2\sqrt{2}}$$

\therefore Height of the shorter person = $\frac{x}{\sqrt{2}}$ mrs.


15. The angle of elevation of a cloud from a point h metres above a lake is β . The angle of depression of its reflection in the lake is 45° . The height of location of the cloud from the lake is

(1) $\frac{h(1 + \tan \beta)}{1 - \tan \beta}$ (2) $\frac{h(1 - \tan \beta)}{1 + \tan \beta}$

(3) $h \tan(45^\circ - \beta)$ (4) none of these

Hint :

Ans : (1)



LE \rightarrow Surface of the lake

P \rightarrow Point of observation

PL = h mrs = CE

A, A' \rightarrow Positions of cloud & its reflection

\therefore AE = A'E = $x + h$

In $\triangle APC$,

$$\tan \beta = \frac{x}{y}$$

$$\Rightarrow y = \frac{x}{\tan \beta} \quad \dots\dots\dots (1)$$

In $\triangle A'PC$,

$$\tan 45^\circ = \frac{x + 2h}{y} \Rightarrow \frac{x + 2h}{y} = 1$$

$$\Rightarrow y = x + 2h \quad \dots\dots\dots (2)$$

\therefore From (1) & (2),

$$x + 2h = \frac{x}{\tan \beta}$$

$$\Rightarrow 2h = \frac{x}{\tan \beta} - x$$

$$\Rightarrow 2h = x \left(\frac{1}{\tan \beta} - 1 \right)$$

$$\Rightarrow 2h = x \left(\frac{1 - \tan \beta}{\tan \beta} \right)$$

$$x = \frac{2h \tan \beta}{1 - \tan \beta}$$

\therefore Height of the cloud = $h + x$

$$= h + \frac{2h \tan \beta}{1 - \tan \beta}$$

$$= h \left[1 + \frac{2 \tan \beta}{1 - \tan \beta} \right]$$

$$= h \left[\frac{1 + \tan \beta}{1 - \tan \beta} \right]$$

UNIT EXERCISE - 6

1. Prove that

$$(i) \cot^2 A \left(\frac{\sec A - 1}{1 + \sin A} \right) + \sec^2 A \left(\frac{\sin A - 1}{1 + \sec A} \right) = 0$$

$$(ii) \frac{\tan^2 \theta - 1}{\tan^2 \theta + 1} = 1 - 2 \cos^2 \theta$$

Solution :**Solution :**

i) LHS

$$\begin{aligned} & \cot^2 A \left(\frac{\sec A - 1}{1 + \sin A} \right) + \sec^2 A \left(\frac{\sin A - 1}{1 + \sec A} \right) \\ &= \frac{1}{\tan^2 A} \left(\frac{\sec A - 1}{1 + \sin A} \right) - \frac{1}{\cos^2 A} \left(\frac{1 - \sin A}{1 + \sec A} \right) \\ &= \frac{1}{(\sec A + 1) \cancel{(\sec A - 1)}} \cdot \left(\frac{\cancel{\sec A - 1}}{1 + \sin A} \right) - \\ & \quad \frac{1}{(1 + \sin A) \cancel{(1 - \sin A)}} \left(\frac{\cancel{1 - \sin A}}{1 + \sec A} \right) \\ &= \frac{1}{(\sec A + 1)(1 + \sin A)} - \frac{1}{(1 + \sin A)(1 + \sec A)} \\ &= 0 \\ &= \text{RHS} \end{aligned}$$

ii) LHS

$$\begin{aligned} &= \frac{\tan^2 \theta - 1}{\tan^2 \theta + 1} \\ &= \frac{\tan^2 \theta - 1}{\sec^2 \theta} \\ &= (\tan^2 \theta - 1) \cos^2 \theta \\ &= \left(\frac{\sin^2 \theta}{\cos^2 \theta} - 1 \right) \cos^2 \theta \\ &= \sin^2 \theta - \cos^2 \theta \\ &= 1 - \cos^2 \theta - \cos^2 \theta \\ &= 1 - 2 \cos^2 \theta \\ &= \text{RHS} \end{aligned}$$

2. Prove that

$$\left(\frac{1 + \sin \theta - \cos \theta}{1 + \sin \theta + \cos \theta} \right)^2 = \frac{1 - \cos \theta}{1 + \cos \theta}$$

Solution :

LHS

$$= \left(\frac{1 + \sin \theta - \cos \theta}{1 + \sin \theta + \cos \theta} \right)^2$$

Take $1 + \sin \theta = a$, $\cos \theta = b$

$$\begin{aligned} &\therefore \frac{(a - b)^2}{(a + b)^2} \\ &= \frac{a^2 + b^2 - 2ab}{a^2 + b^2 + 2ab} \\ &= \frac{(1 + \sin \theta)^2 + \cos^2 \theta - 2(1 + \sin \theta) \cos \theta}{(1 + \sin \theta)^2 + \cos^2 \theta + 2(1 + \sin \theta) \cos \theta} \\ &= \frac{1 + \sin^2 \theta + 2 \sin \theta + \cos^2 \theta - 2(1 + \sin \theta) \cos \theta}{1 + \sin^2 \theta + 2 \sin \theta + \cos^2 \theta + 2(1 + \sin \theta) \cos \theta} \\ &= \frac{2 + 2 \sin \theta - 2(1 + \sin \theta) \cos \theta}{2 + 2 \sin \theta + 2(1 + \sin \theta) \cos \theta} \\ &= \frac{2(1 + \sin \theta) - 2(1 + \sin \theta) \cos \theta}{2(1 + \sin \theta) + 2(1 + \sin \theta) \cos \theta} \\ &= \frac{\cancel{2(1 + \sin \theta)} [1 - \cos \theta]}{\cancel{2(1 + \sin \theta)} [1 + \cos \theta]} \\ &= \frac{1 - \cos \theta}{1 + \cos \theta} \end{aligned}$$

= RHS

Hence proved.

3. If $x \sin^3 \theta + y \cos^3 \theta = \sin \theta \cos \theta$ and $x \sin \theta = y \cos \theta$, then prove that $x^2 + y^2 = 1$.**Solution :**Given $x \sin^3 \theta + y \cos^3 \theta = \sin \theta \cos \theta$,

$$\Rightarrow x \sin \theta (\sin^2 \theta) + y \cos \theta (\cos^2 \theta)$$

$$= \sin \theta \cos \theta$$

$$\Rightarrow x \sin \theta (\sin^2 \theta) + x \sin \theta (\cos^2 \theta)$$

$$= \sin \theta \cos \theta \text{ (given } x \sin \theta = y \cos \theta \text{)}$$

$$\Rightarrow x \sin \theta (\sin^2 \theta + \cos^2 \theta) = \sin \theta \cos \theta$$

$$\Rightarrow x = \cos \theta$$

$$\text{Also given } x \sin \theta = y \cos \theta$$

$$\Rightarrow \cos \theta \cdot \sin \theta = y \cos \theta$$

$$\Rightarrow y = \sin \theta$$

$$\therefore x^2 + y^2 = \cos^2 \theta + \sin^2 \theta \\ = 1$$

Hence proved.

4. If $a \cos \theta - b \sin \theta = c$, then prove that
 $(a \sin \theta + b \cos \theta) = \pm \sqrt{a^2 + b^2 - c^2}$

Solution :

$$\text{Given } a \cos \theta - b \sin \theta = c$$

Squaring on both sides

$$(a \cos \theta - b \sin \theta)^2 = c^2$$

$$\Rightarrow a^2 (1 - \sin^2 \theta) + b^2 (1 - \cos^2 \theta) - 2ab \cos \theta \sin \theta = c^2$$

$$\Rightarrow a^2 - a^2 \sin^2 \theta + b^2 - b^2 \cos^2 \theta - 2ab \cos \theta \sin \theta = c^2$$

$$\Rightarrow a^2 \sin^2 \theta + b^2 \cos^2 \theta + 2ab \cos \theta \sin \theta + a^2 + b^2 - c^2$$


$$(a \sin \theta + b \cos \theta)^2 = a^2 + b^2 - c^2$$

$$\therefore a \sin \theta + b \cos \theta = \pm \sqrt{a^2 + b^2 - c^2}$$

Hence proved.

5. A bird is sitting on the top of a 80 m high tree. From a point on the ground, the angle of elevation of the bird is 45° . The bird flies away horizontally in such away that it remained at a constant height from the ground. After 2 seconds, the angle of elevation of the bird from the same point is 30° . Determine the speed at which the bird flies. ($\sqrt{3} = 1.732$)

Solution :



P → Point of observation

T, T' → Initial & final positions of the bird

TG = T'G' = 80m = height at which the bird is on the tree, from the ground.

$$\angle QTG = \angle QT'T = 30^\circ$$

In ΔPTG ,

$$\tan 45^\circ = \frac{TG}{PG}$$

$$\Rightarrow 1 = \frac{80}{y}$$

$$\Rightarrow y = 80 \quad \dots\dots\dots (1)$$

In ΔPQG ,

$$\tan 30^\circ = \frac{80-x}{y}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{80-x}{80} \quad (\text{from (1)})$$

$$\Rightarrow 80 = 80\sqrt{3} - \sqrt{3}x$$

$$\Rightarrow \sqrt{3}x = 80(\sqrt{3} - 1)$$

$$\therefore x = \frac{80(\sqrt{3} - 1)}{\sqrt{3}} \quad \dots\dots\dots (2)$$

In $\Delta TQT'$

$$\tan 30^\circ = \frac{TQ}{TT'}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{x}{TT'}$$

$$\Rightarrow TT' = \sqrt{3}x$$


$$\Rightarrow TT' = 80(\sqrt{3} - 1) \quad (\text{from (2)})$$

Given, time taken by the bird from T to reach T' = 2 sec.

$$\begin{aligned}\therefore \text{Speed of the bird} &= \frac{\text{Distance}}{\text{Time}} \\ &= \frac{80(\sqrt{3}-1)}{2} \\ &= 40(\sqrt{3}-1) \\ &= 40 \times 1.732 \\ &= 29.28 \text{ m/sec.}\end{aligned}$$

6. An aeroplane is flying parallel to the Earth's surface at a speed of 175 m/sec and at a height of 600 m. The angle of elevation of the aeroplane from a point on the Earth's surface is 37° at a given point. After what period of time does the angle of elevation increase to 53° ? ($\tan 53^\circ = 1.3270$, $\tan 37^\circ$)

Solution :



P → Point of observation.

A, A' → Initial & final positions of the plane.

Speed of the plane = 175 m/sec.

AG = A'G' = 600 m

= height at which the plane is flying.

In $\triangle PAG$,

$$\tan 37^\circ = \frac{AG}{PG}$$

$$\Rightarrow \tan 37^\circ = \frac{600}{y+x}$$

$$\Rightarrow y+x = \frac{600}{\tan 37^\circ}$$

$$\Rightarrow y = \frac{600}{\tan 37^\circ} - x \quad \dots\dots\dots (1)$$

In $\triangle PA'G'$,

$$\tan 53^\circ = \frac{A'G'}{PG'}$$

$$\Rightarrow \tan 53^\circ = \frac{600}{y}$$

$$\Rightarrow y = \frac{600}{\tan 53^\circ} \quad \dots\dots\dots (2)$$

\therefore From (1) & (2)

$$\frac{600}{\tan 53^\circ} = \frac{600}{\tan 37^\circ} - x$$

$$\Rightarrow x = 600 \left(\frac{1}{\tan 37^\circ} - \frac{1}{\tan 53^\circ} \right)$$

$$\Rightarrow x = 600 \left(\frac{\tan 53^\circ - \tan 37^\circ}{\tan 53^\circ \tan 37^\circ} \right)$$

$$= 600 \left(\frac{\tan 53^\circ - \tan 37^\circ}{\tan 53^\circ \tan 37^\circ} \right) \quad (\because \tan 53^\circ = \cot 53^\circ = 1)$$

$$= 600 \left(\frac{\tan 53^\circ - \tan 37^\circ}{1} \right)$$

$$= 600 (1.3270 - 0.7536)$$

$$= 600 \times 0.5734$$

$$= 344.04 \text{ m}$$

$$\text{Time} = \frac{\text{Distance}}{\text{Speed}}$$

$$= \frac{344.04}{175}$$

$$= 1.9659$$

$$\approx 1.97 \text{ sec.}$$

\therefore After 1.97 sec (approx), angle of elevation changes from 37° to 53° .

7. A bird is flying from A towards B at an angle of 35° , a point 30 km away from A. At B it changes its course of flight and heads towards C on a bearing of 48° and distance 32 km away.

(i) How far is B to the North of A?


(ii) How far is B to the West of A?

(iii) How far is C to the North of B?

(iv) How far is C to the East of B?

($\sin 55^\circ = 0.8192$, $\cos 55^\circ = 0.5736$,
 $\sin 42^\circ = 0.6691$, $\cos 42^\circ = 0.7431$)

Solution :



- i) The distance of B to the north of A [BE]

In $\triangle BCE$,

$$\Rightarrow \cos 48^\circ = \frac{BE}{BC}$$

$$\Rightarrow \sin 42^\circ = \frac{BE}{32}$$

$$\begin{aligned} \Rightarrow BE &= 32 (0.6691) \\ &= 21.4112 \\ &\approx 21.41 \text{ km.} \end{aligned}$$

- ii) The distance of B to the west of A [BD]

In $\triangle BAD$,


$$\Rightarrow \sin 35^\circ = \frac{BD}{30}$$

$$\Rightarrow \cos 55^\circ = \frac{BD}{30} \quad (\because \sin(90 - \theta) = \cos \theta)$$

$$\begin{aligned} \Rightarrow BD &= 30^\circ \cdot \cos 55^\circ \\ &= 30 (0.5736) \\ &= 17.208 \\ &\approx 17.21 \text{ km.} \end{aligned}$$

8. Two ships are sailing in the sea on either side of the lighthouse. The angles of depression of two ships as observed from the top of the lighthouse are 60° and 45° respectively. If the distance between the ships is $200 \left(\frac{\sqrt{3} + 1}{\sqrt{3}} \right)$ metres, find the height of the lighthouse.

Solution :



LH = h m = height of the lighthouse

AB = Dist. between 2 ships =

$$x + y = 200 \left(\frac{\sqrt{3} + 1}{\sqrt{3}} \right)$$

In $\triangle LBH$,

$$\tan 45^\circ = \frac{h}{y}$$

$$\Rightarrow 1 = \frac{h}{y}$$

$$\Rightarrow y = h \quad \dots\dots\dots (1)$$

In $\triangle LAH$,

$$\begin{aligned}\tan 60^\circ &= \frac{h}{x} \\ \Rightarrow \sqrt{3} &= \frac{h}{x} \\ \Rightarrow x &= \frac{h}{\sqrt{3}} \quad \dots\dots\dots (2)\end{aligned}$$

\therefore Adding (1) & (2)


$$\begin{aligned}x + y &= h + \frac{h}{\sqrt{3}} \\ \Rightarrow 200 \left(\frac{\sqrt{3} + 1}{\sqrt{3}} \right) &= h \left(1 + \frac{1}{\sqrt{3}} \right) \\ \Rightarrow \frac{200(\sqrt{3} + 1)}{\sqrt{3}} &= h \left(\frac{\sqrt{3} + 1}{\sqrt{3}} \right) \\ \therefore h &= 200\end{aligned}$$

\therefore Height of the light house = 200 m.

9. A building and a statue are in opposite side of a street from each other 35 m apart. From a point on the roof of building the angle of elevation of the top of statue is 24° and the angle of depression of base of the statue is 34° . Find the height of the statue.

($\tan 24^\circ = 0.4452$, $\tan 34^\circ = 0.6745$)

Solution :



Let $CD = x$ m = height of the building = EB

$AB = x + y$ m = height of the statue

In $\triangle CBD$,

$$\begin{aligned}\tan 34^\circ &= \frac{CD}{BD} \\ \Rightarrow \tan 34^\circ &= \frac{x}{35} \\ x &= 35 \tan 34^\circ \quad \dots\dots\dots (1)\end{aligned}$$

In $\triangle ACE$,

$$\begin{aligned}\tan 24^\circ &= \frac{AE}{EC} \\ \Rightarrow \tan 24^\circ &= \frac{y}{35} \\ y &= 35 \tan 24^\circ \quad \dots\dots\dots (2)\end{aligned}$$

Height of the statue = $x + y$

$$\begin{aligned}&= 35 (0.6745 + 0.4452) \\ &= 35 (1.1197) \\ &= 39.1895 \\ &\approx 39.19 \text{ m}\end{aligned}$$

PROBLEMS FOR PRACTICE

- If $7 \sin^2 \theta + 3 \cos^2 \theta = 4$, show that $\tan \theta = \frac{1}{\sqrt{3}}$.
- Prove that $2(\sin^6 \theta + \cos^6 \theta) - 3(\sin^4 \theta + \cos^4 \theta) = 0$
- Prove that $\frac{\cot A + \operatorname{cosec} A - 1}{\cot A - \operatorname{cosec} A + 1} = \frac{1 + \cos A}{\sin A}$
- If $\tan \theta + \sin \theta = p$, $\tan \theta - \sin \theta = q$, prove that $p^2 - q^2 = 4\sqrt{pq}$
- Prove that $\cot \theta - \tan \theta = \frac{2 \cos^2 \theta - 1}{\sin \theta \cdot \cos \theta}$
- If $x = \sec A + \sin A$, $y = \sec A - \sin A$, prove that $\left(\frac{2}{x+y} \right)^2 + \left(\frac{x-y}{2} \right)^2 = 1$

7. If $\sin \theta = \frac{15}{17}$, find the value of $\frac{3-4\sin^2 \theta}{4\cos^2 \theta - 3}$
 (Ans : $\frac{33}{611}$)
8. If $a = \sin \theta + \cos \theta$, $b = \sin^3 \theta + \cos^3 \theta$, then show that $(3a - 2b) = a^3$
9. If $\sec \theta = x + \frac{1}{4x}$ prove that $\sec \theta + \tan \theta = 2x$
10. Prove that $\frac{\sin \theta}{\cot \theta + \operatorname{cosec} \theta} = 2 + \frac{\sin \theta}{\cot \theta - \operatorname{cosec} \theta}$
11. Prove that $(\tan \alpha + \operatorname{cosec} \beta)^2 - (\cot \beta - \sec \alpha)^2 = 2 \tan \alpha \cdot \cot \beta (\operatorname{cosec} \alpha + \sec \beta)$.
12. If $\sin \theta + \sin^2 \theta = 1$, prove that $\cos^{12} \theta + 3\cos^{10} \theta + 3\cos^8 \theta + \cos^6 \theta + 2\cos^4 \theta + 2\cos^2 \theta - 2 = 0$
13. Two vertical lamp - posts of equal height stand on either side of a roadway which is 50m wide. At a point in the road between lamp posts, elevations of the tops of the lamp posts are 60° and 30° . Find the height of each lamp post. (Ans : 21.65 m)
14. The angle of elevation of a jet plane from a point P on the ground is 60° . After 15 seconds the angle of elevation changes to 30° . If the jet is flying at a speed of 720 km/h, find the height at which the jet is flying. (Ans : 2.6 km)
15. From an aeroplane flying horizontally above a straight road, the angles of depression of two consecutive kilometer stones on the road are 45° and 30° respectively. Find the height of the aeroplane above the road when the km stones are i) on the same side of the vertical through aeroplane
 ii) on the opposite sides.
 (Ans : i) 1.366 km ii) 0.366 km)

Objective Type Questions :

1. If $\cos \theta = \frac{a}{b}$, then $\operatorname{cosec} \theta$ is equal to
 a) $\frac{b}{a}$ b) $\frac{b}{\sqrt{b^2 - a^2}}$
 c) $\frac{\sqrt{b^2 - a^2}}{b}$ d) $\frac{a}{\sqrt{b^2 - a^2}}$
 (Ans : (b))
2. If $\sin \theta - \cos \theta = 0$, then the value of $\sin^4 \theta + \cos^4 \theta$ is
 a) $\frac{1}{2}$ b) $\frac{1}{4}$ c) $\frac{3}{4}$ d) 1
 (Ans : (d))
3. The value of $\frac{11}{\cot^2 \theta} - \frac{11}{\cos^2 \theta}$ is
 a) 11 b) 0 c) $\frac{1}{\cos^2 \theta}$ d) -11
 (Ans : (d))
4. If $\tan \theta + \cot \theta = 5$, then the value of $\tan^2 \theta + \cot^2 \theta$ is
 a) 23 b) 25 c) 27 d) 15
 (Ans : (a))
5. The value of $(1 + \cot^2 \theta) \cdot (1 + \cos \theta) (1 - \cos \theta)$ is
 a) $\sin^2 \theta$ b) $\operatorname{cosec}^2 \theta$ c) 1
 d) $\sec^2 \theta$ (Ans : (c))
6. The value of $\tan 10^\circ \cdot \tan 15^\circ \cdot \tan 75^\circ \tan 80^\circ$ is
 a) -1 b) 0 c) 1 d) 4
 (Ans : (c))
7. A pole 6m high casts a shadow $2\sqrt{3}$ m long on the ground, then the sun's elevation is
 a) 60° b) 45° c) 30° d) 90°
 (Ans : (a))

8. The angle of elevation of the top of a tower from a point situated at a distance of 100m from the base of a tower is 30° . The height of the tower is

a) $\frac{100}{\sqrt{3}}$ m b) $100\sqrt{3}$ m
 c) $\frac{50}{\sqrt{3}}$ m d) $50\sqrt{3}$ m

(Ans : (a))

9. The angle of depression of a point on the horizontal from the top of a hill is 60° . If one has to walk 300m to reach the top from this point, then the distance of this point from the base of the hill is

a) $300\sqrt{3}$ m b) 150 m
 c) $150\sqrt{3}$ m d) $\frac{150}{\sqrt{3}}$ m

(Ans : (b))

10. The value of $\sin\theta \cdot \operatorname{cosec}\theta + \cos\theta \sec\theta$ is

a) 1 b) 2 c) 0 d) $\frac{1}{2}$

(Ans : (b))

11. If $5 \cos \theta = 7 \sin \theta$, then the value of $\frac{7 \sin \theta + 5 \cos \theta}{5 \sin \theta + 7 \cos \theta}$ is

a) $\frac{37}{35}$ b) 1 c) $\frac{5}{7}$ d) $\frac{35}{37}$

(Ans : (d))

12. If $\sin A = \frac{1}{\sqrt{5}}$, then $\sec A$ is

a) $\frac{1}{\sqrt{5}}$ b) $\frac{2}{\sqrt{5}}$ c) $\frac{\sqrt{5}}{2}$ d) $\sqrt{5}$

(Ans : (c))

13. The acute angle ' θ ' when $\sec^2 \theta + \tan^2 \theta = 3$ is

a) 30° b) 45° c) 60° d) 90°

(Ans : (b))

a) 45° b) 30° c) 90°

d) none of these **(Ans: (a))**

15. If $\tan \alpha = \sqrt{3}$, $\tan \beta = \frac{1}{\sqrt{3}}$, then $\cot(\alpha + \beta)$ is

a) $\sqrt{3}$ b) 0 c) $\frac{1}{\sqrt{3}}$ d) 1

(Ans : (b))

CHAPTER 7


MENSURATION

I. SURFACE AREA


Key Points

Right Circular and Hollow Cylinder

- ✓ A right circular cylinder is a solid generated by the revolution of a rectangle about one of its sides as axis.
- ✓ If the axis is perpendicular to the radius then the cylinder is called a right circular cylinder.
- ✓ A solid cylinder is an object bounded by two circular plane surfaces and a curved surface.






- ✓ T.S.A. of a right circular cylinder = $2\pi r(h + r)$ sq. units
- ✓ An object bounded by two co-axial cylinders of the same height and different radii is called a 'hollow cylinder'.
- ✓ C.S.A of a hollow cylinder = $2\pi(R + r)h$ sq. units
- ✓ T.S.A. of a hollow cylinder = $2\pi(R + r)(R - r + h)$ sq. units




Right Circular and Hollow Cone

- ✓ A right circular cone is a solid generated by the revolution of a right angled triangle about one of the sides containing the right angle as axis.
- ✓ If the right triangle ABC revolves about AB as axis, the hypotenuse AC generates the curved surface of the cone.
- ✓ The height of the cone is the length of the axis AB, and the slant height is the length of the hypotenuse AC.
- ✓ Curved Surface Area of the cone = Area of the Sector = $\pi r l$ sq. units.
- ✓ T.S.A. of a right circular cone = $\pi r(l + r)$ sq. units, where $l = \sqrt{h^2 + r^2}$.




Sphere & Hemisphere

- ✓ A sphere is a solid generated by the revolution of a semicircle about its diameter as axis.
- ✓ Surface area of a sphere = $4\pi r^2$ sq.units.
- ✓ A section of the sphere cut by a plane through any of its great circle is a hemisphere.
- ✓ C.S.A. of a hemisphere = $2\pi r^2$ sq.units.
- ✓ T.S.A. of a hemisphere = $3\pi r^2$ sq.units.
- ✓ C.S.A. of a hollow hemisphere = $2\pi(R^2 + r^2)$ sq. units.
- ✓ T.S.A. of a hollow hemisphere = $\pi(3R^2 + r^2)$ sq. units.
- ✓ Thickness = $R - r$



Frustum of a Cone

- ✓ When a cone ABC is cut through by a plane parallel to its base, the portion of the cone DECB between the cutting plane and the base is
- ✓ C.S.A. of a frustum = $\pi(R + r)l$ sq. units, where, $l = \sqrt{h^2 + (R - r)^2}$.
- ✓ T.S.A. of a frustum = $\pi(R + r)l + \pi R^2 + \pi r^2$ sq. units.



Example 7.1

A cylindrical drum has a height of 20 cm and base radius of 14 cm. Find its curved surface area and the total surface area.

Solution :

Given that, height of the cylinder $h = 20$ cm ; radius $r = 14$ cm

Now, C.S.A. of the cylinder = $2\pi rh$ sq. units

C.S.A. of the cylinder

$$= 2 \times \frac{22}{7} \times 14 \times 20 = 2 \times 22 \times 2 \times 20 = 1760 \text{ cm}^2$$

T.S.A. of the cylinder = $2\pi r(h + r)$ sq. units

$$= 2 \times \frac{22}{7} \times 14 \times (20 + 14) = 2 \times \frac{22}{7} \times 14 \times 34 = 2992 \text{ cm}^2$$

Therefore, C.S.A.

$$= 1760 \text{ cm}^2 \text{ and T.S.A.} = 2992 \text{ cm}^2$$

Example 7.2

The curved surface area of a right circular cylinder of height 14 cm is 88 cm^2 . Find the diameter of the cylinder.

Solution :

Given that, C.S.A. of the cylinder = 88 sq. cm

$$2\pi rh = 88$$

$$= 2 \times \frac{22}{7} \times r \times 14 = 88 \quad (\text{given } h = 14 \text{ cm})$$

$$2r = \frac{88 \times 7}{22 \times 14} = 2$$

Therefore, diameter = 2 cm

Example 7.3

A garden roller whose length is 3 m long and whose diameter is 2.8 m is rolled to level a garden. How much area will it cover in 8 revolutions?

Solution :

Given that, diameter $d = 2.8$ m and height = 3 m
radius $r = 1.4$ m



Fig. 7.6

area of the cylinder

$$= 2\pi rh \text{ sq. units}$$

$$= 2 \times \frac{22}{7} \times 1.4 \times 3 = 26.4$$

Area covered in 1 revolution = 26.4 m^2

Area covered in 8 revolutions

$$= 8 \times 26.4 = 211.2$$

Therefore, area covered is 211.2 m^2

Example 7.4

If one litre of paint covers 10 m^2 , how many litres of paint is required to paint the internal and external surface areas of a cylindrical tunnel whose thickness is 2 m, internal radius is 6 m and height is 25 m.

Solution :

Given that, height $h = 25$ m; thickness = 2 m.

Internal radius $r = 6$ m

Now, external radius $R = 6 + 2 = 8$ m

C.S.A. of the cylindrical tunnel

= C.S.A. of the hollow cylinder

C.S.A. of the hollow cylinder

$$= 2\pi (R + r) h \text{ sq. units}$$

$$= 2 \times \frac{22}{7} (8 + 6) \times 25$$

Hence, C.S.A. of the cylindrical tunnel

$$= 2200 \text{ m}^2$$

Area covered by one litre of paint = 10 m^2

Number of litres required to paint the tunnel

$$= \frac{2200}{10} = 220$$

Therefore, 220 litres of paint is needed to paint the tunnel.

Example 7.5

The radius of a conical tent is 7 m and the height is 24 m. Calculate the length of the canvas used to make the tent if the width of the rectangular canvas is 4 m?

Solution :

Let r and h be the radius and height of the cone respectively.

Given that, radius $r = 7$ m and height $h = 24$ m

$$\begin{aligned} \text{Hence, } l &= \sqrt{r^2 + h^2} \\ &= \sqrt{49 + 576} \\ l &= \sqrt{625} = 25 \text{ m} \end{aligned}$$

C.S.A. of the conical tent = πrl sq. units

$$\text{Area of the canvas} = \frac{22}{7} \times 7 \times 25 = 550 \text{ m}^2$$

Now, length of the canvas =

$$\frac{\text{Area of canvas}}{\text{Width}} = \frac{550}{4} = 137.5 \text{ m}$$

Therefore, the length of the canvas is 137.5 m

Example 7.6

If the total surface area of a cone of radius 7 cm is 704 cm^2 , then find its slant height.

Solution :

Given that, radius $r = 7 \text{ cm}$

Now, total surface area of the cone

$$= \pi r (l + r) \text{ sq. units}$$

$$\text{T.S.A.} = 704 \text{ cm}^2$$

$$704 = \frac{22}{7} \times 7 (l + 7)$$

$$32 = l + 7 \text{ implies } l = 25 \text{ cm}$$

Therefore, slant height of the cone is 25 cm.

Example 7.7

From a solid cylinder whose height is 2.4 cm and diameter 1.4 cm, a conical cavity of the same height and base is hollowed out (Fig. 7.13). Find the total surface area of the remaining solid.

Solution :

Let h and r be the height and radius of the cone and cylinder.

Let l be the slant height of the cone.

Given that, $h = 2.4 \text{ cm}$ and $d = 1.4 \text{ cm}$;

$$r = 0.7 \text{ cm}$$




Fig. 7.13

Here, total surface area of the remaining solid }
 = C.S.A. of the cylinder + C.S.A. of the cone
 + area of the bottom

$$= 2\pi rh + \pi rl + \pi r^2 \text{ sq. units}$$

Now,

$$l = \sqrt{r^2 + h^2} = \sqrt{0.49 + 5.76} = \sqrt{6.25} = 2.5 \text{ cm}$$

$$l = 2.5 \text{ cm}$$

Area of the remaining solid

$$= 2\pi rh + \pi rl + \pi r^2 \text{ sq. units}$$

$$= \pi r (2h + l + r)$$

$$= \frac{22}{7} \times 0.7 \times [(2 \times 2.4) + 2.5 + 0.7]$$

Therefore, total surface area of the remaining solid is 17.6 m^2

Example 7.8

Find the diameter of a sphere whose surface area is 154 m^2 .

Solution :

Let r be the radius of the sphere.

Given that, surface area of sphere = 154 m^2

$$4\pi r^2 = 154$$

$$4 \times \frac{22}{7} \times r^2 = 154$$

$$\text{gives } r^2 = 154 \times \frac{1}{4} \times \frac{7}{22}$$

$$\text{hence, } r^2 = \frac{49}{4} \text{ we get } r = \frac{7}{2}$$

Therefore, diameter is 7 m

Example 7.9

The radius of a spherical balloon increases from 12 cm to 16 cm as air being pumped into it. Find the ratio of the surface area of the balloons in the two cases.

Solution :

Let r_1 and r_2 be the radii of the balloons.

Given that,

$$\frac{r_1}{r_2} = \frac{12}{16} = \frac{3}{4}$$

Now, ratio of C.S.A. of balloons =

$$\frac{4\pi r_1^2}{4\pi r_2^2} = \frac{r_1^2}{r_2^2} = \left(\frac{r_1}{r_2}\right)^2 = \left(\frac{3}{4}\right)^2 = \frac{9}{16}$$

Example 7.10

If the base area of a hemispherical solid is 1386 sq. metres, then find its total surface area?

Solution :

Let r be the radius of the hemisphere.

Given that, base area = $\pi r^2 = 1386$ sq. m

$$\begin{aligned} \text{T.S.A.} &= 3\pi r^2 \text{ sq.m} \\ &= 3 \times 1386 = 4158 \end{aligned}$$

Therefore, T.S.A. of the hemispherical solid is 4158 m².

Example 7.11

The internal and external radii of a hollow hemispherical shell are 3 m and 5 m respectively. Find the T.S.A. and C.S.A. of the shell.

Solution :

Let the internal and external radii of the hemispherical shell be r and R respectively.




Fig. 7.19

Given that, $R = 5$ m, $r = 3$ m

C.S.A. of the shell = $2\pi(R^2 + r^2)$ sq. units

$$= 2 \times \frac{22}{7} \times (25 + 9) = 213.71$$

T.S.A. of the shell = $\pi(3R^2 + r^2)$ sq. units

$$= \frac{22}{7} (75 + 9) = 264$$

Therefore, C.S.A. = 213.71 m² and

$$\text{T.S.A.} = 264 \text{ m}^2.$$

Example 7.12

A sphere, a cylinder and a cone (Fig.7.20) are of the same radius, where as cone and cylinder are of same height. Find the ratio of their curved surface areas.


Solution :

Fig. 7.20

Required Ratio = C.S.A. of the sphere: C.S.A. of the cylinder : C.S.A. of the cone

$$\begin{aligned} &= 4\pi r^2 : 2\pi rh : \pi rl, \\ &\quad (l = \sqrt{r^2 + h^2} = \sqrt{2r^2} = \sqrt{2}r \text{ units}) \\ &= 4 : 2 : \sqrt{2} = 2\sqrt{2} : \sqrt{2} : 1 \end{aligned}$$

Example 7.13

The slant height of a frustum of a cone is 5 cm and the radii of its ends are 4 cm and 1 cm. Find its curved surface area.

Solution :

Let l , R and r be the slant height, top radius and bottom radius of the frustum.

Given that, $l = 5$ cm, $R = 4$ cm, $r = 1$ cm

Now, C.S.A. of the frustum

$$\begin{aligned} &= \pi(R + r) l \text{ sq. units} \\ &= \frac{22}{7} \times (4 + 1) \times 5 \\ &= \frac{550}{7} \end{aligned}$$

Therefore, C.S.A. = 78.57 cm^2

Example 7.14

An industrial metallic bucket is in the shape of the frustum of a right circular cone whose top and bottom diameters are 10 m and 4 m and whose height is 4 m. Find the curved and total surface area of the bucket.

Solution :

Let h , l , R and r be the height, slant height, outer radius and inner radius of the frustum.




Fig. 7.24

Given that, diameter of the top = 10 m; radius of the top $R = 5$ m. diameter of the bottom = 4 m; radius of the bottom $r = 2$ m, height $h = 4$ m

$$\begin{aligned} \text{Now, } l &= \sqrt{h^2 + (R - r)^2} \\ &= \sqrt{4^2 + (5 - 2)^2} \\ l &= \sqrt{16 + 9} = \sqrt{25} = 5 \text{ m} \end{aligned}$$

Here, C.S.A = $\pi(R + r)l$ sq. units

$$= \frac{22}{7} (5 + 2) \times 5 = 110 \text{ m}^2$$

T.S.A. = $\pi(R + r)l + \pi R^2 + \pi r^2$ sq. units

$$= \frac{22}{7} [(5 + 2) 5 + 25 + 4]$$

$$= \frac{1408}{7} = 201.14$$

Therefore, C.S.A. = 110 m^2 and

2

EXERCISE 7.1

1. The radius and height of a cylinder are in the ratio 5 : 7 and its curved surface area is 5500 sq.cm. Find its radius and height.

Solution:

Given $r : h = 5 : 7$

$$\begin{aligned} \Rightarrow \frac{r}{h} &= \frac{5}{7} \\ \Rightarrow 7r &= 5h \\ \Rightarrow h &= \frac{7r}{5} \end{aligned}$$

CSA of Cylinder = 5500

$$\Rightarrow 2\pi rh = 5500$$

$$\Rightarrow 2 \times \frac{22}{7} \times r \times \frac{7r}{5} = 5500$$

$$\Rightarrow \frac{44}{5} r^2 = 5500$$

$$\Rightarrow r^2 = \frac{5500 \times 5}{44}$$

$$\Rightarrow r = \frac{500 \times 5}{4}$$

$$= 125 \times 5$$

$$= 625$$

$$\therefore r = 25$$

$$\therefore h = \frac{7}{5} \times 25 = 35$$

\therefore Radius = 25 cm, Height = 35 cm

2. A solid iron cylinder has total surface area of 1848 sq.m. Its curved surface area is five – sixth of its total surface area. Find the radius and height of the iron cylinder.

Solution :

$$= 1848 \text{ m}^2 \text{ \&}$$

$$\text{CSA} = \frac{5}{6} (\text{TSA})$$

$$\Rightarrow 2\pi rh = \frac{5}{6} \times 1848$$

$$= 5 \times 308$$

$$2\pi rh = 1540 \quad \dots\dots\dots (1)$$

$$2\pi r (h + r) = 1848$$

$$\Rightarrow 2\pi rh + 2\pi r^2 = 1848$$

$$\Rightarrow 1540 + 2\pi r^2 = 1848$$

$$\Rightarrow 2\pi r^2 = 308$$

$$\Rightarrow 2 \times \frac{22}{7} \times r^2 = 308$$

$$\Rightarrow r^2 = \frac{\overset{14}{\cancel{308}} \times 7}{\cancel{2} \times \underset{1}{22}}$$

$$\Rightarrow r^2 = 49$$

$$r = 7 \text{ m}$$

Sub $r = 7$ in (1)

$$2 \times \frac{22}{7} \times 7 \times h = 1540$$

$$\Rightarrow h = \frac{1540}{2 \times 22}$$

$$h = 35$$

\therefore Radius = 7 m, Height = 35 m.

3. The external radius and the length of a hollow wooden log are 16 cm and 13 cm respectively. If its thickness is 4 cm then find its T.S.A.

Solution :

Given, external radius of

hollow cylinder = 16 cm = R

length of log = 13 cm = R

thickness = 4 cm = t

$$\therefore t = R - r$$

$$\therefore r = R - t$$

$$= 16 - 4$$

$$r = 12$$

\therefore TSA of hollow cylinder


$$= 2\pi (R + r) (R - r + h)$$

$$= 2 \times \frac{22}{7} (28) (4 + 13)$$

$$= 44 \times 4 \times 17$$

$$= 2992 \text{ cm}^2$$

4. A right angled triangle PQR where $\angle Q = 90^\circ$ is rotated about QR and PQ. If QR = 16 cm and PR = 20 cm, compare the curved surface areas of the right circular cones so formed by the triangle.

Solution :

$$\begin{aligned}\therefore PQ &= \sqrt{400 - 256} \\ &= \sqrt{144} = 12\end{aligned}$$

- i) When the Δ is rotated about PQ,

$$h = 12 \text{ cm}, r = 16 \text{ cm}$$

$$\begin{aligned}\therefore \text{CSA of cone} &= \pi r l \\ &= \pi \times 16 \times 20 \\ &= 320\pi \text{ cm}^2\end{aligned}$$

$$h = 16 \text{ cm}, r = 12 \text{ cm}$$

$$\begin{aligned}\therefore \text{CSA of cone} &= \pi r l \\ &= \pi \times 16 \times 12 \\ &= 192\pi \text{ cm}^2\end{aligned}$$

\therefore CSA of the cone when rotated about PQ is larger than that of QR.

5. 4 persons live in a conical tent whose slant height is 19 cm. If each person require 22 cm² of the floor area, then find the height of the tent.

Solution :

Given slant height of the cone $l = 19$ cm

Total floor area of 4 persons = 88 cm²

$$\Rightarrow \pi r^2 = 88$$

$$\Rightarrow \frac{22}{7} \times r^2 = 88$$

$$\Rightarrow r^2 = 28$$

$$\begin{aligned}\therefore h &= \sqrt{l^2 - r^2} \\ &= \sqrt{19^2 - 28} \\ &= \sqrt{361 - 28} \\ &= \sqrt{333}\end{aligned}$$

height of cone $\simeq 18.25$ cm.

6. A girl wishes to prepare birthday caps in the form of right circular cones for her birthday party, using a sheet of paper whose area is 5720 cm², how many caps can be made with radius 5 cm and height 12 cm.

Solution :

Given $r = 5$ cm, $h = 12$ cm in a cone

$$\begin{aligned}\therefore l &= \sqrt{h^2 + r^2} \\ &= \sqrt{144 + 25} \\ &= \sqrt{169} \\ &= 13\end{aligned}$$

$$\begin{aligned}\therefore \text{CSA of cone} &= \pi r l \\ &= \frac{22}{7} \times 5 \times 13 \\ &= \frac{110 \times 13}{7} \text{ cm}^2\end{aligned}$$

Given, area of sheet of paper = 5720 cm²

$$\begin{aligned}\therefore \text{Number of caps} &= \frac{5720 \times 7}{110 \times 3} \\ &= 28 \text{ caps}\end{aligned}$$

7. The ratio of the radii of two right circular cones of same height is 1 : 3. Find the ratio of their curved surface area when the height of each cone is 3 times the radius of the smaller cone.

Solution :

$$\text{Given } r_1 : r_2 = 1 : 3 \quad h_1 = 3r_1, h_2 = 3r_1$$

$$\text{Let } r_1 = x \quad \quad \quad = 3x \quad , \quad = 3x$$

$$r_2 = 3x$$

$$l_1 = \sqrt{h_1^2 + r_1^2}, \quad l_2 = \sqrt{h_2^2 + r_2^2}$$

$$= \sqrt{9x^2 + x^2} \quad = \sqrt{9x^2 + 9x^2}$$

$$= \sqrt{10x} \quad = 3\sqrt{2}x$$

∴ Ratio of their CSA

$$= \frac{\pi r_1 l_1}{\pi r_2 l_2}$$

$$= \frac{1}{3} \times \frac{\sqrt{10}}{3\sqrt{2}} = \frac{\sqrt{5}}{9}$$

$$\therefore \text{Ratio of their CSA} = \sqrt{5} : 9$$

- 8. The radius of a sphere increases by 25%. Find the percentage increase in its surface area.**

Solution :

Let 'r' be the original radius of sphere

$$\therefore \text{Its surface area} = 4\pi r^2$$

If the radius increases by 25%

$$\text{New radius} = r + \frac{25}{100}r$$

$$= r + \frac{1}{4}r$$

$$= \frac{5}{4}r$$

$$\text{New surface area} = 4\pi \left(\frac{5}{4}r \right)^2$$

$$= 4\pi \times \frac{25}{16}r^2$$

$$= \frac{25\pi r^2}{4}$$

$$\therefore \text{Increment in SA} = \frac{25\pi r^2}{4} - 4\pi r^2$$

$$= \frac{9\pi r^2}{4}$$

$$\therefore \text{Percentage inc. in SA} = \frac{\frac{9\pi r^2}{4}}{4\pi r^2} \times 100$$

$$= \frac{9}{16} \times 100$$

$$= \frac{225}{4}$$

$$= 56.25\%$$

- 9. The internal and external diameters of a hollow hemispherical vessel are 20 cm and 28 cm respectively. Find the cost to paint the vessel all over at ₹ 0.14 per cm.**

Solution :

Given in a hollow hemisphere

$$D = 28 \text{ cm}, \quad d = 20 \text{ cm}$$

$$\Rightarrow R = 14 \text{ cm}, \quad r = 10 \text{ cm}$$

∴ TSA of hollow hemisphere

$$= \pi(3R^2 + r^2)$$

$$= \frac{22}{7} (588 + 100)$$

$$= \frac{22}{7} \times 688 \text{ cm}^2$$

Given cost of painting = 0.14 / cm²

∴ Total cost of painting

$$= \frac{22}{7} \times 688 \times 0.14$$

$$= ₹ 302.72$$

10. The frustum shaped outer portion of the table lamp has to be painted including the top part. Find the total cost of painting the lamp if the cost of painting 1 sq.cm is ₹ 2.



Solution :

Given in a frustum shaped lamp

$$R = 12\text{m}, r = 6\text{m}, h = 8\text{m}$$

$$l = \sqrt{(R - r)^2 + h^2}$$

$$= \sqrt{36 + 64}$$

$$= \sqrt{100} = 10\text{ cm}$$

∴ Required portion to be painted =

$$\text{CSA of frustum} + \pi r^2$$

$$= \pi (R + r) l + \pi r^2$$

$$= \pi [18(10) + 36]$$

$$= \frac{22}{7} \times 216$$

$$= \frac{4752}{7}$$

$$= 678.86\text{ m}^2$$

Given cost of painting = Rs.2/m²

∴ Total cost = 678.86 × 2

$$= \text{Rs.}1357.72/-$$

II. VOLUME

Key Points

Right Circular and Hollow Cylinder

- ✓ Volume of a cylinder = $\pi r^2 h$ cu. units.
- ✓ Volume of a hollow cylinder = $\pi(R^2 - r^2)h$ cu. units.
- ✓ Volume of a cone = $\frac{1}{3} \pi r^2 h$ cu. units.

Sphere and Hemi-sphere

- ✓ Volume of a sphere = $\frac{4}{3} \pi r^3$ cu. units.
- ✓ Volume of a hollow sphere = $\frac{4}{3} \pi(R^3 - r^3)$ cu. units.
- ✓ Volume of a solid hemisphere = $\frac{2}{3} \pi r^3$ cu. units.
- ✓ Volume of a hollow hemisphere = $\frac{2}{3} \pi(R^3 - r^3)$

Frustum of a Cone

- ✓ Volume of a frustum = $\frac{\pi h}{3} (R^2 + Rr + r^2)$ cu. units.

Example 7.15

Find the volume of a cylinder whose height is 2 m and whose base area is 250 m^2 .

Solution :

Let r and h be the radius and height of the cylinder respectively.

Given that, height $h = 2 \text{ m}$,

$$\text{base area} = 250 \text{ m}^2$$

Now, volume of a cylinder $= \pi r^2 h$ cu. units

$$= \text{base area} \times h$$

$$= 250 \times 2 = 500 \text{ m}^3$$

Therefore, volume of the cylinder $= 500 \text{ m}^3$

Example 7.16

The volume of a cylindrical water tank is 1.078×10^6 its height.

Solution :

Let r and h be the radius and height of the cylinder respectively.

Given that, volume of the tank

$$= 1.078 \times 10^6 = 1078000 \text{ litre}$$

$$= 1078 \text{ m}^3 \quad \left(\text{since } 1 \text{ l} = \frac{1}{1000} \text{ m}^3 \right)$$

diameter $= 7 \text{ m}$ gives radius $= \frac{7}{2} \text{ m}$

volume of the tank $= \pi r^2 h$ cu. units

$$1078 = \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \times h$$

Therefore, height of the tank is 28 m.

Example 7.17

Find the volume of the iron used to make a hollow cylinder of height 9 cm and whose internal and external radii are 21 cm and 28 cm respectively.

Solution :

Let r , R and h be the internal radius, external radius and height of the hollow cylinder respectively.

Given that, $r = 21 \text{ cm}$, $R = 28 \text{ cm}$, $h = 9 \text{ cm}$

Now, volume of hollow cylinder

$$= \pi(R^2 - r^2) h \text{ cu. units}$$

$$= \frac{22}{7} (28^2 - 21^2) \times 9$$


$$= \frac{22}{7} (784 - 441) \times 9 = 9702$$

Therefore, volume of iron used $= 9702 \text{ cm}^3$.

Example 7.18

For the cylinders A and B

- find out the cylinder whose volume is greater.
- verify whether the cylinder with greater volume has greater total surface area.
- find the ratios of the volumes of the cylinders A and B.

Solution :

(i) Volume of cylinder = $\pi r^2 h$ cu. units

Volume of cylinder A

$$= \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \times 21$$

$$= 808.5 \text{ cm}^3$$

Volume of cylinder B

$$= \frac{22}{7} \times \frac{21}{2} \times \frac{21}{2} \times 7$$

$$= 2425.5 \text{ cm}^3$$

Therefore, volume of cylinder B is greater than volume of cylinder A.

(ii) T.S.A. of cylinder = $2\pi r (h + r)$ sq. units

T.S.A. of cylinder

$$\frac{22}{7} \times \frac{7}{2} \times 2$$

T.S.A. of cylinder

$$B = 2 \times \frac{22}{7} \times \frac{21}{2} \times (7 + 10.5) = 1155 \text{ cm}^2$$

Hence verified that cylinder B with greater volume has a greater surface area.

$$(iii) \frac{\text{Volume of cylinder A}}{\text{Volume of cylinder B}} = \frac{808.5}{2425.5} = \frac{1}{3}$$

Therefore, ratio of the volumes of cylinders A and B is 1:3.

Example 7.19

The volume of a solid right circular cone is 11088 cm^3 . If its height is 24 cm then find the radius of the cone.

Solution :

Let r and h be the radius and height of the cone respectively.

Given that,

Volume of the cone = 11088 cm^3

$$\frac{1}{3} \pi r^2 h = 11088$$

$$\frac{1}{3} \times \frac{22}{7} \times r^2 \times 24 = 11088$$

$$r^2 = 441$$

Therefore, radius of the cone $r = 21 \text{ cm}$

Example 7.20

The ratio of the volumes of two cones is 2 : 3. Find the ratio of their radii if the height of second cone is double the height of the first..

Solution :

Let r_1 and h_1 be the radius and height of the cone-I and let r_2 and h_2 be the radius and height of the

Given $h_2 = 2h_1$ and $\frac{\text{Volume of the cone I}}{\text{Volume of the cone II}} = \frac{2}{3}$

$$\frac{\frac{1}{3} \pi r_1^2 h_1}{\frac{1}{3} \pi r_2^2 h_2} = \frac{2}{3}$$

$$\frac{r_1^2}{r_2^2} \times \frac{h_1}{2h_1} = \frac{2}{3}$$

$$\frac{r_1^2}{r_2^2} = \frac{4}{3} \text{ gives } \frac{r_1}{r_2} = \frac{2}{\sqrt{3}}$$

Therefore, ratio of their radii = $2 : \sqrt{3}$

Example 7.21

The volume of a solid hemisphere is 29106 cm^3 . Another hemisphere whose volume is two-third of the above is carved out. Find the radius of the new hemisphere.

Solution :

Let r be the radius of the hemisphere.

Given that, volume of the hemisphere

$$= 29106 \text{ cm}^3$$

Now, volume of new hemisphere

$$= \frac{2}{3} (\text{Volume of original sphere})$$

$$= \frac{2}{3} \times 29106$$

Volume of new hemisphere = 19404 cm^3

$$\frac{2}{3} \pi r^3 = 19404$$

$$r^3 = \frac{19404 \times 3 \times 7}{2 \times 22} = 9261$$

$$r = \sqrt[3]{9261} = 21 \text{ cm}$$

Therefore, $r = 21 \text{ cm}$.

Example 7.22

Calculate the weight of a hollow brass sphere if the inner diameter is 14 cm and thickness is 3

Solution :

Let r and R be the inner and outer radii of the hollow sphere.

Given that, inner diameter $d = 14 \text{ cm}$; inner radius $r = 7 \text{ cm}$;

$$\text{thickness} = 1 \text{ mm} = \frac{1}{10} \text{ cm}$$

$$\text{Outer radius } R = 7 + \frac{1}{10} = \frac{71}{10} = 7.1 \text{ cm}$$

$$\begin{aligned} \text{Volume of hollow sphere} &= \frac{4}{3} \pi (R^3 - r^3) \text{ cu.cm} \\ &= \frac{4}{3} \times \frac{22}{7} (357.91 - 343) \\ &= 62.48 \text{ cm}^3 \end{aligned}$$

But, weight of brass in $1 \text{ cm}^3 = 17.3 \text{ gm}$

$$\text{Total weight} = 17.3 \times 62.48 = 1080.90 \text{ gm}$$


Therefore, total weight is 1080.90 grams.

Example 7.23

If the radii of the circular ends of a frustum which is 45 cm high are 28 cm and 7 cm, find the volume of the frustum.

Solution :

Let h , r and R be the height, top and bottom radii of the frustum.



Given that, $h = 45 \text{ cm}$, $R = 28 \text{ cm}$, $r = 7 \text{ cm}$

Now,

$$\begin{aligned} \text{Volume} &= \frac{1}{3} \pi [R^2 + Rr + r^2] h \text{ cu.units} \\ &= \frac{1}{3} \times \frac{22}{7} \times [28^2 + (28 \times 7) + 7^2] \times 45 \\ &= \frac{1}{3} \times \frac{22}{7} \times 1029 \times 45 = 48510 \end{aligned}$$

Therefore, volume of the frustum is 48510 cm^3

EXERCISE 7.2

1. A 14 m deep well with inner diameter 10 m is dug and the earth taken out is evenly spread all around the well to form an embankment of width 5 m. Find the height of the embankment.

Solution :

Given radius of well, $r = 5 \text{ m}$

height of well, $h = 14 \text{ m}$

\therefore Volume of earth taken out = $\pi r^2 h$

$$\begin{aligned} &= \frac{22}{7} \times 25 \times 14 \\ &= 1100 \text{ m}^3 \end{aligned}$$

Since it is spread to form embankment which is the form of hollow cylinder,

Inner radius = 5 cm,

Outer radius = 5 + 5 (Given width = 5m)
= 10 m

$$\begin{aligned}\therefore \pi h_1 (R^2 - r^2) &= 1100 \\ \Rightarrow \frac{22}{7} \times h_1 (10^2 - 5^2) &= 1100 \\ \Rightarrow \frac{22}{7} \times h_1 (75) &= 1100 \\ \Rightarrow h_1 &= \frac{1100 \times 7}{22 \times 75} \\ h_1 &= 4.67 \text{ m}\end{aligned}$$

\therefore height of embankment = 4.67 m

2. A cylindrical glass with diameter 20 cm has water to a height of 9 cm. A small cylindrical metal of radius 5 cm and height 4 cm is immersed it completely. Calculate the raise of the water in the glass?

Solution :

Cylindrical Glass	Cylindrical Metal
-------------------	-------------------

R = 10 cm	r = 5 cm
-----------	----------

H = 9 cm	h = 4 cm
----------	----------

When cylindrical metal is immersed completely in glass,

Total volume = Vol. of glass
+ Vol. of metal

$$= \pi[100 \times 9 + 25 \times 4] = 1000\pi \text{ cm}^3$$

Let h_1 be the height of water level

$$\therefore \pi R^2 h_1 = 1000\pi$$

$$\Rightarrow 100 h_1 = 1000$$

$$\therefore h_1 = 10 \text{ cm}$$

$$\therefore \text{Rise in water level} = h_1 - H$$

$$= 10 - 9 = 1 \text{ cm}$$

3. If the circumference of a conical wooden piece is 484 cm then find its volume when its height is 105 cm.

Solution :

Given circumference of a cone = 484 cm

$$\text{ie } 2\pi r = 484$$

$$\Rightarrow 2 \times \frac{22}{7} \times r = 484$$

$$\Rightarrow r = \frac{484 \times 7}{2 \times 22}$$

$$r = 77 \text{ cm}$$

Also given h = 105 cm

$$\therefore \text{Volume of cone} = \frac{1}{3} \pi r^2 h$$

$$= \frac{1}{3} \times \frac{22}{7} \times 77 \times 77 \times 105$$

$$= 652190 \text{ cm}^3$$

4. A conical container is fully filled with petrol. The radius is 10m and the height is 15 m. If the container can release the petrol through its bottom at the rate of 25 cu. meter per minute, in how many minutes the container will be emptied. Round off your answer to the nearest minute.

Solution :

Given r = 10m, h = 15m in a cone

$$\therefore \text{Volume of the cone} = \frac{1}{3} \pi r^2 h$$

$$= \frac{1}{3} \times \frac{22}{7} \times 100 \times 15$$

$$= \frac{11000}{7} \text{ cm}^3$$


Rate of releasing the petrol = 25 cm³/min

∴ Total time taken to empty the can

$$\begin{aligned}
 &= \frac{11000}{7 \times 25} \\
 &= \frac{440}{7} \\
 &= 62.851 \text{ min.} \\
 &\approx 63 \text{ min.}
 \end{aligned}$$

5. A right angled triangle whose sides are 6 cm, 8 cm and 10 cm is revolved about the sides containing the right angle in two ways. Find the difference in volumes of the two solids so formed.

Solution :



- i) When it revolves about AB = 6 cm

$$h = 6 \text{ cm, } r = 8 \text{ cm}$$

∴ Volume of the cone

$$\begin{aligned}
 &= \frac{1}{3} \pi r^2 h \\
 &= \frac{1}{3} \times \frac{22}{7} \times 64 \times 6 \\
 &= \frac{22 \times 64 \times 2}{7} \\
 &= \frac{2816}{7} \text{ cm}^3
 \end{aligned}$$

- ii) When it revolves about BC = 8 cm

$$h = 8 \text{ cm, } r = 6 \text{ cm}$$

∴ Volume of the cone

$$\begin{aligned}
 &= \frac{1}{3} \pi r^2 h \\
 &= \frac{1}{3} \times \frac{22}{7} \times 36 \times 8 \\
 &= \frac{22 \times 12 \times 8}{7} \\
 &= \frac{2112}{7} \text{ cm}^3
 \end{aligned}$$

∴ Difference in volumes

$$\begin{aligned}
 &= \frac{2816}{7} - \frac{2112}{7} \\
 &= \frac{704}{7} \\
 &= 100.58 \text{ cm}^3
 \end{aligned}$$

6. The volumes of two cones of same base

the ratio of heights.

Solution :

Given volumes of 2 cones

$$= 3600 \text{ cm}^3 \text{ \& } 5040 \text{ cm}^3$$

& base radius are equal

$$\therefore \text{Ratio of volumes} = \frac{V_1}{V_2} = \frac{3600}{5040}$$

$$\Rightarrow \frac{\frac{1}{3} \pi r_1^2 h_1}{\frac{1}{3} \pi r_2^2 h_2} = \frac{3600}{5040}$$

$$\Rightarrow \frac{h_1}{h_2} = \frac{40}{56} = \frac{5}{7}$$

$$\therefore h_1 : h_2 = 5 : 7$$

7. If the ratio of radii of two spheres is 4 : 7, find the ratio of their volumes.

Solution :

Given ratio of radii of 2 spheres = 4 : 7

$$\text{ie } \frac{r_1}{r_2} = \frac{4}{7}$$

$$\begin{aligned}\therefore \text{Ratio of their volumes} &= \frac{V_1}{V_2} \\ &= \frac{\frac{4}{3}\pi r_1^3}{\frac{4}{3}\pi r_2^3} \\ &= \left(\frac{r_1}{r_2}\right)^3 \\ &= \left(\frac{4}{7}\right)^3 \\ &= \frac{64}{343}\end{aligned}$$

\therefore Ratio of the volumes = 64 : 343

8. A solid sphere and a solid hemisphere have equal total surface area. Prove that the ratio of their volume is $3\sqrt{3} : 4$.

Solution :

Given TSA of a solid sphere

= TSA of a solid hemisphere

$$\Rightarrow 4\pi R^2 = 3\pi r^2$$

$$\Rightarrow \therefore \frac{R^2}{r^2} = \frac{3}{4}$$

$$\therefore \frac{R}{r} = \frac{\sqrt{3}}{2}$$

$$\begin{aligned}\therefore \text{Ratio of their volumes} &= \frac{\frac{4}{3}\pi R^3}{\frac{2}{3}\pi r^3} \\ &= \frac{2R^3}{r^3}\end{aligned}$$

$$\begin{aligned}&= 2\left[\frac{R}{r}\right]^3 \\ &= 2\left(\frac{\sqrt{3}}{2}\right)^3 \\ &= 2 \times \frac{3\sqrt{3}}{8} \\ &= \frac{3\sqrt{3}}{4}\end{aligned}$$

\therefore Ratio of the volumes = $3\sqrt{3} : 4$

9. The outer and the inner surface areas of a spherical copper shell are $576\pi \text{ cm}^2$ and $324\pi \text{ cm}^2$ respectively. Find the volume of the material required to make the shell.

Solution :

$$\begin{array}{l|l}\text{Given } 4\pi R^2 = 576\pi & 4\pi r^2 = 324\pi \\ R^2 = 144 & r^2 = 81 \\ R = 12 \text{ cm} & r = 9 \text{ cm}\end{array}$$

\therefore Volume of the material

$$\begin{aligned}&= \frac{4}{3}\pi (R^3 - r^3) \\ &= \frac{4}{3} \times \frac{22}{7} (1728 - 729) \\ &= \frac{4}{3} \times \frac{22}{7} \times 999 \\ &= \frac{88 \times 333}{7} = 4186.29 \text{ cm}^3\end{aligned}$$

10. A container open at the top is in the form of a frustum of a cone of height 16 cm with radii of its lower and upper ends are 8 cm and 20 cm respectively. Find the cost of milk which can completely fill a container at the rate of ₹ 40 per litre.

Solution :

Given $R = 20\text{cm}$, $r = 8\text{cm}$, $h = 16\text{cm}$

in frustum of a cone

\therefore Volume of frustum of a cone

$$\begin{aligned}
 &= \frac{\pi h}{3} (R^2 + Rr + r^2) \\
 &= \frac{22}{7} \times \frac{16}{3} (400 + 160 + 64) \\
 &= \frac{22}{7} \times \frac{16}{3} \times \frac{624}{1} \\
 &= \frac{73216}{7} \\
 &= 10,459.42 \text{ cm}^3 \\
 &= \frac{10459.42}{1000} \text{ litres}
 \end{aligned}$$

\therefore Cost of milk at ₹ 40 / lr,

$$= 10.459 \times 40$$

$$= ₹ 418.36$$


III. Volume and Surface Area of Combined Solids

Example 7.24

A toy is in the shape of a cylinder surmounted by a hemisphere. The height of the toy is 25 cm. Find the total surface area of the toy if its common diameter is 12 cm.

Solution :

Let r and h be the radius and height of the cylinder respectively.



Given that, diameter $d = 12 \text{ cm}$, radius $r = 6 \text{ cm}$

Total height of the toy is 25 cm

Therefore, height of the cylindrical portion

$$= 25 - 6 = 19 \text{ cm}$$

T.S.A. of the toy = C.S.A. of the cylinder +

C.S.A. of the hemisphere

+ Base Area of the cylinder

$$\begin{aligned}
 &= 2\pi rh + 2\pi r^2 + \pi r^2 \\
 &= \pi r(2h + 3r) \text{ sq. units} \\
 &= \frac{22}{7} \times 6 \times (38 + 18) \\
 &= \frac{22}{7} \times 6 \times 56 = 1056
 \end{aligned}$$

Therefore, T.S.A. of the toy is 1056 cm^2 .

Example 7.25

A jewel box (Fig. 7.39) is in the shape of a cuboid of dimensions $30 \text{ cm} \times 15 \text{ cm} \times 10 \text{ cm}$ surmounted by a half part of a cylinder as shown in the figure. Find the volume and T.S.A. of the box.




Fig. 7.39

Solution :

Let l , b and h_1 be the length, breadth and height of the cuboid. Also let us take r and h_2 be the radius and height of the cylinder.

Now, Volume of the box =

Volume of the cuboid +

$$\frac{1}{2} (\text{Volume of cylinder})$$

$$\begin{aligned}
 &= (l \times b \times h_1) + \frac{1}{2} (\pi r^2 h_2) \text{ cu. units} \\
 &= (30 \times 15 \times 10) + \frac{1}{2} \left(\frac{22}{7} \times \frac{15}{2} \times \frac{15}{2} \times 30 \right) \\
 &= 4500 + 2651.79 = 7151.79
 \end{aligned}$$

Therefore, Volume of the box = 7151.79 cm³

Now, T.S.A. of the box = C.S.A. of the cuboid +

$$\begin{aligned}
 &\frac{1}{2} (\text{C.S.A. of the cylinder}) \\
 &= 2(l + b)h_1 + \frac{1}{2} (2\pi r h_2) \\
 &= 2(45 \times 10) + \left(\frac{22}{7} \times \frac{15}{2} \times 30 \right) \\
 &= 900 + 707.14 = 1607.14
 \end{aligned}$$

2

Example 7.26

Arul has to make arrangements for the accommodation of 150 persons for his family function. For this purpose, he plans to build a tent which is in the shape of cylinder surmounted by a cone. Each person occupies 4 sq. m of the space on ground and 40 cu. meter of air to breathe. What should be the height of the conical part of the tent if the height of cylindrical part is 8 m?

Solution :

Let h_1 and h_2 be the height of cylinder and cone respectively.



Fig. 7.40

$$\begin{aligned}
 \text{Area for one person} &= 4 \text{ sq. m} \\
 \text{Total number of persons} &= 150
 \end{aligned}$$

$$\text{Therefore total base area} = 150 \times 4$$

$$\pi r^2 = 600$$

$$r^2 = 600 \times \frac{7}{22} = \frac{2100}{11} \quad \dots\dots (1)$$

$$\text{Volume of air required for 1 person} = 40 \text{ m}^3$$

$$\text{Total Volume of air required for 150 persons}$$

$$= 150 \times 40 = 6000 \text{ m}^3$$

$$\pi r^2 h_1 + \frac{1}{3} \pi r^2 h_2 = 6000$$

$$\pi r^2 \left(h_1 + \frac{1}{3} h_2 \right) = 6000$$

$$\frac{22}{7} \times \frac{2100}{11} \left(8 + \frac{1}{3} h_2 \right) = 6000 \quad [\text{using (1)}]$$

$$8 + \frac{1}{3} h_2 = \frac{6000 \times 7 \times 11}{22 \times 2100}$$

$$\frac{1}{3} h_2 = 10 - 8 = 2$$

Therefore, the height of the conical tent h_2 is 6 m

Example 7.27

A funnel consists of a frustum of a cone attached to a cylindrical portion 12 cm long attached at the bottom. If the total height be 20 cm, diameter of the cylindrical portion be 12 cm and the diameter of the top of the funnel be 24 cm. Find the outer surface area of the funnel.

Solution :

Let R , r be the top and bottom radii of the frustum.

Let h_1 , h_2 be the heights of the frustum and cylinder respectively.




Fig. 7.41

Given that, $R = 12$ cm, $r = 6$ cm, $h_2 = 12$ cm

Now, $h_1 = 20 - 12 = 8$ cm

Here, Slant height of the frustum

$$\begin{aligned} l &= \sqrt{(R-r)^2 + h_1^2} \text{ units} \\ &= \sqrt{36 + 64} \\ l &= 10 \text{ cm} \end{aligned}$$

Outer surface area

$$\begin{aligned} &= 2\pi rh_2 + \pi(R+r)l \text{ sq. units} \\ &= \pi[2rh_2 + (R+r)l] \\ &= \pi[(2 \times 6 \times 12) + (18 \times 10)] \\ &= \pi[144 + 180] \\ &= \frac{22}{7} \times 324 = 1018.28 \end{aligned}$$

Therefore, outer surface area of the funnel is 1018.28 cm^2

Example 7.28

A hemispherical section is cut out from one face of a cubical block (Fig. 7.42) such that the diameter l of the hemisphere is equal to side length of the cube. Determine the surface area of the remaining solid.




Fig. 7.42

Solution :

Let r be the radius of the hemisphere.

Given that, diameter of the hemisphere = side of the cube = l

Radius of the hemisphere = $\frac{l}{2}$

TSA of the remaining solid =

Surface area of the cubical part +
C.S.A. of the hemispherical part –
Area of the base of the hemispherical part
 $= 6 \times (\text{Edge})^2 + 2\pi r^2 - \pi r^2$

$$= 6 \times (l)^2 + \pi \left(\frac{l}{2} \right)^2 = \frac{1}{4} (24 + \pi) l^2$$


Total surface area of the remaining solid =

$$\frac{1}{4} (24 + \pi) l^2 \text{ sq. units}$$

EXERCISE 7.3

1. A vessel is in the form of a hemispherical bowl mounted by a hollow cylinder. The diameter is 14 cm and the height of the vessel is 13 cm. Find the capacity of the vessel.

Solution:



Given $AD = BE = 7 \text{ cm} = \text{radius of the vessel} = BC$

$AC = 13 \text{ cm} = \text{height of the vessel}$

$$\therefore AB = 13 - 7$$

$$= 6 \text{ cm} = \text{height of cylindrical part}$$

\therefore Capacity of the vessel

= Capacity of cylinder + Capacity of HS

$$= \pi r^2 h + \frac{2}{3} \pi r^3$$

$$= \pi r^2 \left[h + \frac{2}{3} r \right]$$


$$= \frac{22}{7} \times 49 \left[6 + \frac{14}{3} \right]$$

$$= 154 \times \frac{32}{3}$$

3

2. Nathan, an engineering student was asked to make a model shaped like a cylinder with two cones attached at its two ends. The diameter of the model is 3 cm and its length is 12 cm. If each cone has a height of 2 cm, find the volume of the model that Nathan made.

Solution:



Cone

$$h = 2 \text{ cm}$$

$$r = 1.5 \text{ cm} = \frac{3}{2}$$

Cylinder

$$H = 8 \text{ cm}$$

$$r = 1.5 \text{ cm} = \frac{3}{2}$$

\therefore Volume of the model =

2 (Vol. of Cone) + Vol. of Cylinder

$$= \frac{2}{3} \pi r^2 h + \pi r^2 H$$

$$= \pi r^2 \left[\frac{2h}{3} + H \right]$$

$$= \frac{22}{7} \times \frac{9}{4} \left[\frac{4}{3} + 8 \right]$$

$$= \frac{11 \times 9}{7 \times 2} \left[\frac{28}{3} \right]$$

$$= \frac{11 \times 3 \times 14}{7}$$

$$= 66 \text{ cm}^3$$

3. From a solid cylinder whose height is 2.4 cm and the diameter 1.4 cm, a cone of the same height and same diameter is carved out. Find the volume of the remaining solid to the nearest cm^3 .

Solution:

Volume of the remaining solid

= Vol. of Cylinder – Vol. of Cone


$$= \pi r^2 h - \frac{1}{3} \pi r^2 h$$

$$= \frac{2}{3} \pi r^2 h$$

$$= \frac{2}{3} \times \frac{22}{7} \times 0.7 \times 0.7 \times 2.4$$

$$= 2.46 \text{ cm}^3$$

4. A solid consisting of a right circular cone of height 12 cm and radius 6 cm standing on a hemisphere of radius 6 cm is placed upright in a right circular cylinder full of water such that it touches the bottom. Find the volume of the water displaced out of the cylinder, if the radius of the cylinder is 6 cm and height is 18 cm.

**Solution:**

Cone	Hemisphere	Cylinder
------	------------	----------

$r = 6\text{ cm}$	$r = 6\text{ cm}$	$r = 6\text{ cm}$
-------------------	-------------------	-------------------

$h = 12\text{ cm}$		$H = 18\text{ cm}$
--------------------	--	--------------------

Volume of water displaced out of cylinder =

Volumme of cone + Volume of HS

$$\begin{aligned}
 &= \frac{1}{3}\pi r^2 h + \frac{2}{3}\pi r^3 \\
 &= \frac{1}{3}\pi r^2 [h + 2r] \\
 &= \frac{1}{3} \times \frac{22}{7} \times 36(12 + 12) \\
 &= \frac{22}{7} \times 12 \times 24 \\
 &= 905.14\text{ cm}^3
 \end{aligned}$$

Note : When the conical hemisphere is completely submerged in water inside the cylinder,


Volume of water left in the cylinder.

= Volume of cylinder – [Volume of cone + Vol. of Hemi sphere)

$$= \pi r^2 H - \left[\frac{1}{3}\pi r^2 h + \frac{2}{3}\pi r^3 \right]$$

$$\begin{aligned}
 &= \pi \left[36 \times 18 - \frac{1}{3} \times 36 \times 12 - \frac{2}{3} \times 216 \right] \\
 &= \frac{22}{7} [648 - 144 - 144] \\
 &= \frac{22}{7} \times 360 \\
 &= \frac{7920}{7} \\
 &= 1131.42\text{ cm}^3
 \end{aligned}$$

5. A capsule is in the shape of a cylinder with two hemisphere stuck to each of its ends. If the length of the entire capsule is 12 mm and the diameter of the capsule is 3 mm, how much medicine it can hold?


Solution:**Cylinder****Hemisphere** $H = 9\text{ mm}$ $r = 1.5\text{ mm} = \frac{3}{2}$ $r = 1.5\text{ mm} = \frac{3}{2}$ \therefore Volume of the Capsule =

Vol. of Cylinder + 2 (Vol. of hemisphere)

$$\begin{aligned}
 &= \pi r^2 H + 2 \left(\frac{2}{3}\pi r^3 \right) \\
 &= \frac{22}{7} \left[\frac{9}{4} \times 9 + \frac{4}{3} \times \frac{27}{8} \right] \\
 &= \frac{22}{7} \left[\frac{81}{4} + \frac{9}{2} \right] \\
 &= \frac{22}{7} \left[\frac{81 + 18}{4} \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{22 \times 99}{28} \\
 &= \frac{11 \times 99}{14} \\
 &= 77.78 \text{ mm}^3
 \end{aligned}$$

6. As shown in figure a cubical block of side 7 cm is surmounted by a hemisphere. Find the surface area of the solid.



Solution:

Given side of cube = 7 cm

$$\frac{7}{2}$$


Surface area of the solid = CSA of cube + CSA of hemisphere – Base area of HS

$$\begin{aligned}
 &= 6a^2 + 2\pi r^2 - \pi r^2 \\
 &= 6(49) + \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \\
 &= 294 + \frac{77}{2} \\
 &= 294 + 38.5 \\
 &= 332.5 \text{ cm}^2
 \end{aligned}$$

7. A right circular cylinder just enclose a sphere of radius r units. Calculate
- the surface area of the sphere
 - the curved surface area of the cylinder
 - the ratio of the areas obtained in (i) and (ii).

Solution:


From the given fig, it is clear that $h = 2r$



- Surface area of sphere
 $= 4\pi r^2$ sq. units
- CSA of the cylinder = $2\pi rh$
 $= 2\pi r (2r)$
 $= 4\pi r^2$ sq. units
- Ratio of areas obtained in (i) & (ii)
 $= 4\pi r^2 : 4\pi r^2$
 $= 1 : 1$

8. A shuttle cock used for playing badminton has the shape of a frustum of a cone is mounted on a hemisphere. The diameters of the frustum are 5 cm and 2 cm. The height of the entire shuttle cock is 7 cm. Find its external surface area.

Solution:



$AB = BD =$ radius of hemisphere = 1 cm

radius of frustum = r

$AC = 7\text{ cm} =$ Total length of cock

$\therefore BC = 7 - 1 = 6\text{ cm} =$ height of frustum

$CE = 2.5\text{ cm} = R$

$$\therefore l = \sqrt{h^2 + (R - r)^2} = \sqrt{26 + (1.5)^2} = 6.18$$

\therefore External Surface Area =

CSA of Frustum + CSA of HS

$$= \pi (R + r)l + 2\pi r^2$$

$$= \pi [(2.5 + 1) 6.18 + 2 \times 1]$$

$$= \frac{22}{7} \left[\frac{7}{2} (6.1) + 2 \right]$$

$$= \frac{22}{7} [21.35 + 2]$$

$$= \frac{22 \times 23.35}{7}$$

$$= \frac{513.7}{7}$$

$$= 73.39 \text{ cm}^2$$

IV. Conversion of Solids

Example 7.29

A metallic sphere of radius 16 cm is melted and recast into small spheres each of radius 2 cm. How many small spheres can be obtained?

Solution :

Let the number of small spheres obtained be n .

Let r be the radius of each small sphere and R be the radius of metallic sphere.

Here, $R = 16 \text{ cm}$, $r = 2 \text{ cm}$

Now, n (Volume of a small sphere)

= Volume of big metallic sphere

$$n \left(\frac{4}{3} \pi r^3 \right) = \frac{4}{3} \pi R^3$$

$$n \left(\frac{4}{3} \pi \times 2^3 \right) = \frac{4}{3} \pi \times 16^3$$

$$8n = 4096 \text{ gives } n = 512$$

Therefore, there will be 512 small spheres.

Example 7.30

A cone of height 24 cm is made up of modeling clay. A child reshapes it in the form of a cylinder of same radius as cone. Find the height of the cylinder.

Solution :

Let h_1 and h_2 be the heights of a cone and cylinder respectively.

Also, let r be the radius of the cone.

Given that, height of the cone $h_1 = 24 \text{ cm}$; radius of the cone and cylinder $r = 6 \text{ cm}$

Since, Volume of cylinder = Volume of cone

$$\pi^2 h_2 = \frac{1}{3} \pi r^2 h_1$$

$$h_2 = \frac{1}{3} \times h_1 \text{ gives } h_2 = \frac{1}{3} \times 24 = 8$$

Example 7.31

A right circular cylindrical container of base radius 6 cm and height 15 cm is full of ice cream. The ice cream is to be filled in cones of height 9 cm and base radius 3 cm, having a hemispherical cap. Find the number of cones needed to empty the container.

Solution :

Let h and r be the height and radius of the cylinder respectively.

Given that, $h = 15 \text{ cm}$, $r = 6 \text{ cm}$

Volume of the container $V = \pi r^2 h$ cubic units.

$$= \frac{22}{7} \times 6 \times 6 \times 15$$

Let, $r_1 = 3 \text{ cm}$, $h_1 = 9 \text{ cm}$ be the radius and height of the cone.

Also, $r_1 = 3 \text{ cm}$ is the radius of the hemispherical cap.

Volume of one ice cream cone = (Volume of the cone + Volume of the hemispherical cap)

$$\begin{aligned}
 &= \frac{1}{3} \pi r_1^2 h_1 + \frac{2}{3} \pi r_1^3 \\
 &= \frac{1}{3} \times \frac{22}{7} \times 3 \times 3 \times 9 + \frac{2}{3} \times \frac{22}{7} \times 3 \times 3 \times 3 \\
 &= \frac{22}{7} \times 9(3 + 2) = \frac{22}{7} \times 45
 \end{aligned}$$

Number of cones =

$$\frac{\text{Volume of the cylinder}}{\text{Volume of one ice cream cone}}$$

Number of ice cream cones needed =

$$\begin{aligned}
 &\frac{\frac{22}{7} \times 6 \times 6 \times 15}{\frac{22}{7} \times 45} = 12
 \end{aligned}$$

Thus 12 ice cream cones are required to empty the cylindrical container.

EXERCISE 7.4

1. An aluminium sphere of radius 12 cm is melted to make a cylinder of radius 8 cm. Find the height of the cylinder.

Solution:

Given radius of sphere = 12 cm = R &

radius of cylinder = 8 cm = r

By the data given,

Volume of sphere = Volume of Cylinder

$$\Rightarrow \frac{4}{3} \pi R^3 = \pi r^2 h$$

$$\Rightarrow \frac{4}{3} \times 12 \times 12 \times 12 = 8 \times 8 \times h$$

$$\Rightarrow h = 36 \text{ cm}$$

\therefore Height of the cylinder = 36 cm

2. Water is flowing at the rate of 15 km per hour through a pipe of diameter 14 cm into a rectangular tank which is 50 m long and 44 m wide. Find the time in which the level of water in the tanks will rise by 21 cm.

Solution:

Cylindrical Pipe

Given, Speed of water in the pipe

$$= 15 \text{ Km/hr}$$

$$H = 15000 \text{ m}$$

$$\text{Radius of pipe } r = 7 \text{ cm} = \frac{7}{100} \text{ m}$$

Rectangular Tank

$$l = 50 \text{ m} \quad b = 44 \text{ m}$$

$$\frac{21}{100}$$

$$\begin{aligned}
 \therefore \text{ Required time} &= \frac{\text{Volume of tank}}{\text{Volume of pipe}} \\
 &= \frac{lbh}{\pi r^2 H} \\
 &= \frac{50 \times 44 \times \frac{21}{100}}{\frac{22}{7} \times \frac{7}{100} \times \frac{7}{100} \times 15000} \\
 &= 2 \text{ hrs}
 \end{aligned}$$

3. A conical flask is full of water. The flask has base radius r units and height h units, the water poured into a cylindrical flask of base radius xr units. Find the height of water in the cylindrical flask.

Solution:

By the data given,

Volume of Cylindrical Flask =

Volume of Conical Flask

$$\Rightarrow \pi(xr)^2 H = \frac{1}{3} \pi r^2 h$$

$$\Rightarrow x^2 r^2 H = \frac{1}{3} r^2 h$$

$$\Rightarrow H = \frac{h}{3x^2}$$

$$\therefore \text{Height of the Cylindrical Flask} = \frac{h}{3x^2} \text{ cm}$$

4. A solid right circular cone of diameter 14 cm and height 8 cm is melted to form a hollow sphere. If the external diameter of the sphere is 10 cm, find the internal diameter.

Solution:

Right Circular Cone

$$r = 7 \text{ cm}$$

$$h = 8 \text{ cm}$$

By the problem,

Volume of Hollow Sphere =

Vol. of Right Circular Cone

$$\Rightarrow \frac{4}{3} \pi (R^3 - r^3) = \frac{1}{3} \pi r^2 h$$

$$\Rightarrow 4(125 - r^3) = 49 \times 8$$

$$\Rightarrow 125 - r^3 = 49 \times 2$$

$$\Rightarrow r^3 = 125 - 98$$

$$r^3 = 27$$

$$\therefore r = 3$$

$$\therefore \text{Internal diameter of hollow sphere} = 6 \text{ cm}$$

5. Seenu's house has an overhead tank in the shape of a cylinder. This is filled by pumping water from a sump (under-ground tank) which is in the shape of a cuboid. The sump has dimensions 2 m × 1.5 m × 1 m. The overhead tank has its radius of 60 cm and height 105 cm. Find the volume of the water left in the sump after the overhead tank has been completely filled with water from the sump which has been full, initially.

Solution:

Over head tank

(Cylinder)

$$R = 60 \text{ cm}$$

$$H = 105 \text{ cm}$$

Sump

(Cuboid)

$$l = 2 \text{ m} = 200 \text{ cm}$$

$$b = 1.5 \text{ m} = 150 \text{ cm}$$

$$h = 1 \text{ m} = 100 \text{ cm}$$

Volume of water left

$$= \text{Volume of Sump} - \text{Volume of tank}$$

$$= l b h - \pi R^2 H$$

$$= 200 \times 150 \times 100 - \frac{22}{7} \times 60 \times 60 \times 105$$

$$= 3000000 - 1188000$$

$$= 2812000 \text{ cm}^3$$

6. The internal and external diameter of a hollow hemispherical shell are 6 cm and

cast into a solid cylinder of diameter 14 cm, then find the height of the cylinder.

Solution:

Hollow Hemisphere

$$R = 5 \text{ cm}$$

$$r = 3 \text{ cm}$$

Solid Cylinder

$$r = 7 \text{ cm}$$

$$h = ?$$

\therefore By the problem given,

$$\text{Volume of Solid Cylinder} =$$

$$\text{Volume of Hollow Hemisphere}$$

$$\Rightarrow \pi r^2 h = \frac{2}{3} \pi (R^3 - r^3)$$

$$\Rightarrow 49 \times h = \frac{2}{3} (125 - 27)$$

$$\Rightarrow h = \frac{2}{3} \times \frac{98}{49}$$

$$\therefore h = \frac{4}{3} = 1.33 \text{ cm}$$

$$\therefore \text{Height of Solid Cylinder} = 1.33 \text{ cm}$$

7. A solid sphere of radius 6 cm is melted into a hollow cylinder of uniform thickness. If the external radius of the base of the cylinder is 5 cm and its height is 32 cm, then find the thickness of the cylinder.

Solution:

Solid Sphere	Hollow Cylinder
---------------------	------------------------

$$r = 6 \text{ cm}$$

$$R = 5 \text{ cm}$$

$$H = 32 \text{ cm}$$

$$t = ?$$

By the problem given,

Volume of Hollow Cylinder =

Volume of Solid Sphere

$$\Rightarrow \pi(R^2 - r^2)H = \frac{4}{3}\pi r^3$$

$$\Rightarrow (25 - r^2)32 = \frac{4}{3} \times \cancel{6}^2 \times 6 \times 6$$

$$\Rightarrow 25 - r^2 = \frac{\cancel{4} \times \cancel{2} \times \cancel{6}^3 \times \cancel{6}^3}{\cancel{32}^{\cancel{4} \times \cancel{2}}}$$

$$\Rightarrow 25 - r^2 = 9$$

$$r^2 = 16$$

$$r^2 = 4$$

$$\therefore \text{Thickness} = R - r$$

$$= 5 - 4$$

$$= 1 \text{ cm}$$

8. A hemispherical bowl is filled to the brim with juice. The juice is poured into a cylindrical vessel whose radius is 50% more than its height. If the diameter is same for both the bowl and the cylinder then find the percentage of juice that can be transferred from the bowl into the cylindrical vessel.

Solution:

Hemisphere

$$\text{Radius} = r$$

Cylinder

$$\text{Radius} = r$$

$$= h + \frac{1}{2}h$$

$$r = \frac{3}{2}h$$

$$\therefore \text{Volume of hemisphere} = \frac{2}{3}\pi r^3$$

$$= \frac{2}{3}\pi \times \left(\frac{3}{2}h\right)^3$$

$$= \frac{2}{3}\pi \times \frac{27}{8}h^3$$

$$= \frac{9}{4}\pi h^3$$

$$\text{Volume of Cylinder} = \pi r^2 h$$

$$= \pi \times \left(\frac{3}{2}h\right)^2 h$$

$$= \pi \times \frac{9}{4}h^2 h$$

$$= \frac{9}{4}\pi h^3$$

$$\therefore \text{Vol. of Hemisphere} = \text{Vol. of Cylinder}$$

$$\therefore \% \text{ of juice that can be transferred to the cylindrical vessel} = 100 \%$$

EXERCISE 7.5

Multiple choice questions

- The curved surface area of a right circular cone of height 15 cm and base diameter 16 cm is
 - $60\pi \text{ cm}^2$
 - $68\pi \text{ cm}^2$
 - $120\pi \text{ cm}^2$
 - $136\pi \text{ cm}^2$

Hint :

Ans : (4)

$$h = 15 \text{ cm}, r = 8 \text{ cm}$$

$$\begin{aligned}\Rightarrow l &= \sqrt{h^2 + r^2} \\ &= \sqrt{225 + 64} \\ &= \sqrt{289} \\ &= 17\end{aligned}$$

$$\begin{aligned}\therefore \text{CSA of Cone} &= \pi r l \\ &= \pi \times 8 \times 17 \\ &= 136 \pi \text{ cm}^2\end{aligned}$$

2. If two solid hemispheres of same base radius r units are joined together along their bases, then curved surface area of this new solid is

- (1) $4\pi r^2$ sq. units (2) $6\pi r^2$ sq. units
(3) $3\pi r^2$ sq. units (4) $8\pi r^2$ sq. units

Hint : Ans : (1)

The CSA of the new solid is nothing but the CSA of a sphere = $4\pi r^2$ sq. units

3. The height of a right circular cone whose radius is 5 cm and slant height is 13 cm will be

- (1) 12 cm (2) 10 cm
(3) 13 cm (4) 5 cm

Hint : Ans : (1)

$$r = 5 \text{ cm}, l = 13 \text{ cm}$$

$$\begin{aligned}\therefore h &= \sqrt{l^2 - r^2} \\ &= \sqrt{169 - 25} \\ &= \sqrt{144} \\ &= 12 \text{ cm}\end{aligned}$$

4. If the radius of the base of a right circular cylinder is halved keeping the same height, then the ratio of the volume of the cylinder thus obtained to the volume of original cylinder is

- (1) 1 : 2 (2) 1 : 4
(3) 1 : 6 (4) 1 : 8

Hint : Ans : (2)

$$\frac{\text{Volume of New Cylinder}}{\text{Volume of Original Cylinder}} = \frac{\pi R^2 h}{\pi r^2 h}$$

$$\begin{aligned}\text{where } R &= \frac{r}{2} \\ &= \frac{R^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= \frac{1}{4}\end{aligned}$$

$$\therefore V_1 : V_2 = 1 : 4$$

radius is $\frac{1}{3}$ of its height is

- (1) $\frac{9\pi h^2}{8}$ sq. units (2) $24\pi h^2$ sq. units
(3) $\frac{8\pi h^2}{9}$ sq. units (4) $\frac{56\pi h^2}{9}$ sq. units

Hint : Ans : (3)

$$\text{TSA of Cylinder} = 2\pi r (h + r)$$

$$\begin{aligned}\text{where } r &= \frac{1}{3} h \\ &= 2\pi \times \frac{h}{3} \left(h + \frac{h}{3} \right) \\ &= 2\pi \frac{h}{3} \times \frac{4h}{3} \\ &= \frac{8\pi h^2}{9} \text{ Sq. units}\end{aligned}$$

6. In a hollow cylinder, the sum of the external and internal radii is 14 cm and the width is 4 cm. If its height is 20 cm, the volume of the material in it is

- (1) $5600 \pi \text{ cm}^3$ (2) $11200 \pi \text{ cm}^3$
 (3) $56\pi \text{ cm}^3$ (4) $3600 \pi \text{ cm}^3$

Hint : Ans : (2)

$$R + r = 14 \text{ cm, } h = 20 \text{ cm, } W = 4 \text{ cm}$$

$$R - r = 4 \text{ cm}$$

Volume of hollow cylinder

$$\begin{aligned} &= \pi h (R^2 - r^2) \\ &= \pi h (R + r) (R - r) \\ &= \pi \times 20 \times 14 \times 4 \\ &= 1120\pi \text{ cm}^3 \end{aligned}$$

7. If the radius of the base of a cone is tripled and the height is doubled then the volume is

- (1) made 6 times (2) made 18 times
 (3) made 12 times (4) unchanged

Hint : Ans : (2)

$$\text{Volume of cone} = \frac{1}{3} \pi r^2 h$$

$$\text{When } r \rightarrow 3r, h \rightarrow 2h$$

Volume of new cone

$$\begin{aligned} &= \frac{1}{3} \pi \times 9r^2 \times 2h \\ &= 18 \left(\frac{1}{3} \pi r^2 h \right) \\ &= 18 \text{ times} \end{aligned}$$

8. The total surface area of a hemi-sphere is how much times the square of its radius.

- (1) π (2) 4π (3) 3π
 (4) 2π

Hint : Ans : (3)

$$\text{TSA of a hemisphere} = 3\pi r^2$$

$$= 3\pi (\text{square of its radius})$$

$$= 3\pi \text{ times } r^2$$

9. A solid sphere of radius x cm is melted and cast into a shape of a solid cone of same radius. The height of the cone is

- (1) $3x$ cm (2) x cm
 (3) $4x$ cm (4) $2x$ cm

Hint : Ans : (3)

Volume of sphere = Volume of Cone

$$\Rightarrow \frac{4}{3} \pi r^3 = \frac{1}{3} \pi r^2 h$$

$$\Rightarrow \frac{4}{3} \pi x^3 = \frac{1}{3} \pi x^2 \times h$$

$$\Rightarrow h = \frac{\frac{4}{3} \pi x^3}{\frac{1}{3} \pi x^2} = 4x$$

10. A frustum of a right circular cone is of height 16cm with radii of its ends as 8cm and 20cm. Then, the volume of the frustum is

- (1) $3328\pi \text{ cm}^3$ (2) $3228\pi \text{ cm}^3$
 (3) $3240\pi \text{ cm}^3$ (4) $3340\pi \text{ cm}^3$

Hint : Ans : (1)

Volume of Frustum of a Cone

$$= \frac{\pi h}{3} (R^2 + Rr + r^2)$$

$$= \frac{\pi}{3} \times 16 [400 + 160 + 64]$$

$$= \frac{16\pi}{3} \times 624$$

$$= 16\pi \times 208$$

$$= 3328\pi \text{ cm}^3$$

11. A shuttle cock used for playing badminton has the shape of the combination of
- (1) a cylinder and a sphere
 - (2) a hemisphere and a cone
 - (3) a sphere and a cone
 - (4) frustum of a cone and a hemisphere

Hint :

Ans : (4)

Frustum of a cone & a hemisphere

12. A spherical ball of radius r_1 units is melted to make 8 new identical balls each of radius r_2 units. Then $r_1 : r_2$ is
- (1) 2:1
 - (2) 1:2
 - (3) 4:1
 - (4) 1:4

Hint :

Ans : (1)

Volume of a sphere = 8 (Volume of new identical

$$\frac{4}{3}\pi r_1^3 = 8\left(\frac{4}{3}\pi r_2^3\right)$$

$$\Rightarrow \frac{r_1^3}{r_2^3} = \frac{8}{1}$$

$$\therefore r_1 : r_2 = 2 : 1$$

13. The volume (in cm^3) of the greatest sphere that can be cut off from a cylindrical log of wood of base radius 1 cm and height 5 cm is

- (1) $\frac{4}{3}\pi$
- (2) $\frac{10}{3}\pi$
- (3) 5π
- (4) $\frac{20}{3}\pi$

Hint :

Ans : (1)

$$\text{Volume of sphere} = \frac{4}{3}\pi r^3 \text{ where } r = 1$$


$$= \frac{4}{3}\pi$$

14. The height and radius of the cone of which the frustum is a part are h_1 units and r_1 units respectively. Height of the frustum is h_2 units and radius of the smaller base is r_2 units. If $h_2 : h_1 = 1 : 2$ then $r_2 : r_1$ is

- (1) 1 : 3
- (2) 1 : 2
- (3) 2 : 1
- (4) 3 : 1

Hint :

Ans : (2)

Given $h_2 : h_1 = 1 : 2$

$$\Rightarrow h_2 = \frac{1}{2}h_1 \quad \therefore \frac{r_2}{r_1} = \frac{1}{2}$$

15. The ratio of the volumes of a cylinder, a cone and a sphere, if each has the same diameter and same height is

- (1) 1:2:3
- (2) 2:1:3
- (3) 1:3:2
- (4) 3:1:2

Hint :

Ans : (4)

Ratio of volumes of Cylinder, Cone, Sphere

$$= \pi r^2 h : \frac{1}{3}\pi r^2 h : \frac{4}{3}\pi r^3 h$$

with same height & same radius.

Since each of them has same diameter and same height, $h = 2r$

$$V_1 = \pi r^2(2r) = 2\pi r^3$$

$$V_2 = \frac{1}{3}\pi r^2(2r) = \frac{2}{3}\pi r^3$$

$$V_2 = \frac{4}{3}\pi r^3$$

$$\therefore V_1 : V_2 : V_3 = 2 : \frac{2}{3} : \frac{4}{3}$$

$$= 6 : 2 : 4$$

$$= 3 : 1 : 2$$

UNIT EXERCISE - 7

1. The barrel of a fountain-pen cylindrical in shape, is 7 cm long and 5 mm in diameter. A full barrel of ink in the pen will be used for writing 330 words on an average. How many words can be written using a bottle of

Solution :

Given height of the pen = 7 cm = 70 mm

$$\text{radius} = \frac{5}{2} \text{ mm}$$

\therefore Volume of the pen = $\pi r^2 h$

$$\begin{aligned} &= \frac{22}{7} \times \frac{25}{4} \times \frac{70}{10} \\ &= 1375 \text{ mm}^3 \\ &= 1.375 \text{ cm}^3 \end{aligned}$$

By data given

$1.375 \text{ cm}^3 \rightarrow 330 \text{ words} -$

$$\frac{1}{5} \text{ of a litre} = \frac{1}{5} (1000 \text{ cm}^3)$$

$$\Rightarrow 200 \text{ cm}^3 \rightarrow x \text{ words}$$

$$\therefore x = \frac{200 \times 330}{1.375}$$

$$= 48000 \text{ words}$$

2. A hemi-spherical tank of radius 1.75 m is full of water. It is connected with a pipe which empties the tank at the rate of 7 litre per second. How much time will it take to empty the tank completely?

Solution :

Radius of hemi-spherical tank = 1.75 m

$$r = \frac{7}{4} \text{ m}$$

\therefore Volume of the tank

$$\begin{aligned} &= \frac{2}{3}\pi r^3 \\ &= \frac{2}{3} \times \frac{22}{7} \times \frac{7}{4} \times \frac{7}{4} \times \frac{7}{4} \\ &= \frac{539}{48} \\ &= 11.229 \text{ m}^3 \\ &= 11.229 \times 1000 \text{ litres} \\ &= 11229 \text{ litres} \end{aligned}$$

Water falls at the rate of 7 lrs/second

\therefore Time taken by the pipe to empty the tank

$$\begin{aligned} &= \frac{11229}{7} \text{ Sec} \\ &= 1604 \text{ sec (approx)} \\ &= \frac{1604}{60} \text{ min} \\ &= 27 \text{ min (approx)} \end{aligned}$$

3. Find the maximum volume of a cone that can be carved out of a solid hemisphere of radius r units.

Solution :

Given radius of solid hemisphere = r

Volume of a cone that can be carved

Out of hemisphere

$$= \frac{2}{3}\pi r^3 - \frac{1}{3}\pi r^2 h$$

But volume is maximum (given)


$$\therefore h = r$$

$$\begin{aligned}\therefore \text{R required volume} &= \frac{2}{3}\pi r^3 - \frac{1}{3}\pi r^3 \\ &= \frac{1}{3}\pi r^3\end{aligned}$$

4. An oil funnel of tin sheet consists of a cylindrical portion 10 cm long attached to a frustum of a cone. If the total height is 22 cm, the diameter of the cylindrical portion be 8 cm and the diameter of the top of the

tin sheet required to make the funnel.

Solution :



Area of tin sheet required to make the funnel

where $R = 9$ cm $r = 4$ cm, $H = 10$ cm

$$\begin{aligned}l &= \sqrt{(R-r)^2 + h^2} \\ &= \sqrt{25 + 144} \\ &= \sqrt{169} \\ &= 13\end{aligned}$$

= CSA of Frustum + CSA of Cylinder

$$= \pi (R + r) l + 2\pi r H$$

$$= \pi [13 \times 13 + 2 \times 4 \times 10]$$

$$= \frac{22}{7} [169 + 80]$$

$$= \frac{22}{7} \times 249$$

$$= \frac{5478}{7}$$

$$= 782.57 \text{ cm}^3$$

5. Find the number of coins, 1.5 cm in diameter and 2 mm thick, to be melted to form a right circular cylinder of height 10 cm and diameter 4.5 cm.

Solution :

Given diameter of coin = 1.5 cm

(smaller cylinder)

$$\therefore r = \frac{1.5}{2} = 0.75 \text{ cm}$$

$$h = 2 \text{ mm} = 0.2 \text{ cm}$$

Also, diameter of bigger cylinder = 4.5 cm

$$R = 2.25 \text{ cm}$$

$$H = 10 \text{ cm}$$

\therefore Number of Coins =

$$\frac{\text{Volume of largest cylinder}}{\text{Volume of smallest cylinder}}$$

$$\begin{aligned}
 &= \frac{\pi R^2 H}{\pi r^2 h} \\
 &= \frac{9}{4} \times \frac{9}{4} \times 10 \\
 &= \frac{3}{4} \times \frac{3}{4} \times \frac{2}{10} \\
 &= 450 \text{ coins}
 \end{aligned}$$

6. A hollow metallic cylinder whose external radius is 4.3 cm and internal radius is 1.1 cm and whole length is 4 cm is melted and recast into a solid cylinder of 12 cm long. Find the diameter of solid cylinder.

Solution :

Hollow Cylinder

$$R = 4.3 \text{ cm}$$

$$r = 1.1 \text{ cm}$$

$$H = 4 \text{ cm}$$

Solid Cylinder

$$h = 12 \text{ cm}$$

$$d = ?$$

When hollow cylinder is melted & recast into a solid cylinder,

Volume of hollow cylinder = Volume of solid cylinder

$$\Rightarrow \pi H (R^2 - r^2) = \pi r^2 h$$

$$\Rightarrow 4 [(4.3)^2 - (1.1)^2] = r^2 \times 12$$

$$\Rightarrow r^2 = \frac{4(17.28)}{12}$$

$$r^2 = \frac{17.28}{3}$$

$$= 5.76$$

$$r = 2.4$$

\therefore Diameter of solid cylinder

$$= 2r$$

$$= 4.8 \text{ cm}$$

7. The slant height of a frustum of a cone is 4 m and the perimeter of circular ends are 18 m and 16 m. Find the cost of painting its curved surface area at ₹100 per sq. m.

Solution :

In a frustum of a cone,

$$l = 4 \text{ cm}, 2\pi R = 18, 2\pi r = 16$$

$$\Rightarrow R = \frac{9}{\pi} \quad r = \frac{8}{\pi}$$

\therefore CSA of frustum of a cone

$$= \pi l (R + r)$$

$$= \pi \times 4 \left(\frac{9}{\pi} + \frac{8}{\pi} \right)$$

$$= 4 \times 17$$

$$= 68 \text{ m}^2$$

\therefore Cost of painting at ₹ 100/m²

8. A hemi-spherical hollow bowl has material of volume $\frac{436\pi}{3}$ cubic cm. Its external diameter is 14 cm. Find its thickness.

Solution :

In a hollow hemisphere,

$$\text{Volume} = \frac{436\pi}{3} \text{ cm}^3$$

$$D = 14 \text{ cm}, R = 7 \text{ cm}, t = ?$$

$$\Rightarrow \frac{2}{3}\pi(R^3 - r^3) = \frac{436\pi}{3}$$

$$\Rightarrow 7^3 - r^3 = 218$$

$$\Rightarrow 343 - r^3 = 218$$

$$\therefore r^3 = 125$$

$$\therefore r = 5 \text{ cm}$$

\therefore thickness, $t = R - r$

$$= 7 - 5$$

$$= 2 \text{ cm}$$

9. The volume of a cone is $1005\frac{5}{7}$ cu. cm.
The area of its base is $201\frac{1}{7}$ sq. cm. Find
the slant height of the cone.

Solution :

$$\text{Given volume of a cone} = 1005\frac{5}{7} \text{ cm}^3$$

$$\text{\& base area} = 201\frac{1}{7} \text{ cm}^2$$

$$\therefore \frac{1}{3}\pi r^2 h = \frac{7040}{7} \text{ \& } \pi r^2 = \frac{1408}{7}$$

$$\Rightarrow \frac{1}{3} \times \frac{1408}{\cancel{7}} \times h = \frac{7040}{\cancel{7}}$$

$$\Rightarrow h = \frac{7040}{1408} \times 3$$

$$\Rightarrow h = 5 \times 3$$

$$\text{Also, } \pi r^2 = \frac{1408}{7}$$

$$\Rightarrow \frac{22}{7} \times r^2 = \frac{1408}{7}$$

$$\Rightarrow r^2 = \frac{1408}{7} = 64$$

$$\therefore r = 8$$

$$\therefore l = \sqrt{h^2 + r^2} = \sqrt{15^2 + 8^2} = \sqrt{225 + 64} \\ = \sqrt{289} = 17 \text{ cm}$$

$$\therefore \text{Slant height} = 17 \text{ cm}$$

10. A metallic sheet in the form of a sector of a circle of radius 21 cm has central angle of 216° . The sector is made into a cone by bringing the bounding radii together. Find the volume of the cone formed.

Solution :

$$\text{Given radius of sector} = 21 \text{ cm, } \theta = 216^\circ$$

$$\text{ie } R = 21 = l \text{ (slant height of cone)}$$

When the sector is made into a cone by bringing the radii together.

Length of arc of the sector = Perimeter of base of cone

$$\Rightarrow \frac{\theta}{360} \times 2\pi R = 2\pi r$$

$$\Rightarrow r = \frac{216}{360} \times 21$$

$$\Rightarrow r = \frac{63}{5} = 12.6 \text{ cm}$$

$$\therefore h = \sqrt{l^2 - r^2} \\ = \sqrt{21^2 - (12.6)^2} \\ = \sqrt{441 - 158.76} \\ = \sqrt{282.24} \\ = 16.8$$

\therefore Volume of the cone

$$\frac{1}{3} \times \frac{22}{7} \times \frac{4.2}{\cancel{7}} \times \frac{1.8}{\cancel{7}} \times \frac{1.8}{\cancel{7}} \times 16.8 \\ = 2794.176 \\ \approx 2794.18 \text{ cm}^3$$

PROBLEMS FOR PRACTICE

- A girl empties a cylindrical bucket, full of sand of base radius 18 cm and height 32 cm on the floor, to form a conical heap of sand. If the height of this heap is 24 cm, find the slant height of cone. **(Ans : 43.27 cm)**
- 12 cylindrical pillars of a building have to be cleaned. If the diameter of each pillar is 42 cm, height is 5m, What will be the cost of cleaning at the rate of Rs.5 per m^2 . **(Ans : Rs.396/-)**
- A conical tent of 56m base diameter requires 3080 m^2 of canvas for the cured surface. Find its height. **(Ans : 21m)**

4. A solid is in the form of a cylinder with hemispherical ends. The total height of the solid is 19cm and the diameter of the cylinder is 7 cm. Find the surface area of the solid. **(Ans 418 m²)**

5. A farmer connects a pipe of internal diameter 20 cm from a Canal into a cylindrical tank which is 10m in diameter and 2m deep. If the water flows through the pipe at the rate of 4 Km/hr, in how much time will the tank be completely filled ?

(Ans : 1 hr, 15min)

6. A toy is in the form of a cone of base radius 3.5 cm mounted on a hemisphere of base diameter 7 cm. If the total height of the toy is 15.5 cm, find the total surface area of the toy. **(Ans : 214.5cm²)**

has water upto a height of 9 cm. A metal cube of 8 cm edge is immersed completely. By how much, the water level rise in the tube ? **(Ans : 1.63 cm)**

8. A vessel is in the form of a cone. Its height is 8 cm and radius of its top which is open is 5 cm. It is filled with water up to the rim. When lead shots, each of which is a sphere of diameter 1cm are dropped into vessel, one fourth of the water flows out. Find the number of lead shots dropped into the vessel.

(Ans : (100))

9. A hollow spherical shell has an inner radius of 8cm. If the volume of material is $\frac{1952\pi}{3}$ C.C, Find the thickness of the shell.

(Ans : 2cm)

10. Find the length of arc of the sector formed by opening out a cone of base radius 8cm. What is the central angle if the height of the cone is 6cm.

(Ans : 280°, 50 $\frac{2}{7}$ cm)

11. Find the capacity of a bucket having the radius of the top as 36cm and that of the bottom as 12cm, depth is 35cm.

(Ans 68640 cm³)

12. Water flows through a cylindrical pipe of internal radius 3.5 cm at 5m per sec. Find the volume of water in litres discharged by the pipe in 1 min. **(Ans : 1155 litres)**

13. A rectangular sheet of metal foil with dimension 66 cm × 12 cm is rolled to form a cylinder of height 12cm. Find the volume of the cylinder. **(Ans : 4158 cm³)**

14. Using clay, a student made a right circular cone of height 48cm and base radius 12cm. Another student reshapes it in the form of a sphere. Find the radius of sphere.

(Ans : 12 cm)

15. A solid sphere of diameter 28 cm is melted and recast into smaller solid cones each of diameter $4\frac{2}{3}$ cm and height 3cm. Find the number of cones so formed.

(Ans : 672)

OBJECTIVE TYPE QUESTIONS

1. The radii of 2 cylinders are in the ratio 2 : 3 and their heights are in the ratio 5 : 3. Then the ratio of their volumes is

a) 10 : 9 b) 20 : 27 c) 7 : 6

d) 5 : 2 **(Ans : (b))**

2. If the volume of sphere is $\frac{9}{16} \text{ cm}^3$, its radius is
 a) $\frac{4}{3} \text{ cm}$ b) $\frac{3}{4} \text{ cm}$
 c) $\frac{3}{2} \text{ cm}$ d) $\frac{2}{3} \text{ cm}$
(Ans : (b))
3. The height of a cone whose slant height 26cm and the diameter of the base is 10cm, is
 a) 24.5 cm b) 26.5 cm c) 25.5 cm
 d) 27.5 cm **(Ans : (c))**
4. The total surface area of a cylinder whose height is half the radius is
 a) $6\pi r^2$ b) $8\pi r^2$ c) $2\pi r^2$
 d) $3\pi r^2$ **(Ans : (d))**
5. The base area of a cone is 80cm^2 . If its height is 9cm, then its volume is
 a) 720cm^3 b) $720\pi \text{ cm}^3$
 c) 240 cm^3 d) none **(Ans : (c))**
6. A well of diameter 2.1m is dug to a depth of 4m. The volume of the earth removed is
 a) $4.4\pi \text{ m}^3$ b) 44.1 m^3
 c) 0.441 m^3 d) $4.41 \pi \text{ m}^3$
(Ans : (d))
7. The volume of a hemisphere is 18π . Its radius is
 a) 4cm b) 3cm c) 2cm
 d) 6cm **(Ans : (b))**
8. If a rectangle ABCD is folded by bringing AB and CD together to form a cylinder, then the height of the cylinder is
 a) BC b) AD c) AB
 d) none **(Ans : (c))**
9. The CSA of a solid hemisphere if the TSA of the solid hemisphere is $12\pi \text{ cm}^2$, is
 a) 8π b) 36π c) 6π d) 24π
(Ans : (a))
10. The CSA of a cone whose radius x cm, height y cm is
 a) $\pi r l$ b) $\pi r \sqrt{x^2 + y^2}$
 c) $\pi y \sqrt{x^2 + y^2}$ d) $\pi x \sqrt{x^2 + y^2}$
(Ans : (d))

CHAPTER 8

STATISTICS AND PROBABILITY

I. MEASURES OF DISPERSION

Key Points

- ✓ Measures of Variation (or) Dispersion of a data provide an idea of how observations spread out (or) scattered throughout the data.
- ✓ Different Measures of Dispersion are
 1. Range 2. Mean deviation 3. Quartile deviation 4. Standard deviation 5. Variance 6. Coefficient of Variation

- ✓ Range $R = L - S$

- ✓ Coefficient of range $= \frac{L - S}{L + S}$ where L - Largest value; S - Smallest value.

- ✓ The mean of the squares of the deviations from the mean is called Variance. It is denoted by σ^2 .

$$\text{Variance } \sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

- ✓ The positive square root of Variance is called Standard deviation.

$$\text{Standard deviation } \sigma = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}}$$

- ✓ Formula Table for Standard Deviation (σ).

Data Type	Direct Method	Mean Method	Assumed mean method	Step Deviation method
Ungrouped Data	$\sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2}$	$\sqrt{\frac{\sum d^2}{n}}$	$\sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2}$	$\sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \times C$
Grouped Data	—	$\sqrt{\frac{\sum fd^2}{N}}$ $N = \sum f$	$\sqrt{\frac{\sum fd^2}{N} - \left(\frac{\sum fd}{N}\right)^2}$ $N = \sum f$	$\sqrt{\frac{\sum fd^2}{N} - \left(\frac{\sum fd}{N}\right)^2} \times C$ $N = \sum f$

- ✓ Standard deviation of first 'n' natural numbers

$$\sigma = \sqrt{\frac{n^2 - 1}{12}}$$

- ✓ The value of SD will not be changed if we add (or) subtract some fixed constant to all the values.
- ✓ When we multiply each value of a data by a constant, the value of SD is also multiplied by the same constant.

Example 8.1

Find the range and coefficient of range of the following data: 25, 67, 48, 53, 18, 39, 44.

Solution :

Largest value L = 67; Smallest value S = 18

Range R = L - S = 67 - 18 = 49

Coefficient of range = $\frac{L - S}{L + S}$

Coefficient of range = $\frac{67 - 18}{67 + 18} = \frac{49}{85} = 0.576$

Example 8.2

Find the range of the following distribution.

Age (in years)	16-18	18-20	20-22	22-24	24-26	26-28
Number of students	0	4	6	8	2	2

Solution :

Here Largest value L = 28

Smallest value S = 18

Range R = L - S

R = 28 - 18 = 10 Years.

Example 8.3

The range of a set of data is 13.67 and the largest value is 70.08. Find the smallest value.

Solution :

Range R = 13.67

Largest value L = 70.08

Range R = L - S

13.67 = 70.08 - S

S = 70.08 - 13.67 = 56.41

Therefore, the smallest value is 56.41.

Example 8.4

The number of televisions sold in each day of a week are 13, 8, 4, 9, 7, 12, 10. Find its standard deviation.

Solution :

x_i	x_i^2
13	169
8	64
4	16
9	81
7	49
12	144
10	100
$\Sigma x_i = 63$	$\Sigma x_i^2 = 623$

Standard deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\Sigma x_i^2}{n} - \left(\frac{\Sigma x_i}{n}\right)^2} \\ &= \sqrt{\frac{623}{7} - \left(\frac{63}{7}\right)^2} \\ &= \sqrt{89 - 81} = \sqrt{8}\end{aligned}$$

Hence, $\sigma \approx 2.83$

Example 8.5

The amount of rainfall in a particular season for 6 days are given as 17.8 cm, 19.2 cm, 16.3 cm, 12.5 cm, 12.8 cm and 11.4 cm. Find its standard deviation.

Solution :

Arranging the numbers in ascending order we get, 11.4, 12.5, 12.8, 16.3, 17.8, 19.2. [Number of observations $n = 6$]

$$\begin{aligned}\text{Mean} &= \frac{11.4 + 12.5 + 12.8 + 16.3 + 17.8 + 19.2}{6} \\ &= \frac{90}{6} = 15\end{aligned}$$

x_i	$d_i = x_i - \bar{x}$ $= x - 15$	d_i^2
11.4	-3.6	12.96
12.5	-2.5	6.25
12.8	-2.2	4.84
16.3	1.3	1.69
17.8	2.8	7.84
19.2	4.2	17.64
		$\Sigma d_i^2 = 51.22$

$$\begin{aligned}\text{Standard deviation } \sigma &= \sqrt{\frac{\Sigma d_i^2}{n}} \\ &= \sqrt{\frac{51.22}{6}} = \sqrt{8.53}\end{aligned}$$

$$\text{Hence, } \sigma \approx 2.9$$

Example 8.6

The marks scored by 10 students in a class test are 25, 29, 30, 33, 35, 37, 38, 40, 44, 48. Find the standard deviation.

Solution :

The mean of marks is 35.9 which is not an integer. Hence we take assumed mean, $A = 35$, $n = 10$.

x_i	$d_i = x_i - A$ $d_i = x_i - 35$	d_i^2
25	-10	100
29	-6	36
30	-5	25
33	-2	4
35	0	0
37	2	4
38	3	9
40	5	25
44	9	81
48	13	169
	$\Sigma d_i = 9$	$\Sigma d_i^2 = 453$

Example 8.7

The amount that the children have spent for purchase are 5, 10, 15, 20, 25, 30, 35, 40. Using step deviation method, find the standard deviation of the amount they have spent.

Solution :

We note that all the observations are divisible by 5. Hence we can use the step deviation method. Let the Assumed mean $A = 20$, $n = 8$.

x_i	$d_i = x_i - A$ $d_i = x_i - 20$	$d_i = \frac{x_i - A}{c}$ $c = 5$	d_i^2
5	-15	-3	9
10	-10	-2	4
15	-5	-1	1
20	0	0	0
25	5	1	1
30	10	2	4
35	15	3	9
40	20	4	16
		$\Sigma d_i = 4$	$\Sigma d_i^2 = 44$

Standard Deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum d_i^2}{n} - \left(\frac{\sum d_i}{n}\right)^2} \times c \\ &= \sqrt{\frac{44}{8} - \left(\frac{4}{8}\right)^2} \times 5 = \sqrt{\frac{11}{2} - \frac{1}{4}} \times 5 \\ &= \sqrt{5.5 - 0.25} \times 5 = 2.29 \times 5\end{aligned}$$

$$\sigma \approx 11.45$$

Example 8.8

Find the standard deviation of the following data 7, 4, 8, 10, 11. Add 3 to all the values then find the standard deviation for the new values.

Solution :

Arranging the values in ascending order we get, 4, 7, 8, 10, 11 and $n = 5$

x_i	x_i^2
4	16
7	49
8	64
10	100
11	121
$\sum x_i = 40$	$\sum x_i^2 = 350$

Standard deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2} \\ &= \sqrt{\frac{350}{5} - \left(\frac{40}{5}\right)^2} \\ \sigma &= \sqrt{6} \approx 2.45\end{aligned}$$

When we add 3 to all the values, we get the new values as 7, 10, 11, 13, 14.

x_i	x_i^2
7	49
10	100
11	121
13	169
14	196
$\sum x_i = 55$	$\sum x_i^2 = 635$

Standard deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2} \\ &= \sqrt{\frac{635}{5} - \left(\frac{55}{5}\right)^2} \\ \sigma &= \sqrt{6} \approx 2.45\end{aligned}$$

From the above, we see that the standard deviation will not change when we add some fixed constant to all the values.

Example 8.9

Find the standard deviation of the data 2, 3, 5, 7, 8. Multiply each data by 4. Find the standard deviation of the new values.

Solution :Given, $n = 5$

x_i	x_i^2
2	4
3	9
5	25
7	49
8	64
$\sum x_i = 25$	$\sum x_i^2 = 151$

Standard deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2} \\ \sigma &= \sqrt{\frac{151}{5} - \left(\frac{25}{5}\right)^2} \\ &= \sqrt{30.2 - 25} \\ &= \sqrt{5.2} \approx 2.28\end{aligned}$$

When we multiply each data by 4, we get the new values as 8, 12, 20, 28, 32.

x_i	x_i^2
8	64
12	144
20	400
28	784
32	1024
$\sum x_i = 100$	$\sum x_i^2 = 2416$

Standard deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2} \\ &= \sqrt{\frac{2416}{5} - \left(\frac{100}{5}\right)^2} \\ &= \sqrt{483.2 - 400} \\ &= \sqrt{83.2} \\ \sigma &= \sqrt{16 \times 5.2} \\ &= 4\sqrt{5.2} \approx 9.12\end{aligned}$$

From the above, we see that when we multiply each data by 4 the standard deviation also get multiplied by 4.

Example 8.10

Find the mean and variance of the first n natural numbers.

Solution :

$$\text{Mean } \bar{x} = \frac{\text{Sum of all the observations}}{\text{Number of observations}}$$

$$= \frac{\sum x_i}{n} = \frac{1+2+3+\dots+n}{n} = \frac{n(n+1)}{2 \times n}$$

$$\text{Mean } \bar{x} = \frac{n+1}{2}$$

$$\text{Variance } \sigma^2 = \frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n} \right)^2$$

$$= \left[\frac{\sum x_i^2 = 1^2 + 2^2 + 3^2 + \dots + n^2}{(\sum x_i)^2 = (1+2+3+\dots+n)^2} \right]$$

$$= \frac{n(n+1)(2n+1)}{6 \times n} - \left[\frac{n(n+1)}{2 \times n} \right]^2$$

$$= \frac{2n^2 + 3n + 1}{6} - \frac{n^2 + 2n + 1}{4}$$

$$\text{Variance } \sigma^2 = \frac{4n^2 + 6n + 2 - 3n^2 - 6n - 3}{12}$$

$$= \frac{n^2 - 1}{12}$$

Example 8.11

48 students were asked to write the total number of hours per week they spent on watching television. With this information find the standard deviation of hours spent for watching television.

x	6	7	8	9	10	11	12
f	3	6	9	13	8	5	4

Solution :

x_i	f_i	$x_i f_i$	$d_i = x_i - \bar{x}$	d_i^2	$f_i d_i^2$
6	3	18	-3	9	27
7	6	42	-2	4	24
8	9	72	-1	1	9
9	13	117	0	0	0
10	8	80	1	1	8
11	5	55	2	4	20
12	4	48	3	9	36
	$N=48$	$\sum x_i f_i = 432$	$\sum d_i = 0$		$\sum f_i d_i^2 = 124$

$$\text{Mean } \bar{x} = \frac{\sum x_i f_i}{N} = \frac{432}{48} = 9 \quad (\text{Since } N = \sum f_i)$$

Standard deviation

$$\sigma = \sqrt{\frac{\sum f_i d_i^2}{N}} = \sqrt{\frac{124}{48}} = \sqrt{2.58}$$

$$\sigma \approx 1.6$$

Example 8.12

The marks scored by the students in a slip test are given below.

x	4	6	8	10	12
f	7	3	5	9	5

Find the standard deviation of their marks.

Solution :

Let the assumed mean, $A = 8$

x_i	f_i	$d_i = x_i - A$	$f_i d_i$	$f_i d_i^2$
4	7	-4	-28	112
6	3	-2	-6	12
8	5	0	0	0
10	9	2	18	36
12	5	4	20	80
	$N=29$		$\sum f_i d_i = 4$	$\sum f_i d_i^2 = 240$

Standard deviation

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum f_i d_i^2}{N} - \left(\frac{\sum f_i d_i}{N}\right)^2} \\ &= \sqrt{\frac{240}{29} - \left(\frac{4}{29}\right)^2} = \sqrt{\frac{240 \times 29 - 16}{29 \times 29}} \\ \sigma &= \sqrt{\frac{6944}{29 \times 29}}; \sigma \approx 2.87\end{aligned}$$

Example 8.13

Marks of the students in a particular subject of a class are given below.

Marks	0-10	10-20	20-30	30-40
Number of students	8	12	17	14
Marks	40-50	50-60	60-70	-
Number of students	9	7	4	-

Find its standard deviation.

Solution :

Let the assumed mean, $A = 35$, $c = 10$

Marks	Mid value (x_i)	f_i	$d_i = \frac{x_i - A}{c}$	$d_i = \frac{x_i - A}{c}$	$f_i d_i$	$f_i d_i^2$
0-10	5	8	-30	-3	-24	72
10-20	15	12	-20	-2	-24	48
20-30	25	17	-10	-1	-17	17
30-40	35	14	0	0	0	0
40-50	45	9	10	1	9	9
50-60	55	7	20	2	14	28
60-70	65	4	30	3	12	36
		N=71			$\sum f_i d_i = -30$	$\sum f_i d_i^2 = 210$

$$\begin{aligned}\text{Standard deviation } \sigma &= c \times \sqrt{\frac{\sum f_i d_i^2}{N} - \left(\frac{\sum f_i d_i}{N}\right)^2} \\ \sigma &= 10 \times \sqrt{\frac{210}{71} - \left(\frac{-30}{71}\right)^2} = 10 \times \sqrt{\frac{210}{71} - \frac{900}{5041}} \\ &= 10 \times \sqrt{2.779}; \sigma \approx 16.67\end{aligned}$$

Example 8.14

The mean and standard deviation of 15 observations are found to be 10 and 5 respectively. On rechecking it was found that one of the observation with value 8 was incorrect. Calculate the correct mean and standard deviation if the correct observation value was 23?

Solution :

$$n = 15, \bar{x} = 10, \sigma = 5;$$

$$\bar{x} = \frac{\sum x}{n}; \sum x = 15 \times 10 = 150$$

Wrong observation value = 8,

Correct observation value = 23.

Correct total = $150 - 8 + 23 = 165$

$$\text{Correct mean } \bar{x} = \frac{165}{15} = 11$$

$$\text{Standard deviation } \sigma = \sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2}$$

$$\text{Incorrect value of } \sigma = 5 = \sqrt{\frac{\sum x^2}{15} - (10)^2}$$

$$25 = \frac{\sum x^2}{15} - 100 \text{ gives } \frac{\sum x^2}{15} = 125$$

Incorrect value of $\sum x^2 = 1875$

Correct value of $\sum x^2 = 1875 - 8^2 + 23^2 = 2340$

$$\text{Correct standard deviation } \sigma = \sqrt{\frac{2340}{15} - (11)^2}$$

$$\sigma = \sqrt{156 - 121} = \sqrt{35} \quad \sigma \approx 5.9$$

EXERCISE 8.1

1. Find the range and coefficient of range of the following data.

(i) 63, 89, 98, 125, 79, 108, 117, 68

(ii) 43.5, 13.6, 18.9, 38.4, 61.4, 29.8

Solution:

$$\begin{aligned} \text{i) Range} &= L - S \\ &= 125 - 63 \\ &= 62 \end{aligned}$$

$$\begin{aligned} \text{Coefficient of range} &= \frac{L - S}{L + S} \\ &= \frac{125 - 63}{125 + 63} \\ &= \frac{62}{188} \\ &= 0.33 \end{aligned}$$

$$\begin{aligned} \text{ii) Range} &= L - S \\ &= 61.4 - 13.6 \\ &= 47.8 \end{aligned}$$

$$\begin{aligned} \text{Coefficient of range} &= \frac{L - S}{L + S} \\ &= \frac{61.4 - 13.6}{61.4 + 13.6} \\ &= \frac{47.8}{75} \\ &= 0.64 \end{aligned}$$

2. If the range and the smallest value of a set of data are 36.8 and 13.4 respectively, then find the largest value.

Solution :

Given ; range = 36.8

Smallest value = 13.4

$$\therefore R = L - S$$

$$\therefore L = R + S$$

$$= 36.8 + 13.4$$

$$= 50.2$$

3. Calculate the range of the following data.

Income	400-450	450-500	500-550
Number of workers	8	12	30
Income	550-600	600-650	-
Number of workers	21	6	-

Solution :

Here, Largest value = $L = 650$

Smallest value = $S = 450$

$$\begin{aligned} \therefore \text{Range} &= L - S \\ &= 650 - 450 \\ &= 200 \end{aligned}$$

4. A teacher asked the students to complete 60 pages of a record note book. Eight students have completed only 32, 35, 37, 30, 33, 36, 35 and 37 pages. Find the standard deviation of the pages yet to be completed by them.

Solution :

The pages yet to be completed by them are

60-32, 60-35, 60-37, 60-30, 60-33, 60-36, 60-35, 60-37, 60-37

= 28, 25, 23, 30, 27, 24, 25, 23

To find the SD of the data 28, 25, 23, 30, 27, 24, 25, 23

Arrange them in ascending order

x	$d = x - A$	d^2
23	-2	4
23	-2	4
24	-1	1
25	0	0
25	0	0

27	2	4
28	3	9
30	5	25
	5	47
	$\Sigma d = 5$	$\Sigma d^2 = 47$

$$\begin{aligned}
 \therefore \sigma &= \sqrt{\frac{\Sigma d^2}{n} - \left(\frac{\Sigma d}{n}\right)^2} \\
 &= \sqrt{\frac{47}{8} - \left(\frac{5}{8}\right)^2} \\
 &= \sqrt{\frac{47}{8} - \frac{25}{64}} \\
 &= \sqrt{\frac{376 - 25}{64}} \\
 &= \frac{\sqrt{351}}{8} \\
 &= \frac{18.74}{8} \\
 &= 2.34
 \end{aligned}$$

5. Find the variance and standard deviation of the wages of 9 workers given below:

₹310, ₹290, ₹320, ₹280, ₹300, ₹290, ₹320, ₹310, ₹280.

Solution :

Given wages of a workers are ₹310, ₹290, ₹320, ₹280, ₹300, ₹290, ₹320, ₹310, ₹280

To find the variance and SD, arrange them in ascending order.

x	$d = \frac{x - 300}{10}$	d^2
280	-2	4
280	-2	4
290	-1	1

290	-1	1
300	0	0
310	1	1
310	1	1
320	2	4
320	2	4
	0	20
	$\Sigma d = 0$	$\Sigma d^2 = 20$

$$d = \frac{x - A}{c}$$

A - Assumed Mean C - Common divisor

$$\begin{aligned}
 \sigma^2 &= \frac{\Sigma d^2}{n} - \left(\frac{\Sigma d}{n}\right)^2 \times c^2 \\
 &= \frac{20}{9} \\
 &= \frac{2000}{9} \\
 \sigma^2 &= 222.2
 \end{aligned}$$

$$\therefore \text{Variance} = 222.2$$

$$\begin{aligned}
 \therefore \text{S.D} &= \sqrt{222.2} \\
 &= 14.906 \\
 &\approx 14.91
 \end{aligned}$$

6. A wall clock strikes the bell once at 1 o' clock, 2 times at 2 o' clock, 3 times at 3 o' clock and so on. How many times will it strike in a particular day. Find the standard deviation of the number of strikes the bell make a day.

Solution :

A clock strikes bell at 1 o' clock once
twice at 2 o' clock,
3 times at 3 o' clock

\therefore Number of times it strikes in a particular day

$$= 2(1 + 2 + 3 + \dots + 12)$$

$$= 2\left(\frac{12 \times 13}{2}\right)$$

$$= 156 \text{ times}$$

To find the S.D of 2 (1, 2, 3,12)

$$= 2\left[\sqrt{\frac{n^2 - 1}{12}}\right]$$

$$= 2\left[\sqrt{\frac{144 - 1}{12}}\right]$$

$$= 2\sqrt{\frac{143}{12}} = 2\sqrt{11.91}$$

$$= 2(3.45)$$

$$= 6.9$$

natural numbers.

Solution :

SD of first 21 natural numbers

$$= \sqrt{\frac{n^2 - 1}{12}}$$

$$= \sqrt{\frac{441 - 1}{12}}$$

$$= \sqrt{\frac{440}{12}}$$

$$= \sqrt{36.66}$$

$$= 6.0547$$

$$\approx 6.05$$

- 8. If the standard deviation of a data is 4.5 and if each value of the data is decreased by 5, then find the new standard deviation.**

Solution :

Given, S.D of a data = 4.5

Since each value is decreased by 5, then the new SD = 4.5

(\because S.D will not be changed when we add (or) subtract fixed constant to all the values of the data).

- 9. If the standard deviation of a data is 3.6 and each value of the data is divided by 3, then find the new variance and new standard deviation.**

Solution :

Given, S.D of a data = 3.6

Since each value is divided by 3 then the new S.D = $\frac{3.6}{3}$

$$= 1.2$$

New Variance = $(1.2)^2$

$$= 1.44$$

- 10. The rainfall recorded in various places of five districts in a week are given below.**

Rainfall (in mm)	45	50	55	60	65	70
Number of places	5	13	4	9	5	4

Find its standard deviation.

Solution :

x	f	$d = \frac{x - 60}{5}$	d^2	$f \cdot d$	$f \cdot d^2$
45	5	-3	9	-15	45
50	13	-2	4	-26	52
55	4	-1	1	-4	4
60	9	0	0	0	0
65	5	1	1	5	5
70	4	2	4	8	16
				-32	122

$$\Sigma f = N = 40, \quad \Sigma fd = -32 \quad \Sigma f \cdot d^2 = 122$$

$$c = 5$$

$$\begin{aligned}
 \therefore \sigma &= \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2} \times c \\
 &= \sqrt{\frac{122}{40} - \left(\frac{-32}{40}\right)^2} \times 5 \\
 &= \sqrt{\frac{122}{40} - \frac{1024}{1600}} \times 5 \\
 &= \sqrt{\frac{4880 - 1024}{1600}} \times 5 \\
 &= \frac{\sqrt{3856}}{40} \times 5 \\
 &= \frac{62.096}{8} = 7.76
 \end{aligned}$$

$$\therefore \text{S.D} = 7.76$$

11. In a study about viral fever, the number of people affected in a town were noted as

Age in years	0-10	10-20	20-30	30-40
Number of people affected	3	5	16	18
Age in years	40-50	50-60	60-70	-
Number of people affected	12	7	4	

Solution :

C.I	mid value (x)	f	$d = \frac{x-35}{10}$	d ²	f · d	f · d ²
0-10	5	3	-3	9	-9	27
10-20	15	5	-2	4	-10	20
20-30	25	16	-1	1	-16	16
30-40	35-A	18	0	0	0	0
40-50	45	12	1	1	12	12
50-60	55	7	2	4	14	28
60-70	65	4	3	9	12	36
					3	139

$$\therefore \sum f = 65, \sum fd = 3, \sum fd^2 = 139 \text{ and } c = 10$$

$$\begin{aligned}
 \therefore \sigma &= \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2} \times c \\
 &= \sqrt{\frac{139}{65} - \left(\frac{3}{65}\right)^2} \times 10 \\
 &= \sqrt{\frac{139}{65} - \frac{9}{65^2}} \times 10 \\
 &= \sqrt{\frac{9035 - 9}{65^2}} \times 10 \\
 &= \frac{\sqrt{9026}}{65} \times 10 \\
 &= \frac{95.005}{65} \times 10 \\
 &= 1.46 \times 10 \\
 &= 14.6
 \end{aligned}$$

12. The measurements of the diameters (in cms) of the plates prepared in a factory are given below. Find its standard deviation.

Diameter (cm)	21-24	25-28	29-32	33-36	37-40	41-44
Number of plates	15	18	20	16	8	7

Solution :

C.I	mid value (x)	f	$d = \frac{x-34.5}{4}$	d ²	f · d	f · d ²
21-24	22.5	15	-3	9	-45	135
25-28	26.5	18	-2	4	-36	72
29-32	30.5	20	-1	1	-20	20
33-36	34.5	16	0	0	0	0
37-40	38.5	8	1	1	8	8
41-44	42.5	7	2	4	14	28
		84			-79	263

$$\therefore \sum f = 94, \sum fd = -79, \sum fd^2 = 263, c = 4$$

$$\begin{aligned}
 \therefore \sigma &= \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2} \times c \\
 &= \sqrt{\frac{263}{84} - \left(\frac{-79}{84}\right)^2} \times 4 \\
 &= \sqrt{\frac{263}{84} - \frac{6241}{84^2}} \times 4 \\
 &= \sqrt{\frac{22092 - 6241}{84^2}} \times 4 \\
 &= \frac{\sqrt{15851}}{84} \times 4 \\
 &= \frac{125.9}{21} \\
 &= 5.995 \\
 &\approx 6
 \end{aligned}$$

plete a 100 meter race are given below.
Find its standard deviation

Time taken (sec)	8.5- 9.5	9.5- 10.5	10.5- 11.5	11.5- 12.5	12.5- 13.5
Number of students	6	8	17	10	9

Solution :

C.I	mid value (x)	f	d = x - 11	d ²	f · d	f · d ²
8.5-9.5	9	6	-2	4	-12	24
9.5-10.5	10	8	-1	1	-8	8
10.5-11.5	11	17	0	0	0	0
11.5-12.5	12	10	1	1	10	10
12.5-13.5	13	9	2	4	18	36
		50			8	78

$$\therefore \sum f = 50, \sum fd = 8, \sum fd^2 = 78 \text{ and } c = 1$$

$$\begin{aligned}
 \therefore \sigma &= \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2} \\
 &= \sqrt{\frac{78}{50} - \left(\frac{8}{50}\right)^2} \\
 &= \sqrt{\frac{78}{50} - \frac{64}{50^2}} \\
 &= \sqrt{\frac{3900 - 64}{50^2}} \\
 &= \frac{\sqrt{3836}}{50} \\
 &= \frac{61.935}{65} \\
 &= 1.238 \\
 &\approx 1.24
 \end{aligned}$$

and standard deviation of their marks were found to be 60 and 15 respectively. Later on it was found that the scores 45 and 72 were wrongly entered as 40 and 27. Find the correct mean and standard deviation.

Solution :

$$\text{Given } n = 100, \bar{x} = 60, \sigma = 15$$

$$\therefore \frac{\sum x}{n} = 60$$

$$\Rightarrow \frac{\sum x}{100} = 60$$

$$\Rightarrow \sum x = 6000$$

$$\begin{aligned}
 \therefore \text{Corrected } \sum x &= 6000 - (40 + 27) + (45 + 72) \\
 &= 6000 - 67 + 117 \\
 &= 6050
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{Corrected mean} &= \frac{6050}{100} \\
 &= 60.5
 \end{aligned}$$

$$\text{Variance} = \sigma^2 = \frac{\sum x^2}{n} - \left(\frac{\sum x}{n} \right)^2$$

$$225 = \frac{\sum x^2}{100} - 60^2$$

$$\therefore \frac{\sum x^2}{100} = 3825$$

$$\Rightarrow \sum x^2 = 382500$$

$$\therefore \text{The correct } \sum x^2 = 382500$$

$$\therefore \text{Corrected } \sum x^2$$

$$= \text{Incorrect } \sum x^2 - 40^2 - 27^2 + 45^2 + 72^2$$

$$= 382500 - 1600 - 729 + 2025 + 5184$$

$$= 387380$$

$$\therefore \text{Corrected } \sigma^2 = \frac{\text{Corrected } \sum x^2}{n} - (\text{Corr. mean})^2$$

$$= \frac{387380}{n} - (60.5)^2$$

$$= 3873.80 - 3660.25$$

$$= 213.55$$

$$\therefore \text{Corrected SD} = \sqrt{213.55}$$

$$= 14.6$$

15. The mean and variance of seven observations are 8 and 16 respectively. If five of these are 2, 4, 10, 12 and 14, then find the remaining two observations.

Solution :

$$\text{Given } n = 7, \bar{x} = 8, \sigma^2 = 16$$

5 of the observations are 2, 4, 10, 12, 14

Let the remaining 2 observations be a, b .

$$\therefore \bar{x} = 8 \Rightarrow \frac{\sum x}{n} = 8$$

$$\Rightarrow \frac{42 + a + b}{7} = 8$$

$$\Rightarrow a + b = 56 - 42$$

$$\Rightarrow a + b = 14 \quad \dots(1)$$

$$\text{Also, } \sigma^2 = 16$$

$$\Rightarrow \frac{\sum x^2}{n} - \left(\frac{\sum x}{n} \right)^2 = 16$$

$$\Rightarrow \frac{\sum x^2}{7} - 8^2 = 16$$

$$\Rightarrow \frac{\sum x^2}{7} = 80$$

$$\Rightarrow \sum x^2 = 560$$

$$\Rightarrow 2^2 + 4^2 + 10^2 + 12^2 + 14^2 + a^2 + b^2 = 560$$

$$\Rightarrow 460 + a^2 + b^2 = 560$$

$$\Rightarrow a^2 + b^2 = 100$$

$$\Rightarrow a^2 + (14 - a)^2 = 100 \quad (\text{from (1)})$$

$$\Rightarrow a^2 + 196 + a^2 - 28a = 100$$

$$\Rightarrow 2a^2 - 28a + 96 = 0$$

$$\Rightarrow a^2 - 14a + 48 = 0$$

$$\Rightarrow (a - 8)(a - 6) = 0$$

$$a = 8, \quad a = 6$$

$$\therefore b = 6, \quad b = 8$$

II. COEFFICIENT OF VARIATION :

Key Points

- ✓ Coefficient of variation, $CV = \frac{\sigma}{\bar{x}} \times 100$.
- ✓ If the C.V value is less, then the observations of \bar{x} corresponding data are consistent.
- ✓ If the C.V value is more, then the observations of corresponding data are inconsistent.

Example 8.15

The mean of a data is 25.6 and its coefficient of variation is 18.75. Find the standard deviation.

Solution :

Mean $\bar{x} = 25.6$,

Coefficient of variation, C.V. = 18.75

Coefficient of variation, $C.V. = \frac{\sigma}{\bar{x}} \times 100\%$

$$18.75 = \frac{\sigma}{25.6} \times 100 ; \quad \sigma = 4.8$$

Example 8.16

The following table gives the values of mean and variance of heights and weights of the 10th standard students of a school.

	Height	Weight
Mean	155 cm	46.50 kg ²
Variance	72.25 cm ²	28.09 kg ²

Which is more varying than the other?

Solution :

For comparing two data, first we have to find their coefficient of variations

Mean $\bar{x}_1 = 155$ cm, variance $\sigma_1^2 = 72.25$ cm²

Therefore standard deviation $\sigma_1 = 8.5$

Coefficient of variation $C.V_1 = \frac{\sigma_1}{\bar{x}_1} \times 100\%$

$$C.V_1 = \frac{8.5}{155} \times 100\% = 5.48\% \quad (\text{for heights})$$

Mean $\bar{x}_2 = 46.50$ kg, variance $\sigma_2^2 = 28.09$ kg²

Standard deviation $\sigma_2 = 5.3$ kg

Coefficient of variation $C.V_2 = \frac{\sigma_2}{\bar{x}_2} \times 100\%$

$$C.V_2 = \frac{5.3}{46.50} \times 100\% = 11.40\% \quad (\text{for weights})$$

$C.V_1 = 5.48\%$ and $C.V_2 = 11.40\%$

Since $C.V_2 > C.V_1$, the weight of the students is

Example 8.17

The consumption of number of guava and orange on a particular week by a family are given below.

Number of Guavas	3	5	6	4	3	5	4
Number of Oranges	1	3	7	9	2	6	2

Which fruit is consistently consumed by the family?

Solution :

First we find the coefficient of variation for guavas and oranges separately.

x_i	x_i^2
3	9
5	25
6	36
4	16
3	9
5	25
4	16
$\Sigma x_i = 30$	$\Sigma x_i^2 = 136$

Number of guavas, $n = 7$

$$\text{Mean } \bar{x}_1 = \frac{30}{7} = 4.29$$

$$\text{Standard deviation } \sigma_1 = \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2}$$

$$\sigma_1 = \sqrt{\frac{136}{7} - \left(\frac{30}{7}\right)^2} = \sqrt{19.43 - 18.40} \approx 1.01$$

Coefficient of variation for guavas

$$C.V_1 = \frac{\sigma_1}{\bar{x}_1} \times 100\% = \frac{1.01}{4.29} \times 100\% = 23.54\%$$

x_i	x_i^2
1	1
3	9
7	49
9	81
6	36
2	4
$\sum x_i = 30$	$\sum x_i^2 = 184$

Number of oranges $n = 7$

$$\text{Mean } \bar{x}_2 = \frac{30}{7} = 4.29$$

$$\text{Standard deviation } \sigma_2 = \sqrt{\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2}$$

$$\sigma_2 = \sqrt{\frac{184}{7} - \left(\frac{30}{7}\right)^2} = \sqrt{26.29 - 18.40} = 2.81$$

Coefficient of variation for oranges :

$$C.V_2 = \frac{\sigma_2}{\bar{x}_2} \times 100\% = \frac{2.81}{4.29} \times 100\% = 65.50\%$$

$$C.V_1 = 23.54\% \text{ and } C.V_2 = 65.50\%$$

Since $C.V_1 < C.V_2$, we can conclude that the consumption of guavas is more consistent than oranges.

EXERCISE 8.2

1. The standard deviation and mean of a data are 6.5 and 12.5 respectively. Find the coefficient of variation.

Solution :

$$\text{Given } \sigma = 6.5, \bar{x} = 12.5$$

$$\begin{aligned} \therefore C.V &= \frac{\sigma}{\bar{x}} \times 100 \\ &= \frac{6.5}{12.5} \times 100 \\ &= \frac{13}{25} \times 100 \\ &= 13 \times 4 \\ &= 52\% \end{aligned}$$

2. The standard deviation and coefficient of variation of a data are 1.2 and 25.6 respectively. Find the value of mean.

Solution :

$$\text{Given } \sigma = 1.2, C.V = 25.6$$

$$\begin{aligned} \therefore C.V &= \frac{\sigma}{\bar{x}} \times 100 \\ 25.6 &= \frac{1.2}{\bar{x}} \times 100 \\ \Rightarrow \bar{x} &= \frac{120}{25.6} \\ \bar{x} &= 4.69 \end{aligned}$$

3. If the mean and coefficient of variation of a data are 15 and 48 respectively, then find the value of standard deviation.

Solution :

$$\text{Given } \bar{x} = 15, CV = 48, \sigma = ?$$

$$\therefore \text{C.V} = \frac{\sigma}{x} \times 100$$

$$\Rightarrow 48 = \frac{\sigma}{15} \times 100$$

$$\sigma = \frac{15 \times 48}{100} = \frac{720}{100} = 7.2$$

4. If $n = 5$, $\bar{x} = 6$, $\sum x^2 = 765$, then calculate the coefficient of variation.

Solution :

Given $n = 5$, $\bar{x} = 6$, $\sum x^2 = 765$, $CV = ?$

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2} \\ &= \sqrt{\frac{765}{5} - (6)^2} \\ &= \sqrt{\frac{765 - 180}{5}} \\ &= \sqrt{\frac{585}{5}} \\ &= \sqrt{117} \\ &= 10.82\end{aligned}$$

$$\begin{aligned}\therefore \text{C.V} &= \frac{\sigma}{x} \times 100 \\ &= \frac{10.82}{6} \times 100 \\ &= \frac{1082}{6} \\ &= 180.33\%\end{aligned}$$

5. Find the coefficient of variation of 24, 26, 33, 37, 29, 31.

Solution :

Given data is 24, 26, 33, 37, 29, 31.

$$\therefore \text{C.V} = \frac{\sigma}{x} \times 100$$

$$\bar{x} = \frac{24 + 26 + 33 + 37 + 29 + 31}{6}$$

$$= \frac{180}{6}$$

$$= 30$$

To find σ_1 arrange them in ascending order.

x	$d = x - 31$	d^2
24	-7	49
26	-5	25
29	-2	4
31	0	0
33	2	4
37	6	36
	-6	118

$$\begin{aligned}\therefore \sigma &= \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \\ &= \sqrt{\frac{118}{6} - \left(\frac{-6}{6}\right)^2} \\ &= \sqrt{\frac{118}{6} - 1} \\ &= \sqrt{\frac{112}{6}} \\ &= \sqrt{18.6} \\ \sigma &= 4.31\end{aligned}$$

$$\begin{aligned}\therefore \text{C.V} &= \frac{4.31}{30} \times 100 \\ &= \frac{43.1}{3} \\ &= 14.36 \\ &\approx 14.4\%\end{aligned}$$

6. The time taken (in minutes) to complete a homework by 8 students in a day are given by 38, 40, 47, 44, 46, 43, 49, 53. Find the coefficient of variation.

Solution :

Given data is 38, 40, 47, 44, 46, 43, 49, 53.

$$\begin{aligned}\bar{x} &= \frac{38 + 40 + 47 + 44 + 46 + 43 + 49 + 53}{8} \\ &= \frac{360}{8} \\ &= 45\end{aligned}$$

To find σ , arrange them in ascending order.

x	$d = x - 46$	d^2
38	-8	64
40	-6	36
43	-3	9
44	-2	4
46	0	0
47	1	1
49	3	9
53	7	49
	-8	172

$$\begin{aligned}\therefore \sigma &= \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \\ &= \sqrt{\frac{172}{8} - \left(\frac{-8}{8}\right)^2} \\ &= \sqrt{\frac{172}{8} - 1} = \sqrt{\frac{164}{8}} = \sqrt{20.5} = 4.53\end{aligned}$$

$$\begin{aligned}\therefore C.V &= \frac{4.53}{45} \times 100 \\ &= \frac{453}{45} \\ &= 10.07\%\end{aligned}$$

7. The total marks scored by two students Sathya and Vidhya in 5 subjects are 460 and 480 with standard deviation 4.6 and 2.4 respectively. Who is more consistent in performance?

Solution :

Sathya

$$\sum x_1 = 460$$

$$n = 5$$

$$\therefore \bar{x}_1 = \frac{460}{5}$$

$$= 92$$

$$\sigma_1 = 4.6$$

Vidhya

$$\sum x_2 = 480$$

$$n = 5$$

$$\therefore \bar{x}_2 = \frac{480}{5}$$

$$= 96$$

$$\sigma_2 = 2.4$$

$$\therefore C.V_1 = \frac{\sigma_1}{\bar{x}_1} \times 100$$

$$= \frac{4.6}{92}$$

$$= \frac{460}{92}$$

$$= 5$$

$$\therefore C.V_2 = \frac{\sigma_2}{\bar{x}_2} \times 100$$

$$= \frac{2.4}{96} \times 100$$

$$= \frac{240}{96}$$

$$= 2.5$$

$$\therefore C.V_2 < C.V_1$$

\therefore Vidhya is more consistent than Sathya.

8. The mean and standard deviation of marks obtained by 40 students of a class in three subjects Mathematics, Science and Social Science are given below.

Subject	Mean	SD
Mathematics	56	12
Science	65	14
Social Science	60	10

Which of the three subjects shows highest variation and which shows lowest variation in marks?

Solution :

$$C.V = \frac{\sigma}{x} \times 100$$

$$\text{For Maths, } C.V = \frac{12}{56} \times 100 = 21.428$$

$$\text{For Science, } C.V = \frac{14}{65} \times 100 = 21.538$$

$$\text{For Social Science, } C.V = \frac{10}{60} \times 100 = 16.67$$

Highest variation in Science.

Lowest variation in Social Science.

9. The temperature of two cities A and B in a winter season are given below.

Temperature of city A (in degree Celsius)	18	20	22	24	26
Temperature of city B (in degree Celsius)	11	14	15	17	18

Find which city is more consistent in temperature changes?

Solution :

Temperature of City 'A' :

18, 20, 22, 24, 26

$$\bar{x} = \frac{110}{5} = 22$$

x	$d = \frac{x-22}{2}$	d^2
18	-2	4
20	-1	1
22	0	0
24	1	1
26	2	4
	0	10

$$\begin{aligned} \therefore \sigma &= \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \times C \\ &= \sqrt{\frac{10}{5} - 0^2} \times 2 \\ &= 2\sqrt{2} \end{aligned}$$

$$\begin{aligned} \therefore \text{CV for city A} &= \frac{\sigma}{x} \times 100 \\ &= \frac{2\sqrt{2}}{22} \\ &= \frac{100 \times 1.414}{22} \\ &= 6.427 \end{aligned}$$

Temperature of City B 11, 14, 15, 17, 18

$$\bar{x} = \frac{75}{5} = 15$$

x	$d = x - 15$	d^2
11	-4	16
14	-1	1
15	0	0
17	2	4
18	3	9
	0	30

$$\begin{aligned} \therefore \sigma &= \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \times C \\ &= \sqrt{\frac{30}{5} - 0^2} \\ &= \sqrt{6} \end{aligned}$$

$$\begin{aligned}
 \therefore \text{CV for city B} &= \frac{\sigma}{x} \times 100 \\
 &= \frac{\sqrt{6}}{15} \times 100 \\
 &= \frac{2.45}{15} \times 100 \\
 &= 16.34
 \end{aligned}$$

\therefore CV for City A < CV for City B.

\therefore City A is more consistent in temperature changes.

III. PROBABILITY :

Key Points

- ✓ A **random experiment** is an experiment in which
 - (i) The set of all possible outcomes are known (ii) Exact outcome is not known.
 - ✓ The set of all possible outcomes in a random experiment is called a **sample space**. It is generally denoted by S.
 - ✓ In a random experiment, each possible outcome is called an **event**.
 - ✓ An event will be a subset of the sample space.
 - ✓ If an event E consists of only one outcome then it is called an **elementary event**.
 - ✓ $P(E) = \frac{n(E)}{n(S)}$
 - ✓ $P(S) = \frac{n(S)}{n(S)} = 1$. The probability of sure event is 1.
 - ✓ $P(\phi) = \frac{n(\phi)}{n(s)} = \frac{0}{n(s)} = 0$. The probability of impossible event is 0.
 - ✓ E is a subset of S and ϕ is a subset of any set.
- $$\phi \subseteq E \subseteq S \quad P(\phi) \leq P(E) \leq P(S) \quad 0 \leq P(E) \leq 1$$
- ✓ The complement event of E is \bar{E} .
 - ✓ $P(E) + P(\bar{E}) = 1$.


Example 8.18

Express the sample space for rolling two dice using tree diagram.

Solution :

When we roll two dice, since each die contain 6 faces marked with 1,2,3,4,5,6 the tree diagram will look like

Hence, the sample space can be written as



$$S = \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6) \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6) \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6) \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6) \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6) \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6)\}$$

Example 8.19

A bag contains 5 blue balls and 4 green balls. A ball is drawn at random from the bag. Find the probability that the ball drawn is (i) blue (ii) not blue.

Solution :

Total number of possible outcomes

$$n(S) = 5 + 4 = 9$$

- (i) Let A be the event of getting a blue ball.

Number of favourable outcomes for the event A. Therefore, $n(A) = 5$

Hence we get, $y = x + 11$ gives

$$x - y + 11 = 0.$$

Probability that the ball drawn is blue.

$$\text{Therefore, } P(A) = \frac{n(A)}{n(S)} = \frac{5}{9}$$

- (ii) \bar{A} will be the event of not getting a blue ball.

$$\text{So } P(\bar{A}) = 1 - P(A) = 1 - \frac{5}{9} = \frac{4}{9}$$

Example 8.20

Two dice are rolled. Find the probability that the sum of outcomes is (i) equal to 4 (ii) greater than

Solution :

When we roll two dice, the sample space is given by

$$S = \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6) \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6) \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6) \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6) \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6) \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6)\}; \\ n(S) = 36$$

- (i) Let A be the event of getting the sum of outcome values equal to 4.

$$\text{Then } A = \{(1,3), (2,2), (3,1)\}; n(A) = 3.$$

Probability of getting the sum of outcomes equal to 4 is

$$P(A) = \frac{n(A)}{n(S)} = \frac{3}{36} = \frac{1}{12}$$

- (ii) Let B be the event of getting the sum of outcome values greater than 10.

Then $B = \{(5,6), (6,5), (6,6)\}$; $n(B) = 3$

Probability of getting the sum of outcomes greater than 10 is

$$P(B) = \frac{n(B)}{n(S)} = \frac{3}{36} = \frac{1}{12}$$

(iii) Let C be the event of getting the sum of outcomes less than 13. Here all the outcomes have the sum value less than 13. Hence $C = S$.

Therefore, $n(C) = n(S) = 36$

Probability of getting the total value less than 13 is

$$P(C) = \frac{n(C)}{n(S)} = \frac{36}{36} = 1$$

Example 8.21

Two coins are tossed together. What is the prob-

Solution :

When two coins are tossed together, the sample space is

$$S = \{HH, HT, TH, TT\}; n(S) = 4$$

Let A be the event of getting different faces on the coins.

$$A = \{HT, TH\}; n(A) = 2$$

Probability of getting different faces on the coins is

$$P(A) = \frac{n(A)}{n(S)} = \frac{2}{4} = \frac{1}{2}$$

Example 8.22

From a well shuffled pack of 52 cards, one card is drawn at random. Find the probability of getting (i) red card (ii) heart card (iii) red king (iv) face card (v) number card

Solution :

$$n(S) = 52$$

(i) Let A be the event of getting a red card.

$$n(A) = 26$$

Probability of getting a red card is

$$P(A) = \frac{26}{52} = \frac{1}{2}$$

(ii) Let B be the event of getting a heart card.

$$n(B) = 13$$

Probability of getting a heart card is

$$P(B) = \frac{n(B)}{n(S)} = \frac{13}{52} = \frac{1}{4}$$

(iii) Let C be the event of getting a red king card. A red king card can be either a diamond king or a heart king.

$$n(C) = 2$$

Probability of getting a red king card is

$$P(C) = \frac{n(C)}{n(S)} = \frac{2}{52} = \frac{1}{26}$$

(iv) Let D be the event of getting a face card. The face cards are Jack (J), Queen (Q), and King (K).

$$n(D) = 4 \times 3 = 12$$

Probability of getting a face card is

$$P(D) = \frac{n(D)}{n(S)} = \frac{12}{52} = \frac{3}{13}$$

(v) Let E be the event of getting a number card. The number cards are 2, 3, 4, 5, 6, 7, 8, 9 and 10.

$$n(E) = 4 \times 9 = 36$$

Probability of getting a number card is

$$P(E) = \frac{n(E)}{n(S)} = \frac{36}{52} = \frac{9}{13}$$

Example 8.23

What is the probability that a leap year selected at random will contain 53 Saturdays.

(Hint: $366 = 52 \times 7 + 2$)

Solution :

A leap year has 366 days. So it has 52 full weeks and 2 days. 52 Saturdays must be in 52 full weeks.

The possible chances for the remaining two days will be the sample space.

$S = \{(\text{Sun-Mon, Mon-Tue, Tue-Wed, Wed-Thu, Thu-Fri, Fri-Sat, Sat-Sun})\}$

$$n(S) = 7$$

Let A be the event of getting 53rd Saturday.

Then $A = \{\text{Fri-Sat, Sat-Sun}\}$; $n(A) = 2$

year is

$$P(A) = \frac{n(A)}{n(S)} = \frac{2}{7}$$

Example 8.24

A die is rolled and a coin is tossed simultaneously. Find the probability that the die shows an odd number and the coin shows a head.

Solution :

Sample space

$S = \{1H, 1T, 2H, 2T, 3H, 3T, 4H, 4T, 5H, 5T, 6H, 6T\}$;

$n(S) = 12$

Let A be the event of getting an odd number and a head.

$A = \{1H, 3H, 5H\}$; $n(A) = 3$

$$P(A) = \frac{n(A)}{n(S)} = \frac{3}{12} = \frac{1}{4}$$

Example 8.25

A bag contains 6 green balls, some black and red balls. Number of black balls is as twice as the number of red balls. Probability of getting a green ball is thrice the probability of getting a red ball. Find (i) number of black balls (ii) total number of balls.

Solution :

Number of green balls is $n(G) = 6$

Let number of red balls is $n(R) = x$

Therefore, number of black balls is $n(B) = 2x$

Total number of balls $n(S) = 6 + x + 2x = 6 + 3x$

It is given that, $P(G) = 3 \times P(R)$

$$\frac{6}{6 + 3x} = 3 \times \frac{x}{6 + 3x}$$

$3x = 6$ gives, $x = 2$.

(i) Number of black balls $= 2 \times 2 = 4$

(ii) Total number of balls $= 6 + (3 \times 2) = 12$

Example 8.26

A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the numbers 1, 2, 3, ... 12. What is the probability that it will point to (i) 7 (ii) a prime number (iii) a composite number?

Solution :

Sample space

$S = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$; $n(S) = 12$

(i) Let A be the event of resting in 7. $n(A) = 1$

$$P(A) = \frac{n(A)}{n(S)} = \frac{1}{12}$$

(ii) Let B be the event that the arrow will come to rest in a prime number.

$B = \{2, 3, 5, 7, 11\}$; $n(B) = 5$

$$P(B) = \frac{n(B)}{n(S)} = \frac{5}{12}$$

- (iii) Let C be the event that arrow will come to rest in a composite number.


$$C = \{4, 6, 8, 9, 10, 12\}; n(C) = 6$$

$$P(C) = \frac{n(C)}{n(S)} = \frac{6}{12} = \frac{1}{2}$$

EXERCISE 8.3

1. Write the sample space for tossing three coins using tree diagram.

Solution :




Sample space = {(HHH), (HHT), (HTH), (HTT), (THH), (THT), (TTH), (TTT)}

2. Write the sample space for selecting two balls from a bag containing 6 balls numbered 1 to 6 (using tree diagram).

Solution :

$$S = \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6) \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6) \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6) \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6) \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6) \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6)\}$$



3. If A is an event of a random experiment such that $P(A) : P(\bar{A}) = 17:15$ and $n(S) = 640$ then find (i) $P(\bar{A})$ (ii) $n(A)$.

Solution :

$$\text{Given } P(A) : P(\bar{A}) = 17 : 15$$

$$\Rightarrow \frac{1 - P(\bar{A})}{P(A)} = \frac{17}{15}$$

$$\Rightarrow 15 - 15P(A) = 17P(\bar{A})$$

$$\Rightarrow 32P(\bar{A}) = 15$$

$$\Rightarrow P(A) = \frac{17}{32}$$

$$\Rightarrow \frac{n(A)}{n(S)} = \frac{17}{32}$$

$$\Rightarrow n(A) = \frac{17}{32} \times 640 = 340$$

4. A coin is tossed thrice. What is the probability of getting two consecutive tails?

Solution :

When a coin is tossed thrice,

$S = \{(HHH), (HHT), (HTH), (HTT), (THH), (THT), (TTH), (TTT)\}$

$$n(S) = 8$$

Let A be the event of getting 2 tails continuously,

$A = \{(HTT), (TTH), (TTT)\}$

$$n(A) = 3$$

$$P(A) = \frac{n(A)}{n(S)} = \frac{3}{8}$$

5. At a fete, cards bearing numbers 1 to 1000, one number on one card are put in a box. Each player selects one card at random and that card is not replaced. If the selected card has a perfect square number greater than 500, the player wins a prize. What is the probability that (i)

and player wins a prize, if the first has won?

Solution :

$$n(S) = 1000$$

- i) Let A be the event of getting perfect squares between 500 and 1000

$$A = \{23^2, 24^2, 25^2, 26^2, \dots, 31^2\}$$

$$n(A) = 9$$

$$P(A) = \frac{9}{1000}$$

is the probability for the 1st player to win a prize.

- ii) When the card which was taken first is not replaced.

$$n(S) = 999$$

$$n(B) = 8$$

$$P(B) = \frac{8}{999}$$

6. A bag contains 12 blue balls and x red balls. If one ball is drawn at random (i) what is the probability that it will be a red ball? (ii) If 8 more red balls are put in the bag, and if the probability of drawing a red ball will be twice that of the probability in (i), then find x .

Solution :

Total number of balls in the bag

$$= x + 12. (x \rightarrow \text{red } 12 \rightarrow \text{black})$$

- i) Let A be the event of getting red balls

$$P(A) = \frac{n(A)}{n(S)} = \frac{x}{x+12}$$

- ii) If 8 more red balls are added in the bag.

$$n(S) = x + 20$$

$$\text{By the problem, } \frac{x+8}{x+20} = 2 \left(\frac{x}{x+12} \right)$$

$$\Rightarrow (x+8)(x+12) = 2x^2 + 40x$$

$$\Rightarrow x^2 + 20x + 96 = 2x^2 + 40x$$

$$\Rightarrow x^2 + 20x - 96 = 0$$

$$\Rightarrow (x+24)(x-4) = 0$$

$$\therefore x = -24, 4$$

$$\therefore x = 4$$

$$\therefore P(A) = \frac{4}{16} = \frac{1}{4}$$

7. Two unbiased dice are rolled once. Find the probability of getting

- (i) a doublet (equal numbers on both dice)
- (ii) the product as a prime number
- (iii) the sum as a prime number
- (iv) the sum as 1

Solution :

$$S = \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6)$$

$$(2,1), (2,2), (2,3), (2,4), (2,5), (2,6)$$

$$(3,1), (3,2), (3,3), (3,4), (3,5), (3,6)$$

(4,1),(4,2),(4,3),(4,4),(4,5),(4,6)
 (5,1),(5,2),(5,3),(5,4),(5,5),(5,6)
 (6,1),(6,2),(6,3),(6,4),(6,5),(6,6)}

i) Let A be the event of getting a doublet

$$A = \{(1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6)\}$$

$$n(A) = 6$$

$$\therefore P(A) = \frac{n(A)}{n(S)} = \frac{6}{36} = \frac{1}{6}$$

ii) Let B be the event of getting the product as a prime number.

$$B = \{(1, 2), (1, 3), (1, 5), (2, 1), (3, 1), (5, 1)\}$$

$$n(B) = 6$$

$$\therefore P(B) = \frac{n(B)}{n(S)} = \frac{6}{36} = \frac{1}{6}$$

iii) Let C be the event of getting the sum of numbers on the dice is prime.

$$C = \{(1, 1), (1, 2), (1, 4), (2, 1), (2, 3), (2, 5), (3, 2), (3, 4), (4, 1), (4, 3), (5, 2), (5, 6), (6, 1), (6, 5)\}$$

$$n(C) = 14$$

$$\therefore P(C) = \frac{n(C)}{n(S)} = \frac{7}{36}$$

iv) Let D be the event of getting sum of numbers is 1.

$$n(D) = 0$$

$$P(D) = 0$$

8. Three fair coins are tossed together. Find the probability of getting

(i) all heads (ii) atleast one tail

(iii) atleast one head (iv) atleast two tails

Solution :

When 3 fair coins are tossed,

$$S = \{(HHH), (HHT), (HTH), (HTT), (THH), (THT), (TTH), (TTT)\}$$

$$n(S) = 8$$

i) Let A be the event of getting all heads.

$$A = \{(HHH)\}$$

$$n(A) = 1$$

$$\therefore P(A) = \frac{n(A)}{n(S)} = \frac{1}{8}$$

ii) Let B be the event of getting atleast one tail.

$$B = \{(HHT), (HTH), (HTT), (THH), (THT), (TTH), (TTT)\}$$

$$n(B) = 7$$

$$P(B) = \frac{7}{8}$$

head.

$$C = \{(HTT), (THT), (TTH), (TTT)\}$$

$$n(C) = 4$$

$$P(C) = \frac{4}{8} = \frac{1}{2}$$

iv) Let D - atleast 2 tails

$$D = \{(HHH), (HHT), (HTT), (HTH), (THH), (THT), (TTH)\}$$

$$n(D) = 7$$

$$P(D) = \frac{7}{8}$$

9. Two dice are numbered 1,2,3,4,5,6 and 1,1,2,2,3,3 respectively. They are rolled and the sum of the numbers on them is noted. Find the probability of getting each sum from 2 to 9 separately.

Solution :

$$S = \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6), (2,1), (2,2), (2,3), (2,4), (2,5), (2,6)\}$$

(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)
 (4,1),(4,2),(4,3),(4,4),(4,5),(4,6)
 (5,1),(5,2),(5,3),(5,4),(5,5),(5,6)
 (6,1),(6,2),(6,3),(6,4),(6,5),(6,6)}

i) Let A - Sum of 2

$$n(A) = 2$$

$$\therefore P(A) = \frac{2}{36}$$

ii) Let B - Sum of 3

$$n(B) = 4$$

$$P(B) = \frac{4}{36}$$

iii) Let C - Sum of 4

$$n(C) = 6$$

$$P(C) = \frac{6}{36}$$

iv) Let D - Sum of 5

$$n(D) = 6$$

$$P(D) = \frac{6}{36}$$

v) Let E - Sum of 6

$$n(E) = 6$$

$$P(E) = \frac{6}{36}$$

vi) Let F - Sum of 7

$$n(F) = 6$$

$$P(F) = \frac{6}{36}$$

vii) Let G - Sum of 8

$$n(G) = 4$$

$$P(G) = \frac{4}{36}$$

viii) Let H - Sum of 9

$$n(H) = 2$$

$$P(H) = \frac{2}{36}$$

$$P(H) = \frac{2}{36}$$

10. A bag contains 5 red balls, 6 white balls, 7 green balls, 8 black balls. One ball is drawn at random from the bag. Find the probability that the ball drawn is (i) white (ii) black or red (iii) not white (iv) neither white nor black

Solution :

i) Let A - White ball

$$n(A) = 6$$

$$P(A) = \frac{6}{26} = \frac{3}{13}$$

ii) Let B - Black (or) red

$$n(B) = 5 + 8 = 13$$

$$P(B) = \frac{13}{26} = \frac{1}{2}$$

iii) Let C - not white

$$n(C) = 20$$

$$P(C) = \frac{20}{26} = \frac{10}{13}$$

iv) Let D - Neither white nor black

$$n(D) = 12$$

$$P(D) = \frac{12}{26} = \frac{6}{13}$$

11. In a box there are 20 non-defective and some defective bulbs. If the probability that a bulb selected at random from the box found to be defective is $\frac{3}{8}$ then, find the number of defective bulbs.

Solution :

Let x be the number of defective bulbs.

$$\therefore n(S) = x + 20$$

Let A be the event of selecting defective balls

$$\therefore n(A) = x$$

$$P(A) = \frac{x}{x + 20}$$

$$\text{Given } \frac{x}{x + 20} = \frac{3}{8}$$

$$\Rightarrow 8x = 3x + 60$$

$$\Rightarrow 5x = 60$$

$$x = 12$$

\therefore Number of defective balls = 12.

12. The king and queen of diamonds, queen and jack of hearts, jack and king of spades are removed from a deck of 52 playing cards and then well shuffled. Now one card is drawn at random from the remaining cards. Determine the probability that the card is (i) a clavor (ii) a queen of red card (iii) a king of black card

Solution :

Solution :

By the data given,

$$n(S) = 52 - 2 - 2 - 2 = 46$$

- i) Let A be the event of selecting clubber card.
- $$n(A) = 13$$

$$P(A) = \frac{13}{46}$$

- ii) Let B - queen of red card.

$$n(B) = 0$$

$$P(B) = 0$$


(queen diamond and heart are included in S)

- iii) Let C - King of black cards

$$n(C) = 1 \text{ (enclusing spade king)}$$

$$\therefore P(C) = \frac{1}{46}$$

13. Some boys are playing a game, in which the stone thrown by them landing in a circular region (given in the figure) is considered as win and landing other than the circular region is considered as loss. What is the probability to win the



Solution :

$$\begin{aligned} \text{Area of the rectangular region} &= 4 \times 3 \\ &= 12\text{ft}^2 \end{aligned}$$

$$\begin{aligned} \text{Area of the circular region} &= \pi r^2 \\ &= \pi \times 1^2 \\ &= \pi\text{ft}^2 \end{aligned}$$

$$\begin{aligned} \therefore \text{Probability to win the game} &= \frac{\pi}{12} \\ &= \frac{3.14}{12} \\ &= \frac{314}{1200} \\ &= \frac{157}{600} \end{aligned}$$

14. Two customers Priya and Amuthan are visiting a particular shop in the same week (Monday to Saturday). Each is equally likely to visit the shop on any one day as on another day. What is the probability that both will visit the shop on
(i) the same day (ii) different days
(iii) consecutive days?

Solution :

Given $n(S) = 6$. (Monday - Saturday)

- i) Prob. that both of them will visit the shop on the same day = $\frac{1}{6}$
 ii) Prob. that both of them will visit the shop in different days = $\frac{5}{6}$.
 (\because if one visits on Monday, other one visit the shop out of remaining 5 days).

consecutive days.

$A = \{(\text{Mon, Tue}), (\text{Tue, Wed}), (\text{Wed, Thu}), (\text{Thu, Fri}), (\text{Fri, Sat})\}$

$n(A) = 5$

$$P(A) = \frac{5}{6}$$

15. In a game, the entry fee is ₹150. The game consists of tossing a coin 3 times. Dhana bought a ticket for entry. If one or two heads show, she gets her entry fee back. If she throws 3 heads, she receives double the entry fees. Otherwise she will lose. Find the probability that she (i) gets double entry fee (ii) just gets her entry fee (iii) loses the entry fee.

Solution :

$S = \{(\text{HHH}), (\text{HHT}), (\text{HTH}), (\text{THH}), (\text{HTT}), (\text{THT}), (\text{TTH}), (\text{TTT})\}$

$n(S) = 8$

- i) $P(\text{gets double entry fee}) = \frac{1}{8}$ (\because 3 heads)
 ii) $P(\text{just gets for her entry fee}) = \frac{6}{8} = \frac{3}{4}$
 (\because 1 (or) 2 heads)
 iii) $P(\text{loses the entry fee}) = \frac{1}{8}$
 (\because 3 no heads (TTT) only)

IV. ALGEBRA OF EVENTS:


Key Points

- ✓ $A \cap \bar{A} = \phi$ $A \cup \bar{A} = S$
- ✓ If A, B are mutually exclusive events, the $P(A \cup B) = P(A) + P(B)$.
- ✓ $P(\text{Union of mutually exclusive events}) = \sum (\text{Probability of events})$

Theorem 1

If A and B are two events associated with a random experiment, then prove that

- (i) $P(A \cap \bar{B}) = P(\text{only A}) = P(A) - P(A \cap B)$
 (ii) $P(\bar{A} \cap B) = P(\text{only B}) = P(B) - P(A \cap B)$

Proof

(i) By Distributive property of sets,

$$1. (A \cap B) \cup (A \cap \bar{B}) = A \cap (B \cup \bar{B}) = A \cap S = A$$

$$2. (A \cap B) \cap (A \cap \bar{B}) = A \cap (B \cap \bar{B}) = A \cap \phi = \phi$$

Therefore, $P(A) = P[(A \cap B) \cup (A \cap \bar{B})]$

$$P(A) = P(A \cap B) + P(A \cap \bar{B})$$

Therefore, $P(A \cap \bar{B}) = P(A) - P(A \cap B)$

(ii) By Distributive property of sets,

$$1. (A \cap B) \cup (\bar{A} \cap B) = (A \cup \bar{A}) \cap B = S \cap B = B$$

$$2. (A \cap B) \cap (\bar{A} \cap B) = (A \cap \bar{A}) \cap B = \phi \cap B = \phi$$

Therefore, the events $A \cap B$ and $\bar{A} \cap B$ are mutually exclusive whose union is B.

$$P(B) = P[(A \cap B) \cup (\bar{A} \cap B)]$$

$$P(B) = P(A \cap B) + P(\bar{A} \cap B)$$

Therefore, $P(\bar{A} \cap B) = P(B) - P(A \cap B)$

That is, $P(\bar{A} \cap B) = P(\text{only } B) = P(B) - P(A \cap B)$

Theorem 2

(i) If A and B are any two events then

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

(ii) If A, B and C are any three events then

$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$

$$- P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$$

Proof

(i) Let A and B be any two events of a random experiment with sample space S.

From the Venn diagram, we have the events only A, $A \cap B$ and only B are mutually exclusive and their union is $A \cup B$

Therefore,


$$P(A \cup B) = P[(\text{only } A) \cup (A \cap B) \cup (\text{only } B)]$$

$$= P(\text{only } A) + P(A \cap B) + P(\text{only } B)$$

$$= [P(A) - P(A \cap B)] + P(A \cap B) + [P(B) - P(A \cap B)]$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

(ii) Let A, B, C are any three events of a random experiment with sample space S.



Let $D = B \cup C$

$$P(A \cup B \cup C) = P(A \cup D)$$

$$= P(A) + P(D) - P(A \cap D)$$

$$= P(A) + P(B \cup C) - P[A \cap (B \cup C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) -$$

$$P[(A \cap B) \cup (A \cap C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) - P[(A \cap B)$$

$$- P(A \cap C) + P[(A \cap B) \cap (A \cap C)]$$

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) -$$

$$P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

Example 8.27

If $P(A) = 0.37$, $P(B) = 0.42$, $P(A \cap B) = 0.09$ then find $P(A \cup B)$.

Solution :

$$P(A) = 0.37, P(B) = 0.42, P(A \cap B) = 0.09$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cup B) = 0.37 + 0.42 - 0.09 = 0.7$$

Example 8.28

What is the probability of drawing either a king or a queen in a single draw from a well shuffled pack of 52 cards?

Solution :

Total number of cards = 52

Number of king cards = 4

$$\text{Probability of drawing a king card} = \frac{4}{52}$$

Number of queen cards = 4

$$\text{Probability of drawing a queen card} = \frac{4}{52}$$

Both the events of drawing a king and a queen are mutually exclusive

$$\Rightarrow P(A \cup B) = P(A) + P(B)$$

Therefore, probability of drawing either a king or a queen = $\frac{4}{52} + \frac{4}{52} = \frac{2}{13}$

Example 8.29

Two dice are rolled together. Find the probability of getting a doublet or sum of faces as 4.

Solution :

When two dice are rolled together, there will be $6 \times 6 = 36$ outcomes. Let S be the sample space. Then $n(S) = 36$

Let A be the event of getting a doublet and B be the event of getting face sum 4.

$$\text{Then } A = \{(1,1), (2,2), (3,3), (4,4), (5,5), (6,6)\}$$

$$B = \{(1,3), (2,2), (3,1)\}$$

Therefore, $A \cap B = \{(2,2)\}$

Then, $n(A) = 6$, $n(B) = 3$, $n(A \cap B) = 1$.

$$P(A) = \frac{n(A)}{n(S)} = \frac{6}{36}$$

$$P(B) = \frac{n(B)}{n(S)} = \frac{3}{36}$$

$$P(A \cap B) = \frac{n(A \cap B)}{n(S)} = \frac{1}{36}$$

Therefore,

$P(\text{getting a doublet or a total of 4}) = P(A \cup B)$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{6}{36} + \frac{3}{36} - \frac{1}{36} = \frac{8}{36} = \frac{2}{9}$$

Hence, the required probability is $\frac{2}{9}$

If A and B are two events such that $P(A) = \frac{1}{4}$,

$P(B) = \frac{1}{2}$ and $P(A \text{ and } B) = \frac{1}{8}$, find

(i) $P(A \text{ or } B)$ (ii) $P(\text{not } A \text{ and not } B)$.

Solution :

(i) $P(A \text{ or } B) = P(A \cup B)$

$$= P(A) + P(B) - P(A \cap B)$$

$$P(A \text{ or } B) = \frac{1}{4} + \frac{1}{2} - \frac{1}{8} = \frac{5}{8}$$

(ii) $P(\text{not } A \text{ and not } B) = P(\overline{A \cap B})$

$$= P(\overline{A \cup B})$$

$$= 1 - P(A \cup B)$$

$$P(\text{not } A \text{ and not } B) = 1 - \frac{5}{8} = \frac{3}{8}$$

Example 8.31

A card is drawn from a pack of 52 cards. Find the probability of getting a king or a heart or a red card.

Solution :

Total number of cards = 52; $n(S) = 52$

Let A be the event of getting a king card.

$$n(A) = 4$$

$$P(A) = \frac{n(A)}{n(S)} = \frac{4}{52}$$

Let B be the event of getting a heart card.

$$n(B) = 13$$

$$P(B) = \frac{n(B)}{n(S)} = \frac{13}{52}$$

$$n(C) = 26$$

$$P(C) = \frac{n(C)}{n(S)} = \frac{26}{52}$$

$$P(A \cap B) = P(\text{getting heart king}) = \frac{1}{52}$$

$$P(B \cap C) = P(\text{getting red and heart}) = \frac{13}{52}$$

$$P(A \cap C) = P(\text{getting red king}) = \frac{2}{52}$$

$$P(A \cap B \cap C) = P(\text{getting heart, king which is red})$$

$$= \frac{1}{52}$$

Therefore, required probability is

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

$$= \frac{4}{52} + \frac{13}{52} + \frac{26}{52} - \frac{1}{52} - \frac{13}{52} - \frac{2}{52} + \frac{1}{52} = \frac{28}{52} = \frac{7}{13}$$

Example 8.32

In a class of 50 students, 28 opted for NCC, 30 opted for NSS and 18 opted both NCC and NSS. One of the students is selected at random. Find the probability that

- The student opted for NCC but not NSS.
- The student opted for NSS but not NCC.
- The student opted for exactly one of them.

Solution:

Total number of students $n(S) = 50$.

Let A and B be the events of students opted for NCC and NSS respectively.

$$n(A) = 28, n(B) = 30, n(A \cap B) = 18$$

$$P(A) = \frac{n(A)}{n(S)} = \frac{28}{50}$$

$$P(B) = \frac{n(B)}{n(S)} = \frac{30}{50}$$

$$P(A \cap B) = \frac{n(A \cap B)}{n(S)} = \frac{18}{50}$$

- Probability of the students opted for NCC but not NSS

$$P(A \cap \bar{B}) = P(A) - P(A \cap B) = \frac{28}{50} - \frac{18}{50} = \frac{10}{50} = \frac{1}{5}$$

- Probability of the students opted for NSS but not NCC.

$$P(A \cap \bar{B}) = P(B) - P(A \cap B) = \frac{30}{50} - \frac{18}{50} = \frac{12}{50} = \frac{6}{25}$$

- Probability of the students opted for exactly one of them

$$= P[(A \cap \bar{B}) \cup (\bar{A} \cap B)]$$

$$= P(A \cap \bar{B}) + P(\bar{A} \cap B) = \frac{1}{5} + \frac{6}{25} = \frac{11}{25}$$

(Note that $(A \cap \bar{B})$, $(\bar{A} \cap B)$ are mutually exclusive events)

Example 8.33

A and B are two candidates seeking admission to IIT. The probability that A getting selected is 0.5 and the probability that both A and B getting selected is 0.3. Prove that the probability of B being selected is at most 0.8.

Solution:

$$P(A) = 0.5, P(A \cap B) = 0.3$$

We have $P(A \cup B) \leq 1$

$$P(A) + P(B) - P(A \cap B) \leq 1$$

$$0.5 + P(B) - 0.3 \leq 1$$

$$P(B) \leq 1 - 0.2$$

$$P(B) \leq 0.8$$

Therefore, probability of B getting selected is at most 0.8.

EXERCISE 8.4

1. If $P(A) = \frac{2}{3}$, $P(B) = \frac{2}{5}$, $P(A \cup B) = \frac{1}{3}$ then find $P(A \cap B)$.

Solution :

$$P(A \cap B) = P(A) + P(B) - P(A \cup B)$$

$$= \frac{2}{3} + \frac{2}{5} - \frac{1}{3}$$

$$= \frac{10 + 6 - 5}{15}$$

$$= \frac{11}{15}$$

2. A and B are two events such that, $P(A) = 0.42$, $P(B) = 0.48$, and $P(A \cap B) = 0.16$. Find (i) $P(\text{not } A)$ (ii) $P(\text{not } B)$ (iii) $P(A \text{ or } B)$

Solution :

$$\begin{aligned} \text{a) } P(\text{not } A) &= P(\bar{A}) = 1 - P(A) \\ &= 1 - 0.42 \end{aligned}$$

$$= 0.58$$

$$\begin{aligned} \text{b) } P(\text{not } B) &= P(\bar{B}) = 1 - P(B) \\ &= 1 - 0.48 \end{aligned}$$

$$= 0.52$$

$$\begin{aligned} \text{c) } P(A \text{ or } B) &= P(A \cup B) \\ &= P(A) + P(B) - P(A \cap B) \\ &= 0.42 + 0.48 - 0.16 \\ &= 0.74 \end{aligned}$$

3. If A and B are two mutually exclusive events of a random experiment and $P(\text{not } A) = 0.45$, $P(A \cup B) = 0.65$, then find

Solution :

Given A and B are mutually exclusive events

$$P(A \cap B) = 0$$

$$\text{Also, } P(\text{not } A) = 0.45$$

$$\therefore P(\bar{A}) = 0.45$$

$$1 - P(A) = 0.45$$

$$P(A) = 0.55$$

$$P(A \cup B) = P(A) + P(B)$$

$$\therefore P(B) = P(A \cup B) - P(A)$$

$$= 0.65 - 0.55$$

$$= 0.10$$

4. The probability that atleast one of A and B occur is 0.6. If A and B occur simultaneously with probability 0.2, then find $P(\bar{A}) + P(\bar{B})$.

Solution :

$$\text{Given } P(A \cup B) = 0.6, P(A \cap B) = 0.2$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\Rightarrow 0.6 = P(A) + P(B) - 0.2$$

$$\therefore P(A) + P(B) = 0.8$$

$$\therefore P(\bar{A}) + P(\bar{B})$$

$$= 1 - P(A) + 1 - P(B)$$

$$= 2 - (P(A) + P(B))$$

$$= 2 - 0.8$$

$$= 1.2$$

5. The probability of happening of an event A is 0.5 and that of B is 0.3. If A and B are mutually exclusive events, then find the probability that neither A nor B happen.

Solution :

$$\text{Given } P(A) = 0.5, P(B) = 0.3, P(A \cap B) = 0$$

$$P(\text{neither A nor B})$$

$$= P(\bar{A} \cap \bar{B})$$

$$= P(\overline{A \cup B})$$

$$= 1 - P(A \cup B)$$

$$= 1 - [P(A) + P(B) - P(A \cap B)]$$

$$= 1 - (0.8)$$

$$= 0.2$$

6. Two dice are rolled once. Find the probability of getting an even number on the first die or a total of face sum 8.

Solution :

$$\begin{aligned} S = & \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6) \\ & (2,1), (2,2), (2,3), (2,4), (2,5), (2,6) \\ & (3,1), (3,2), (3,3), (3,4), (3,5), (3,6) \\ & (4,1), (4,2), (4,3), (4,4), (4,5), (4,6) \\ & (5,1), (5,2), (5,3), (5,4), (5,5), (5,6) \\ & (6,1), (6,2), (6,3), (6,4), (6,5), (6,6)\} \end{aligned}$$

$$n(S) = 36$$

Let A be the event of getting even number on the 1st die.

$$\begin{aligned} A = & \{(2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6) \\ & (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6) \\ & (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6)\} \end{aligned}$$

$$n(A) = 18$$

$$P(A) = \frac{18}{36}$$

Let B - Total of face sum as 8.

$$B = \{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}$$

$$n(B) = 5, P(B) = \frac{5}{36}$$

$$A \cap B = \{(2, 6), (4, 4), (6, 2)\}$$

$$n(A \cap B) = 3$$

$$\frac{3}{36}$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{18}{36} + \frac{5}{36} - \frac{3}{36}$$

$$= \frac{20}{36}$$

$$= \frac{5}{9}$$

7. From a well-shuffled pack of 52 cards, a card is drawn at random. Find the probability of it being either a red king or a black queen.

Solution :

$$n(S) = 52$$

Let A - Red King

$$n(A) = 2$$

$$P(A) = \frac{2}{52}$$

Let B - Black Queen

$$n(B) = 2$$

$$P(B) = \frac{2}{52}$$

Here A and B are mutually exclusive

$$\therefore P(A \cup B) = P(A) + P(B)$$

$$= \frac{4}{52}$$

$$= \frac{1}{13}$$

8. A box contains cards numbered 3, 5, 7, 9, ... 35, 37. A card is drawn at random from the box. Find the probability that the drawn card have either multiples of 7 or a prime number.

Solution :

$$S = \{3, 5, 7, 9, \dots, 35, 37\}$$

Let A - multiple of 7.

$$A = \{7, 14, 21, 28, 35\}$$

$$n(A) = 5$$

$$P(A) = \frac{5}{18}$$

Let B - a prime number

$$B = \{3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37\}$$

$$n(B) = 11$$

$$P(B) = \frac{11}{18}$$

$$\text{Here } A \cap B = \{7\}$$

$$n(A \cap B) = 1$$

$$P(A \cap B) = \frac{1}{18}$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\begin{aligned} &= \frac{5}{18} + \frac{11}{18} - \frac{1}{18} \\ &= \frac{15}{18} \\ &= \frac{5}{6} \end{aligned}$$

9. Three unbiased coins are tossed once. Find the probability of getting atmost 2 tails or atleast 2 heads.

Solution :

$$S = \{(HHH), (HHT), (HTH), (THH), (HTT), (THT), (TTH), (TTT)\}$$

$$n(S) = 8$$

Let A - at most 2 tails

$$A = \{(HHT), (HTH), (THH), (HTT), (THT), (TTH), (HHH)\}$$

$$n(A) = 7$$

$$P(A) = \frac{7}{8}$$

Let B - atleast 2 heads

$$B = \{(HHH), (HHT), (HTH), (THH)\}$$

$$n(B) = 4$$

$$P(B) = \frac{4}{8}$$

$$\therefore A \cap B = \{(HHH), (HHT), (HTH), (THH)\}$$

$$n(A \cap B) = 4, P(A \cap B) = \frac{4}{8}$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\begin{aligned} &= \frac{7}{8} + \frac{4}{8} - \frac{4}{8} \\ &= \frac{7}{8} \end{aligned}$$

10. The probability that a person will get an electrification contract is $\frac{3}{5}$ and the probability that he will not get plumbing contract is $\frac{5}{8}$. The probability of getting atleast one contract is $\frac{5}{7}$. What is the probability that he will get both?

Solution :

Let A - electrification contract

B - not plumbing contract

Given

$$P(A) = \frac{3}{5}, P(\bar{B}) = \frac{5}{8}, P(A \cup B) = \frac{5}{7}$$

$$\Rightarrow P(B) = 1 - \frac{5}{8}$$

$$\frac{3}{8}$$

$$\therefore P(A \cap B) = P(A) + P(B) - P(A \cup B)$$

$$= \frac{3}{5} + \frac{3}{8} - \frac{5}{7}$$

$$= \frac{168 + 105 - 200}{280}$$

$$= \frac{73}{280}$$

11. In a town of 8000 people, 1300 are over 50 years and 3000 are females. It is known that 30% of the females are over 50 years. What is the probability that a chosen individual from the town is either a female or over 50 years?

Solution :

Let A - Female

B - Over 50 years

Given $n(S) = 8000$, $n(A) = 3000$,

$$n(B) = 1300 \text{ and } n(A \cap B) = \frac{30}{100} \times 3000 = 900$$

$$\therefore P(A) = \frac{3000}{8000}, P(B) = \frac{1300}{8000}, P(A \cap B) = \frac{900}{8000}$$

$$\therefore P(\text{either a female (or) over 50 years})$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{3000 + 1300 - 900}{8000}$$

$$= \frac{3400}{8000}$$

$$= \frac{34}{80}$$

$$= \frac{17}{40}$$

12. A coin is tossed thrice. Find the probability of getting exactly two heads or atleast one tail or two consecutive heads.

Solution :

$$S = \{(HHH), (HHT), (HTH), (THH), (TTH), (THT), (HTT), (TTT)\}$$

$$n(S) = 8$$

Let A - exactly 2 heads

$$A = \{(HHT), (HTH), (THH)\}$$

$$n(A) = 3$$

$$P(A) = \frac{3}{8}$$

Let B - atleast one tail

$$B = \{(HHT), (HTH), (THH), (TTH), (THT), (HTT), (TTT)\}$$

$$n(B) = 7$$

$$P(B) = \frac{7}{8}$$

Let C - Consecutively 2 heads

$$C = \{(HHH), (HHT), (THH)\}$$

$$n(C) = 3$$

$$P(C) = \frac{3}{8}$$

$$A \cap B = \{(HHT), (HTH), (THH)\}$$

$$n(A \cap B) = 3$$

$$P(A \cap B) = \frac{3}{8}$$

$$B \cap C = \{(HHT), (THH)\}$$

$$n(B \cap C) = 2$$

$$P(B \cap C) = \frac{2}{8}$$

$$C \cap A = \{(HHT), (THH)\}$$

$$n(C \cap A) = 2$$

$$P(C \cap A) = \frac{2}{8}$$

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

$$= \frac{3}{8} + \frac{7}{8} + \frac{3}{8} - \frac{3}{8} - \frac{2}{8} - \frac{2}{8} + \frac{2}{8}$$

$$= \frac{8}{8} = 1$$

13. If A, B, C are any three events such that probability of B is twice as that of probability of A and probability of C is thrice as that of probability of A and if

$$P(A \cap B) = \frac{1}{6}, P(B \cap C) = \frac{1}{4}, P(A \cap C) = \frac{1}{8},$$

$$P(A \cup B \cup C) = \frac{9}{10}, P(A \cap B \cap C) = \frac{1}{15}, \text{ then}$$

find $P(A)$, $P(B)$ and $P(C)$?

Solution :

$$\text{Given } P(B) = 2 \cdot P(A), P(C) = 3 \cdot P(A)$$

$$P(A \cap B) = \frac{1}{6}, P(B \cap C) = \frac{1}{4}, P(A \cap C) = \frac{1}{8},$$

$$P(A \cup B \cup C) = \frac{9}{10}, P(A \cap B \cap C) = \frac{1}{15}$$

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

$$\Rightarrow \frac{9}{10} = P(A) + 2 \cdot P(A) + 3 \cdot P(A) - \frac{1}{6} - \frac{1}{4} - \frac{1}{8} + \frac{1}{15}$$

$$\Rightarrow 6 \cdot P(A) = \frac{9}{10} + \frac{1}{6} + \frac{1}{4} + \frac{1}{8} - \frac{1}{15}$$

$$\Rightarrow 6 \cdot P(A) = \frac{108 + 20 + 30 + 15 - 8}{120}$$

$$\Rightarrow 6 \cdot P(A) = \frac{165}{120}$$

$$\Rightarrow P(A) = \frac{165}{720} = \frac{11}{48}$$

$$\therefore P(A) = \frac{11}{48}$$

$$\therefore P(B) = 2 \cdot P(A) = 2 \times \frac{11}{48} = \frac{11}{24}$$

$$P(C) = 3 \cdot P(A) = 3 \times \frac{11}{48} = \frac{11}{16}$$

from 1 to 35. The ratio of boys to girls is 4:3. The roll numbers of students begin with boys and end with girls. Find the probability that a student selected is either a boy with prime roll number or a girl with composite roll number or an even roll number.

Solution :

Given $n(S) = 35$ and ratio of boys and girls = 4:3

$$\text{No. of boys} = \frac{4}{7} \times 35 = 20$$

$$\text{No. of girls} = \frac{3}{7} \times 35 = 15$$

Let A - a boy with prime roll no

$A = \{2, 3, 5, 7, 11, 13, 19\}$ (\because only 20 boys)

$$n(A) = 7$$

$$P(A) = \frac{7}{35}$$

Let B - a girl with composite roll no.

$B = \{21, 22, 24, 25, 26, 27, 28, 30, 32, 33, 34, 35\}$

$n(B) = 12$

$$\therefore P(B) = \frac{12}{25}$$

Let C - even roll no.

$B = \{2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34\}$

$n(C) = 17$

$$\therefore P(C) = \frac{17}{35}$$

$A \cap B = \{ \}, n(A \cap B) = 0, P(A \cap B) = 0$

$B \cap C = \{22, 24, 26, 28, 30, 32, 34\}$

$$\therefore n(B \cap C) = 7 \Rightarrow P(B \cap C) = \frac{7}{35}$$

$C \cap A = \{2\} \Rightarrow n(C \cap A) = 1$

$$\frac{1}{35}$$

at $P(A \cap B \cap C) = 0$

$$\therefore P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

$$= \frac{7}{35} + \frac{12}{35} + \frac{17}{35} - 0 - \frac{7}{35} - \frac{1}{35} + 0$$

$$= \frac{28}{35}$$

$$= \frac{4}{5}$$

EXERCISE 8.5

Multiple choice questions :

1. Which of the following is not a measure of dispersion?

(1) Range (2) Standard deviation
(3) Arithmetic mean (4) Variance

Ans : (3)

Hint :

A.M is not a measure of dispersion and it is a measure of central tendency.

2. The range of the data 8, 8, 8, 8, 8 . . . 8 is

(1) 0 (2) 1 (3) 8 (4) 3

Ans : (1)

Hint :

$$\begin{aligned} \text{Range} &= L - S \\ &= 8 - 8 = 0 \end{aligned}$$

3. The sum of all deviations of the data from its mean is

(1) Always positive
(2) always negative
(3) zero
(4) non-zero integer

Ans : (3)

Hint :

Sum of all deviations of the data from the mean = 0

$$\text{ie } \sum (x - \bar{x}) = 0$$

4. The mean of 100 observations is 40 and their standard deviation is 3. The sum of squares of all deviations is

(1) 40000 (2) 160900
(3) 160000 (4) 30000

Ans : (2)

Hint :

$$\bar{x} = 40, n = 100, \sigma = 3$$

$$\sigma^2 = \frac{\sum x^2}{n} - \left(\frac{\sum x}{n} \right)^2$$

$$9 = \frac{\sum x^2}{100} - (40)^2$$

$$\frac{\sum x^2}{100} = 1609$$

$$\Rightarrow \sum x^2 = 160900$$

5. Variance of first 20 natural numbers is

- (1) 32.25 (2) 44.25 (3) 33.25 (4) 30

Ans : (3)**Hint :**

Variance for first 20 natural numbers

$$\begin{aligned} & \frac{n^2 - 1}{12} \\ &= \frac{400 - 1}{12} \\ &= \frac{399}{12} \\ &= 33.25 \end{aligned}$$

6. The standard deviation of a data is 3. If each value is multiplied by 5 then the new variance is

- (1) 3 (2) 15 (3) 5 (4) 225

Ans : (4)**Hint :** $\sigma = 3$ of a data.

If each value is multiplied by 5,

then the new SD = 15

$$\therefore \text{Variance} = (\text{SD})^2$$

$$= 15^2$$

$$= 225$$

7. If the standard deviation of x, y, z is p then the standard deviation of $3x + 5, 3y + 5, 3z + 5$ is

- (1)
- $3p + 5$
- (2)
- $3p$
- (3)
- $p + 5$
- (4)
- $9p + 15$

Ans : (2)**Hint :**SD of $x, y, z = p$

$$\Rightarrow \text{SD of } 3x, 3y, 3z = 3p$$

$$\Rightarrow \text{SD of } 3x + 5, 3y + 5, 3z + 5 = 3p.$$

8. If the mean and coefficient of variation of a data are 4 and 87.5% then the standard deviation is

- (1) 3.5 (2) 3 (3) 4.5 (4) 2.5

Ans : (1)**Hint :**

$$\bar{x} = 4, CV = 87.5, \sigma = ?$$

$$CV = \frac{\sigma}{\bar{x}} \times 100$$

$$87.5 = \frac{\sigma}{4} \times 100$$

$$\begin{aligned} \therefore \sigma &= \frac{87.5}{25} \\ &= 3.5 \end{aligned}$$

9. Which of the following is incorrect?

- (1) $P(A) > 1$ (2) $0 \leq P(A) \leq 1$
 (3) $P(\phi) = 0$ (4) $P(A) + P(\bar{A}) = 1$

Ans : (1)**Hint :** $P(A) > 1$ is incorrect.since $0 \leq P(A) \leq 1$

10. The probability a red marble selected at random from a jar containing p red, q blue and r green marbles is

$$(1) \frac{q}{p+q+r} \quad (2) \frac{p}{p+q+r}$$

$$(3) \frac{p+q}{p+q+r} \quad (4) \frac{p+r}{p+q+r}$$

Ans : (2)

Hint :

$$n(\text{Red}) = p, n(S) = p + q + r$$

$$\text{Required probability} = \frac{p}{p+q+r}$$

11. A page is selected at random from a book. The probability that the digit at units place of the page number chosen is less than 7 is

$$\frac{3}{10} \quad \frac{7}{10} \quad \frac{3}{9} \quad \frac{7}{9}$$

Hint :

Ans : (2)

$$P(\text{digit at unit's place of the page is less than 7}) = \frac{7}{10}$$

$$(\because n(S) = 10, A = \{0, 1, 2, 3, 4, 5, 6\}, n(A) = 7)$$

12. The probability of getting a job for a person is $\frac{x}{3}$. If the probability of not getting the job is $\frac{2}{3}$ then the value of x is

$$(1) 2 \quad (2) 1 \quad (3) 3 \quad (4) 1.5$$

Hint :

Ans : (2)

$$\text{Given } P(A) = \frac{x}{3}, P(\bar{A}) = \frac{2}{3}$$

$$P(A) + P(\bar{A}) = 1$$

$$\Rightarrow \frac{x+2}{3} = 1$$

$$\Rightarrow x+2 = 3$$

$$\Rightarrow x = 1$$

13. Kamalam went to play a lucky draw contest. 135 tickets of the lucky draw were sold. If the probability of Kamalam winning is $\frac{1}{9}$, then the number of tickets bought by Kamalam is

$$(1) 5 \quad (2) 10 \quad (3) 15 \quad (4) 20$$

Hint :

Ans : (3)

$$n(S) = 135 \quad n(A) = x$$

$$\therefore P(A) = \frac{x}{135} = \frac{1}{9} \text{ (given)}$$

$$\Rightarrow x = \frac{135}{9} = 15$$

14. If a letter is chosen at random from the English alphabets $\{a, b, \dots, z\}$, then the probability that the letter chosen precedes x

$$(1) \frac{12}{13} \quad (2) \frac{1}{13} \quad (3) \frac{23}{26} \quad (4) \frac{3}{26}$$

Hint :

Ans : (3)

$$n(S) = 26 \quad n(A) = 23 \quad (\because 26 - 3)$$

$$P(A) = \frac{23}{26}$$

15. A purse contains 10 notes of ₹2000, 15 notes of ₹500, and 25 notes of ₹200. One note is drawn at random. What is the probability that the note is either a ₹500 note or ₹200 note?

$$(1) \frac{1}{5} \quad (2) \frac{3}{10} \quad (3) \frac{2}{3} \quad (4) \frac{4}{5}$$

Hint :

Ans : (4)

$$n(S) = 50, n(A) = 10, n(B) = 15, n(C) = 25$$

$$P(B \cup C) = P(B) + P(C) \quad (\because B \text{ \& } C \text{ are mutually exclusive})$$

$$= \frac{15}{50} + \frac{25}{50}$$

$$= \frac{40}{50}$$

$$= \frac{4}{5}$$

UNIT EXERCISE - 8

1. The mean of the following frequency distribution is 62.8 and the sum of all frequencies is 50. Compute the missing frequencies f_1 and f_2 .

Class Interval	0-20	20-40	40-60	60-80	80-100	100-120
Frequency	5	f_1	10	f_2	7	8

Solution :

$$\text{Given } \bar{x} = 62.8, \quad \Sigma f = 50$$

$$\Rightarrow f_1 + f_2 + 30 = 50$$

$$\Rightarrow f_1 + f_2 = 20$$

$$\Rightarrow f_2 = 20 - f_1$$

C.I.	x	f	$d = \frac{x-70}{20}$	fd
0-20	10	5	-3	-15
20-40	30	f_1	-2	$-2f_1$
40-60	50	10	-1	-10
60-80	70	$20-f_1$	0	0
80-100	90	7	1	7
100-120	110	8	2	16
		50		$-2f_1-2$

$$\bar{x} = A + \left(\frac{\Sigma fd}{\Sigma f} \times c \right)$$

$$62.8 = 70 + \left(\frac{-2f_1 - 2}{50} \times 20 \right)$$

$$\Rightarrow 62.8 = 70 + \left(\frac{-4f_1 - 4}{5} \right)$$

$$\Rightarrow 314 = 350 - 4f_1 - 4 \Rightarrow -4f_1 = -32$$

$$f_1 = \frac{32}{4} = 8$$

$$\therefore f_1 = 8, \quad f_2 = 20 - f_1$$

$$= 20 - 8 = 12.$$

2. The diameter of circles (in mm) drawn in a design are given below.

Diameters	33-36	37-40	41-44	45-48	49-52
Number of circles	15	17	21	22	25

Calculate the standard deviation.

Solution :

C.I.	x	f	$d = \frac{x-42.5}{4}$	d^2	$f.d$	$f.d^2$
32.5-36.5	34.5	15	-2	4	-30	60
36.5-40.5	38.5	17	-1	1	-17	17
40.5-44.5	42.5	21	0	0	0	0
44.5-48.5	46.5	22	1	1	22	22
48.5-52.5	50.5	25	2	4	50	100
		100			25	199

2

$$\begin{aligned} \therefore \sigma &= \sqrt{\frac{\Sigma fd^2}{\Sigma f} - \left(\frac{\Sigma fd}{\Sigma f} \right)^2} \times 4 \\ &= \sqrt{\frac{199}{100} - \left(\frac{25}{100} \right)^2} \times 4 \\ &= \sqrt{\frac{19900 - 625}{100^2}} \times 4 \\ &= \frac{\sqrt{19275}}{100} \times 4 \\ &= \frac{138.83}{25} \\ &= 5.55 \end{aligned}$$

$$\therefore \text{S.D} = 5.55$$

3. The frequency distribution is given below.

x	k	$2k$	$3k$	$4k$	$5k$	$6k$
f	2	1	1	1	1	1

In the table, k is a positive integer, has a variance of 160. Determine the value of k .

Solution :

x	f	$d = \frac{x-A}{k}$	d^2	$f.d$	$f.d^2$
k	2	-3	9	-6	18
$2k$	1	-2	4	-2	4
$3k$	1	-1	1	-1	1
$4k$	1	0	0	0	0
$5k$	1	1	1	1	1
$6k$	1	2	4	2	4
	7			-6	28

Given variance = 160

$$\therefore k^2 \left(\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f} \right)^2 \right) = 160$$

$$\Rightarrow k^2 \left[\frac{28}{7} - \left(\frac{-6}{7} \right)^2 \right] = 160$$

$$\Rightarrow k^2 \left[4 - \frac{36}{49} \right] = 160$$

$$\Rightarrow k^2 \left[\frac{160}{49} \right] = 160$$

$$\Rightarrow k^2 = \frac{16 \times 40}{16}$$

$$\Rightarrow k^2 = 49$$

$$\therefore k = 7 \quad (\because k \text{ is positive})$$

4. The standard deviation of some temperature data in degree celsius ($^{\circ}\text{C}$) is 5. If the data were converted into degree Fahrenheit ($^{\circ}\text{F}$) then what is the variance?

Solution : Given $\sigma_c = 5$

$$F = \frac{9c}{5} + 32$$

$$\Rightarrow \sigma_F = \frac{9}{5} \sigma_c$$

$$= \frac{9}{5} \times 5$$

$$= 9 \quad (\because \text{Add (or) subtract the value to a data won't effect the SD})$$

$$\therefore \sigma_F^2 = 9^2 = 81.$$

5. If for a distribution, $\sum(x - 5) = 3$, $\sum(x - 5)^2 = 43$, and total number of observations is 18, find the mean and standard deviation.

Solution :

$$\text{Given } \sum(x - 5) = 3, \quad \sum(x - 5)^2 = 43, \quad n = 18$$

$$\Rightarrow \sum x - \sum 5 = 3 \quad \Rightarrow \sum(x^2 - 10x + 25) = 43$$

$$\Rightarrow \sum x - 5 \cdot \sum 1 = 3 \quad \Rightarrow \sum x^2 - 10 \cdot \sum x + 25 \sum 1 = 43$$

$$\Rightarrow \sum x - 5(18) = 3 \quad \Rightarrow \sum x^2 - 10(93) + 25(18) = 43$$

$$\Rightarrow \sum x = 93 \quad \Rightarrow \sum x^2 = 523$$

i) Mean :

$$\bar{x} = \frac{\sum x}{n} = \frac{93}{18} = 5.17$$

ii) SD :

$$\begin{aligned} \sigma &= \sqrt{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n} \right)^2} = \sqrt{\frac{523}{18} - \left(\frac{93}{18} \right)^2} \\ &= \sqrt{\frac{523}{18} - \frac{8649}{324}} \\ &= \sqrt{\frac{9414 - 8649}{324}} \\ &= \frac{\sqrt{765}}{18} = \frac{27.65}{18} = 1.536 \end{aligned}$$

6. Prices of peanut packets in various places of two cities are given below. In which city, prices were more stable?

Prices in City A	20	22	19	23	16
Prices in city B	10	20	18	12	15

Solution :**CV for prices in City A**

Given data is 20, 22, 19, 23, 16

$$\therefore \bar{x} = \frac{100}{5} = 20$$

To find σ_1 arrange them in ascending order.

x	$d = x - 20$	d^2
16	-4	16
19	-1	1
20	0	0
22	2	4
23	3	9
	0	30

$$\begin{aligned}\therefore \sigma &= \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \\ &= \sqrt{\frac{30}{5}} \\ &= \sqrt{6} \\ &= 2.44\end{aligned}$$

$$\begin{aligned}\therefore C.V &= \frac{\sigma}{x} \times 100 \\ &= \frac{2.44}{20} \times 100 \\ &= 12.24\end{aligned}$$

CV for prices in City B

Given data is 10, 20, 18, 12, 15

$$\therefore \bar{x} = \frac{75}{5} = 15$$

To find σ arrange them in ascending order.

x	$d = x - 15$	d^2
10	-5	25
12	-3	9
15	0	0
18	3	9
20	5	25
	0	68

$$\begin{aligned}\therefore \sigma &= \sqrt{\frac{\sum d^2}{n} - \left(\frac{\sum d}{n}\right)^2} \\ &= \sqrt{\frac{68}{5}} \\ &= \sqrt{13.6} \\ &= 3.68\end{aligned}$$

$$\begin{aligned}\therefore C.V &= \frac{\sigma}{x} \times 100 \\ &= \frac{3.68}{15} \times 100 \\ &= 24.53\end{aligned}$$

\therefore C.V for price in City A < City B

\therefore Prices are very stable in City A.

7. If the range and coefficient of range of

find the largest and smallest values of the data.

Solution :

Given range = 20, Co. eff. of range = 0.2

$$\Rightarrow L - S = 20 \quad \dots(1)$$

$$\frac{L - S}{L + S} = 0.2$$

$$\Rightarrow \frac{20}{L + S} = 0.2$$

$$\Rightarrow L + S = 100 \quad \dots(2)$$

Solving (1) and (2)

$$L = 60, \quad S = 40$$

8. If two dice are rolled, then find the probability of getting the product of face value 6 or the difference of face values 5.

Solution :

$$\begin{aligned}S &= \{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6) \\ &\quad (2,1), (2,2), (2,3), (2,4), (2,5), (2,6)\end{aligned}$$

(3,1),(3,2),(3,3),(3,4),(3,5),(3,6)
 (4,1),(4,2),(4,3),(4,4),(4,5),(4,6)
 (5,1),(5,2),(5,3),(5,4),(5,5),(5,6)
 (6,1),(6,2),(6,3),(6,4),(6,5),(6,6)}

$$n(S) = 36$$

Let A - Product of face value is 6.

$$A = \{(1, 6), (2, 3), (3, 2), (6, 1)\}$$

$$n(A) = 4$$

$$P(A) = \frac{4}{36}$$

Let B - Difference of face value is 5.

$$B = \{(6, 1)\}$$

$$n(B) = 1$$

$$P(B) = \frac{1}{36}$$

$$n(A \cap B) = 1$$

$$P(A \cap B) = \frac{1}{36}$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{4}{36} + \frac{1}{36} - \frac{1}{36} = \frac{4}{36} = \frac{1}{9}$$

- 9. In a two children family, find the probability that there is at least one girl in a family.**

Solution :

$$S = \{(BB), (BG), (GB), (GG)\}$$

$$n(S) = 4$$

Let A be the event of getting atleast one girl.

$$A = \{(BG), (GB), (GG)\}$$

$$\therefore n(A) = 3$$

$$\therefore P(A) = \frac{3}{4}$$

- 10. A bag contains 5 white and some black balls. If the probability of drawing a black ball from the bag is twice the probability of drawing a white ball then find the number of black balls.**

Solution :

$$\text{Given } n(S) = 5 + x, \quad \begin{array}{l} 5 \text{ white balls} \\ x \text{ black balls} \end{array}$$

By daa given,

$$P(B) = 2 \cdot P(W)$$

$$\Rightarrow \frac{x}{5+x} = 2 \cdot \left(\frac{5}{5+x} \right)$$

$$\Rightarrow x = 10$$

$$\therefore \text{No. of black balls} = 10$$

- 11. The probability that a student will pass the final examination in both English and Tamil is 0.5 and the probability of passing neither is 0.1. If the probability of passing the English examination is 0.75, what is the probability of passing the Tamil examination?**

Solution :

$$\text{Given } P(E \cap T) = 0.5 ; \quad P(\bar{E} \cap \bar{T}) = 0.1$$

$$\& P(E) = 0.75 \quad \Rightarrow P(\overline{E \cup T}) = 0.1$$

$$\Rightarrow P(E \cup T) = 1 - 0.1 = 0.9$$

$$P(E \cup T) = P(E) + P(T) - P(E \cap T)$$

$$0.9 = 0.75 + P(T) - 0.5$$

$$P(T) = 0.9 - 0.25$$

$$= 0.65$$

$$= \frac{65}{100}$$

$$= \frac{13}{20}$$

- 12. The King, Queen and Jack of the suit spade are removed from a deck of 52 cards. One card is selected from the remaining cards. Find the probability of getting (i) a diamond (ii) a queen (iii) a spade (iv) a heart card bearing the number 5.**

Solution :

$$n(S) = 52 - 3 = 49$$

- i) Let A - a diamond card

$$n(A) = 13$$

$$\therefore P(A) = \frac{n(A)}{n(S)} = \frac{13}{49}$$

- ii) Let B - a queen card

$$n(B) = 3 \quad (\text{except spade queen out of 4})$$

$$\frac{3}{49}$$

- iii) Let C - a spade card

$$n(C) = 10 \quad (13 - 3 = 10)$$

$$\therefore P(C) = \frac{10}{49}$$

- iv) Let D - 5 of heart

$$n(D) = 1$$

$$\therefore P(D) = \frac{1}{49}$$

PROBLEMS FOR PRACTICE

1. Find the SD of the data
- i) 45, 60, 62, 60, 50, 65, 58, 68, 44, 48
- ii) 8, 10, 15, 20, 22
- iii) 18, 11, 10, 13, 17, 20, 12, 19

(Ans: (i) 8.14, (ii) 5.44, (iii) 3.67)

2. Find the variance of the wages :
Rs.210, Rs.190, Rs.220, Rs.180, Rs.200,
Rs. 190, Rs.200, Rs.210, Rs.180
(Ans : 172.8)
3. Find the range of the heights of 12 girls in a class given in cm.
120, 110, 150, 100, 130, 145, 150, 100, 140, 150, 135, 125
(Ans : 50)
4. The variance of 5 values is 36. If each value is doubled, find the SD of new values.
(Ans : 12)
5. For a group of 200 students, the mean and SD of scores were found to be 40 and 15
scores 43, 35 were misread as 34, 53 respectively. Find the correct mean, SD.
(Ans : 30.955, 14.995)
6. Mean of 100 items is 48 and their S.D is 10. Find the sum of all the items and the sum of the squares of all the items.
(Ans : 4800, 240400)
7. If the coefficient of variation of a collection of data is 57 and its SD is 6.84, find the mean.
(Ans : 12)

8. Calculate S.D from the data :

Marks :	10	20	30	40	50	60
No. of students :	8	12	20	10	7	3

(Ans : 13.45)

9. Find the SD for the data.

Age (in years) : 18 22 21 23 19

No. of students: 100 120 140 150 80

(Ans : 1.84)

10. The following table gives the distribution of income of 100 families in a village. Find the variance.

Income : 0-1000 1000-2000 2000-3000

No. of families : 18 26 30

3000-4000 4000-5000 5000-6000

12 10 4

(Ans : 1827600)

11. Find the coefficient of variation :

(Ans : 20.412)

12. Find the coefficient of variation of the data

Size (in cms) : 10-15 15-20 20-25

No. of items : 2 8 20

25-30 30-35 35-40

35 20 15

(Ans : 21.86)

13. Which of the following cricketers A or B is more consistent player, who scored runs in a cricket season.

A : 58 59 60 54 65 66 52 75 69 52

B : 84 56 92 65 86 78 44 54 78 68

(Ans : Player 'A')

14. Find the missing frequencies of the distribution whose mean is 28.2.

C.V 0-10 10-20 20-30 30-40 40-50

f : 5 f_1 15 f_2 6

(Ans : $f_1 = 8, f_2 = 16$)

15. A number is selected at random from 1 to 100. Find the probability that it is a perfect cube.

(Ans : 1/25)

16. A two digit number is formed of the digits 2, 5 and 9. Find the probability that it is divisible by 2 (or) 5, without repetition.

(Ans : 2/3)

17. From a set of whole numbers less than 40, find the probability of getting a number not divisible by 5 or 7.

(Ans : 12/41)

18. Two dices are thrown together. What is the probability that only odd numbers turn upon both the dices.

(Ans : 5/6)

19. What is the probability that a leap year to contain 53 sundays ?

(Ans : 2/7)

20. The probability that A, B and C can solve a problem are $4/5$, $2/3$, $3/7$ respectively. The probability of the problem being solved by A and B is $8/15$, B and C is $2/7$, A and C is $12/35$. The probability of the problem being solved by all the 3 is $8/35$. Find the probability that the problem can be solved by atleast one of them.

(Ans : 101/105)

OBJECTIVE TYPE QUESTIONS

1. The range of first 20 whole numbers is
a) 19 b) 38 c) 20 d) 19.5

Ans : (c)

2. Variance of 1, 2, 3 is
a) $\frac{2}{3}$ b) 2 c) 0 d) $\frac{\sqrt{2}}{3}$

Ans : (a)

3. The sum of the squares deviations for 10 observations taken from their mean 50 is 250. The coefficient of variation is
a) 10% b) 40% c) 50% d) 15%

Ans : (a)

4. If A and B are mutually exclusive and S is the sample space such that $P(A) = \frac{1}{3}$, $P(B)$ and $S = A \cup B$, their $P(A)$ is
a) $\frac{1}{4}$ b) $\frac{1}{2}$ c) $\frac{4}{3}$ d) $\frac{3}{2}$

5. If the first 10 positive integers, in which we multiply each no. by -1 and then add 1 to each, the variance of the numbers so obtained is
a) 8.25 b) 6.5 c) 3.87 d) 8.25

Ans : (c)

6. The variance of 15 observations is 4. If each observation is increased by 9, the variance of the new data is
a) 13 b) 36 c) 4 d) 16

Ans : (c)

7. Consider the numbers from 1 to 10. If 1 is added to each number, the variance of the numbers so obtained is
a) 6.5 b) 2.87 c) 3.87 d) 8.25

Ans : (c)

8. The probability of drawing neither an ace nor a king is
a) $\frac{2}{13}$ b) $\frac{11}{13}$ c) $\frac{4}{13}$ d) $\frac{8}{13}$

Ans : (b)

9. A number is chosen from 40 to 75. Find the prob. that it is divisible by 7 and 11.

a) $\frac{2}{9}$ b) $\frac{1}{9}$ c) $\frac{3}{9}$ d) $\frac{4}{9}$ **Ans : (a)**

10. The probability of selecting a rotten apple randomly from a heap of 900 apples is 0.18. The number of rotten apples is

a) 0.872 b) 1620 c) 162 d) 172

Ans : (c)

11. In a family of 3 children, probability of having atleast one boy is

a) $\frac{1}{3}$ b) $\frac{7}{8}$ c) $\frac{3}{8}$ d) $\frac{1}{2}$ **Ans : (b)**

12. If a card is drawn at random from 30 cards, the probability that the number on the card is not divisible by 3 is

a) $\frac{2}{3}$ b) $\frac{1}{3}$ c) $\frac{27}{30}$ d) none**Ans : (a)**

13. 3 digit numbers are made using the digits 4, 5, 9 without repetition. If a number is selected at random, the prob. that the number will be ended with 9 is

a) $\frac{1}{3}$ b) $\frac{5}{9}$ c) $\frac{1}{2}$ d) none**(Ans : (a))**

14. A letter of english alphabet is chosen at random. The prob. that the letter chosen is a consonant.

a) $\frac{5}{26}$ b) $\frac{21}{26}$ c) $\frac{7}{13}$ d) 1**Ans : (b)**

15. To prob. of a card to be a club card when it is taken from 52 cards where all red face cards are removed is

a) $\frac{3}{23}$ b) $\frac{13}{46}$ c) $\frac{10}{23}$ d) $\frac{13}{40}$ **Ans : (b)**